

# *Acta Medica Okayama*

---

*Volume 47, Issue 4*

1993

*Article 9*

AUGUST 1993

---

## Cellular immunocompetence in aortitis syndrome.

Akihiro Hosotani\*

Hatsuzo Uchida†

Shigeru Teramoto‡

\*Okayama University,

†Okayama University,

‡Okayama University,

# Cellular immunocompetence in aortitis syndrome.\*

Akihiro Hosotani, Hatsuzo Uchida, and Shigeru Teramoto

## Abstract

Cellular immunocompetence was investigated in 17 cases of aortitis syndrome (3 active, 14 inactive stage). Both the active and inactive groups demonstrated significantly lower interleukin-2 (IL-2) production than healthy volunteers. The active aortitis syndrome group produced significantly more interleukin-1 beta (IL-1 beta) than the inactive group. The proportion of CD11b+ CD8+ cells was significantly lower in the active aortitis syndrome group. Further, the proportions of CD11b- CD8+ cells and CD57+ CD16- cells in the aortitis syndrome patients were significantly higher than the healthy volunteers. These results suggest that there are intrinsic qualitative abnormalities in the T cells that produce IL-2 in aortitis syndrome. Pathogenesis of aortitis syndrome is considered as follows: during the active stage, diminished IL-2 production impairs differentiation and proliferation of suppressor T cells, thus creating abnormalities in the inhibitory functions of immunoregulation and promoting the proliferation of cytotoxic T and natural killer (NK) cells. This presumably initiates inflammation of the aorta and/or artery.

**KEYWORDS:** aortitis syndrome, immunocompetence, interleukin-2, interleukin-1?, lymphocyte subsets

---

\*PMID: 7692704 [PubMed - indexed for MEDLINE]

Copyright (C) OKAYAMA UNIVERSITY MEDICAL SCHOOL

## Cellular Immunocompetence in Aortitis Syndrome

Akihiro Hosotani\*, Hatsuzo Uchida<sup>a</sup> and Shigeru Teramoto

*Second Department of Surgery, Okayama University Medical School and <sup>a</sup>Department of Cardiovascular Surgery, Okayama University Medical School, Okayama 700, Japan*

Cellular immunocompetence was investigated in 17 cases of aortitis syndrome (3 active, 14 inactive stage). Both the active and inactive groups demonstrated significantly lower interleukin-2 (IL-2) production than healthy volunteers. The active aortitis syndrome group produced significantly more interleukin-1 $\beta$  (IL-1 $\beta$ ) than the inactive group. The proportion of CD11b<sup>+</sup> CD8<sup>+</sup> cells was significantly lower in the active aortitis syndrome group. Further, the proportions of CD11b<sup>-</sup> CD8<sup>+</sup> cells and CD57<sup>+</sup> CD16<sup>-</sup> cells in the aortitis syndrome patients were significantly higher than the healthy volunteers. These results suggest that there are intrinsic qualitative abnormalities in the T cells that produce IL-2 in aortitis syndrome. Pathogenesis of aortitis syndrome is considered as follows: during the active stage, diminished IL-2 production impairs differentiation and proliferation of suppressor T cells, thus creating abnormalities in the inhibitory functions of immunoregulation and promoting the proliferation of cytotoxic T and natural killer (NK) cells. This presumably initiates inflammation of the aorta and/or artery.

**Key words :** aortitis syndrome, immunocompetence, interleukin-2, interleukin-1 $\beta$ , lymphocyte subsets

Aortitis syndrome, a generic term, refers to a nonspecific chronic inflammation of unknown etiology that occurs in the aortic arch or its major branches, and may extend to the thoracic, abdominal aorta, or to the pulmonary artery in some cases. As regards the etiology, immunologic abnormalities, particularly diminished cellular immunocompetence, have been addressed (1-3). In the present study, the authors focussed on interleukin-2 (IL-2) and interleukin-1 $\beta$  (IL-1 $\beta$ ), which are important for the regulation and expression of cellular immunity (4, 5), and on the lymphocyte subsets that constitute indices of cellular immunity functions. IL-2 and IL-1 $\beta$  production, plasma concentrations of IL-1 $\beta$ , and lymphocyte subsets were measured in the peripheral blood from patients with aortitis syndrome, and compared with corresponding data from healthy volunteers.

### Materials and Methods

**Subjects.** The present study included 17 cases of aortitis syndrome diagnosed by overall assessment of clinical symptoms, laboratory test results, angiographic and histological findings. The two men and 15 women were aged from 30 to 66 years (mean 46.7 years). Those patients presenting with a 1-h erythrocyte sedimentation rate (ESR) of more than 40 mm associated with a persistent positive reaction for C-reactive protein (CRP) were classified in the active group. Patients with a 1-h ESR less than 40 mm and a negative CRP reaction were classified in the inactive group. One out of the 3 cases in the active group, and 6 of the 14 cases in the inactive group were receiving steroid medication.

Sixteen healthy volunteers were included as controls. All of the them were women, aged from 32 to 66 years (mean 48.4 years).

**Measurement of IL-2 production.** Lymphocytes were separated from heparinized peripheral blood by density gradient centrifugation, and resuspended in culture medium (RPMI-1640 containing 5 % fetal bovine serum (FBS)) to achieve a lymphocyte count of  $2 \times 10^6$  cells/ml. Then, 0.5 ml of the lymphocyte suspension plus 0.5 ml of concanavalin A (ConA) were dispensed to each

\* To whom correspondence should be addressed.

well of a 24-well multiplate, thereby adjusting the lymphocyte count to  $10^6$  cells/well and the ConA concentration to  $5.0 \mu\text{g/ml}$ . The plates were incubated at  $37^\circ\text{C}$  in a 5 carbon dioxide atmosphere for 21h. Then, the supernatant was collected and the IL-2 concentration measured by a radioimmunoassay of two-antibodies method using an Amersham Interleukin-2 [ $I^{125}$ ] Assay Kit.

**Measurement of IL-1 $\beta$  production.** Mononuclear cells (MNC) were separated from heparinized peripheral blood by density gradient centrifugation, and resuspended in culture medium (RPMI-1640 containing 10% FBS) to achieve an MNC count of  $1 \times 10^6$  cells/ml. Aliquots of 0.5ml of the MNC suspension plus 0.5ml of lipopolysaccharide (LPS) were dispensed to each well of a 24-well multiplate, and the LPS concentration adjusted to  $10 \mu\text{g/ml}$ . The plates were incubated at  $37^\circ\text{C}$  in a 5 carbon dioxide atmosphere for 24h. Then, the supernatant was collected and the IL-1 $\beta$  concentration measured by ELISA using a Cistron IL-1 $\beta$  ELISA Kit. The detection limit of this kit is 0.5ng/ml.

**Assay of plasma IL-1 $\beta$  concentration.** Plasma was obtained from peripheral blood samples by centrifugation, and the IL-1 $\beta$  concentration was directly measured by ELISA.

**Determination of lymphocyte subsets.** Fluorescein isothiocyanate (FITC) labelled monoclonal antibody and phycoerythrin (PE) labelled monoclonal antibody  $50 \mu\text{l}$  each were added to  $100 \mu\text{l}$  of the heparinized peripheral blood samples, and the mixture was incubated at  $4^\circ\text{C}$  for 30min. After reaction, 3ml of an erythrocytolytic agent was added, and the mixture was standstill for 15 min at room temperature. Phosphate buffered saline (PBS) was added, and the preparation was washed by centrifugation at 1,000 rpm and  $4^\circ\text{C}$  for 10min. One ml PBS was added to resuspend the pellet, and the lymphocytes subsets were counted with a flow cytometer (FCM-1D, Nippon Bunko).

The monoclonal antibodies used in the measurements comprised FITC labelled Leu7 (CD57) and Leu4 (CD3) antibodies, PE labelled Leu2a (CD8), Leu11c (CD16) and HLA-DR antibodies (Becton & Dickinson), FITC labelled T4 (CD4), Mo1 (CD11b) and PE labelled 2H4 (CD45RA) antibodies (Coulter), and FITC labelled NUT2 (CD25) antibody (Nichirei). The antibody combinations employed in two-color analysis were CD4 plus CD45RA, CD11b plus CD8, CD57 plus CD16, CD3 plus HLA-DR, CD4 plus HLA-DR, and CD8 plus HLA-DR. CD25 antibodies were employed for single-color analysis.

**Statistical analysis.** The results were expressed as mean  $\pm$  standard deviation, and differences between the two groups were assessed by Student's *t*-test. Difference was considered statistically significant when the P value was level less than 5 percent.

## Results

The IL-2 production of the aortitis syndrome patients ( $192.1 \pm 100.8 \text{ pg/ml}$ ) was significantly lower ( $p < 0.01$ ) than the controls' ( $390.5 \pm 144.9 \text{ pg/ml}$ ) (Fig. 1). Regarding, active *versus* inactive status, the IL-2 production of

both active ( $191.7 \pm 86.2 \text{ pg/ml}$ ) and inactive ( $192.2 \pm 106.6 \text{ pg/ml}$ ) groups was significantly lower ( $p < 0.05$  and  $p < 0.01$ , respectively) than the controls' (Fig. 2). The




Fig. 1 IL-2 production in aortitis syndrome patients and healthy subjects.




Fig. 2 IL-2 production in active and inactive stages of aortitis syndrome.

IL-2 production for both steroid users  $138.4 \pm 73.0$  pg/ml and non-users  $229.7 \pm 103.4$  pg/ml was significantly lower ( $p < 0.01$ ) than the controls'.

IL-1 $\beta$  production among the aortitis syndrome patients ( $12.0 \pm 8.7$  ng/ml) tended to be higher than the corresponding level among the healthy subjects ( $10.3 \pm 4.1$  ng/ml), but this difference was not statistically significant (Fig. 3). However, the IL-1 $\beta$  production in the active group ( $23.7 \pm 10.4$  ng/ml) was significantly higher ( $p < 0.01$ ) than the inactive group ( $9.5 \pm 6.2$  ng/ml) (Fig. 4). Neither the active nor the inactive group showed a significant difference related to the healthy group. Further, the IL-1 $\beta$  production in both the steroid-treated group ( $12.1 \pm 9.5$  ng/ml) and the non-steroid group ( $11.9 \pm 8.7$  ng/ml) was not significantly different from that of the healthy subjects. No significant difference between the steroid and non-steroid groups was observed.

Plasma IL-1 $\beta$  level was below the detection sensitivity of the kit in all 33 cases assessed in this series.

The proportion of CD4<sup>+</sup> CD45RA<sup>+</sup> cells was  $13.4 \pm 4.5$  % among the aortitis syndrome patients and  $14.0 \pm 6.8$  % among the healthy subjects, but this difference was not statistically significant. Also, the proportions in both the active group ( $13.7 \pm 6.6$  %) and the inactive group ( $13.4 \pm 4.3$  %) were not significantly different from that of the healthy group. The proportion of CD4<sup>+</sup> CD45RA<sup>-</sup> cells was  $30.5 \pm 6.0$  % for the aortitis syndrome patients and  $31.4 \pm 6.0$  % for the healthy subjects, showing no significant difference. The proportion was higher for the active group ( $34.3 \pm 3.5$  %) than for the inactive group ( $29.7 \pm 6.2$  %), but neither group differed significantly from the healthy group.

The proportion of CD11b<sup>+</sup> CD8<sup>+</sup> cells for the aortitis syndrome patients ( $6.3 \pm 4.6$  %) did not differ significantly from the healthy subjects' ( $6.1 \pm 2.8$  %). However, the proportion in the active group ( $2.3 \pm 1.5$  %) was significantly lower ( $p < 0.05$ ) than the healthy subjects (Fig. 5). CD11b<sup>-</sup> CD8<sup>+</sup> cell proportions demonstrated no significant difference between the active aortitis syndrome group ( $24.8 \pm 7.8$  %) and the inactive group ( $23.1 \pm 7.5$  %). However, the proportion for the overall aortitis syndrome patients ( $23.4 \pm 7.3$  %) was significantly higher ( $p < 0.05$ ) than the healthy subjects' ( $19.0 \pm 4.4$  %) (Fig. 6).

The proportion of CD57<sup>+</sup> CD16<sup>+</sup> cells was  $8.9 \pm 5.6$  % for the aortitis syndrome patients and  $11.5 \pm 5.9$  % for the healthy subjects. Moreover, neither the correspond-




Fig. 3 IL-1 $\beta$  production in aortitis syndrome patients and healthy subjects.




Fig. 4 IL-1 $\beta$  production in active and inactive stages of aortitis syndrome.

ing values for the active group ( $7.3 \pm 2.7\%$ ) nor those for the inactive group ( $9.2 \pm 6.1\%$ ) had significant differences from those of controls. The proportion of  $CD57^+ CD16^-$  cells was significantly increased ( $p < 0.05$ ) among the aortitis syndrome patients ( $13.4 \pm 6.9\%$ ) as compared with the healthy subjects ( $8.8 \pm 4.0\%$ ) (Fig. 7). However, there was no significant difference in this aspect between the active group ( $10.6 \pm 3.7\%$ ) and the inactive group ( $13.9 \pm 7.4\%$ ). Regarding the proportion of  $CD57^- CD16^+$  cells, the values for the aortitis syndrome patients ( $4.4 \pm 1.8\%$ ) and the healthy subjects ( $4.7 \pm 1.5$

$\%$ ) were not significantly different, but those for the active group ( $2.1 \pm 0.5\%$ ) were significantly lower ( $p < 0.05$ ) than the inactive group' ( $4.8 \pm 1.6\%$ ).

The proportions of  $CD3^+ HLA-DR^+$  cells among the aortitis syndrome patients ( $6.2 \pm 2.7\%$ ) and the healthy subjects ( $6.2 \pm 3.6\%$ ) were not significantly different, also the corresponding proportions in the active group ( $4.7 \pm 3.0\%$ ) and the inactive group ( $6.5 \pm 2.7\%$ ) were not significantly different from those of the healthy subjects. Likewise, the proportion of  $CD4^+ HLA-DR^+$  cells in the aortitis syndrome patients ( $2.1 \pm 0.9\%$ ) was not




Fig. 5 Proportion of  $CD11b^+ CD8^+$  cells in active and inactive stages of aortitis syndrome and in healthy subjects.




Fig. 6 Proportion of  $CD11b^- CD8^+$  cells in active and inactive stages of aortitis syndrome and in healthy subjects.




Fig. 7 Proportion of CD57<sup>+</sup> CD16<sup>+</sup> cells in active and inactive stages of aortitis syndrome and in healthy subjects.

significantly different from that in the healthy subjects ( $2.5 \pm 1.2\%$ ), and neither the proportion for the active group ( $2.0 \pm 1.3\%$ ) nor that for the inactive group ( $2.2 \pm 0.9\%$ ) showed a statistically significant difference from that for the healthy subjects. The proportion of CD8<sup>+</sup> HLA-DR<sup>+</sup> cells observed in the aortitis syndrome patients ( $3.8 \pm 2.0\%$ ) showed no significant difference from the healthy subjects' ( $3.8 \pm 2.8\%$ ). However, the proportions in the active group ( $2.6 \pm 1.9\%$ ) tended to be lower than those in the inactive group ( $4.0 \pm 2.0\%$ ). Finally, the proportions of CD25<sup>+</sup> cells among the aortitis syndrome patients ( $1.8 \pm 0.9\%$ ) and those among the healthy subjects ( $1.9 \pm 1.4\%$ ) were not significantly different, and there was no significant difference between the proportions in the active group ( $1.7 \pm 1.0\%$ ) and the inactive group ( $1.9 \pm 0.9\%$ ).

## Discussion

The aortitis syndrome is characterized by obstructive lesions due to nonspecific chronic inflammation at the aorta or at its major branching sites, and the etiology of this syndrome still remains obscure. More recently, attention has been paid in relation to the etiology of this syndrome. Among the causes, infection including tuberculosis, genetic factors, and hormonal abnormalities (6)

have been pointed out. Since Judge *et al.* (7) asserted that autoimmunity is involved in the pathogenesis of this syndrome, many reports have appeared in this connection, supporting the idea of immunological mechanisms to play a role in this disorder. Particularly, studies on cellular immunity have revealed that patients in the active stage of the aortitis syndrome present characteristics of decreased response in delayed skin reactions, such as the tuberculin reaction and the dinitrochlorobenzene (DNCB) skin reaction (1), as well as decreased reactivity of peripheral blood lymphocytes with respect to the phytohemagglutinin (PHA) induced blast transformation (2). Many other forms of depressed cellular immunocompetence; decreased proportions of T cells among the peripheral blood lymphocytes and decreased PHA-induced lymphoblast transformation rates, reportedly occur in aortitis syndrome patients in the active and inactive stages (3).

IL-2 manifests important actions; promoting proliferation of T cells, activating cytotoxic T cells, inducing differentiation of B cells, activating NK cells, inducing lymphokine-activated killer (LAK) cells and inducing formation of IL-2 receptors, and thereby plays a key role in cellular immunoregulation (4). Alcocer-Varela *et al.* (8) have reported that, compared to healthy individuals, the IL-2 production of aortitis syndrome patients in both the active and inactive stages is depressed, and in particular is significantly depressed in the active stage. In the present

study also, although no significant differences between the active and inactive groups were observed, both groups had significantly lower IL-2 production than the healthy subjects.

HLA-DR<sup>+</sup> cells are regarded as activated T cells (9). The results of the present study revealed that the proportions of CD3<sup>+</sup> HLA-DR<sup>+</sup> cells, CD4<sup>+</sup> HLA-DR<sup>+</sup> cells, and CD8<sup>+</sup> HLA-DR<sup>+</sup> cells in both active and inactive stages of aortitis syndrome were not significantly different from the corresponding proportions in healthy subjects, indicating that the proportion of activated T cells is not particularly elevated even in active stages. Hence, the conjecture that *in vitro* decreases in IL-2 production are due to prior *in vivo* activation and consequent exhaustion of the IL-2 production of the T cells before *in vitro* cultivation appears implausible.

CD4<sup>+</sup> cells are divided into the two subsets of helper and suppressor-inducer T cells on the basis of reactivity with CD45RA antibody (10). Of these two categories, the helper T cells are considered to be primarily responsible for IL-2 production (10). Alcocer-Varela *et al.* showed that both the proportion and absolute number of CD4<sup>+</sup> cells were lower in aortitis syndrome patients than in the controls (8). However, they did not subclassify CD4<sup>+</sup> cells into helper or suppressor-inducer T cells. Our results for CD4<sup>+</sup> CD45RA<sup>-</sup> cell proportion (helper T cells) was not significantly different, in either the active or inactive aortitis syndrome group, from the healthy subjects'. Hence, an explanation attributing the results to a decrease in the number of IL-2-producing helper T cells also appears implausible.

The subset of CD8<sup>+</sup> cells can be subdivided into CD11b<sup>+</sup> CD8<sup>+</sup> cells, regarded as being suppressor T cells, and CD11b<sup>-</sup> CD8<sup>+</sup> cells, regarded as being cytotoxic T cells (11). However, in the present study, the proportion of CD11b<sup>+</sup> CD8<sup>+</sup> cells, (*i.e.*, putative suppressor T cells) among all the aortitis syndrome patients was not significantly different from the healthy subjects'. Because the proportion of suppressor T cells in the active group was actually lower, the conjecture that IL-2 production diminished through inhibition by suppressor T cells also seems improbable. Further, CD25<sup>+</sup> cells are regarded as low-affinity IL-2 receptors (12), but in our study, there was no significant difference in the proportion of CD25<sup>+</sup> cells among the active, inactive, and healthy group'. This may exclude the conjecture attributing decreased IL-2 production to diminished expression of low-affinity IL-2 receptors. Production of

IL-2 by T-lymphocytes requires IL-1 production by antigen-presenting cells, mainly macrophages (13), but results of this study indicated that IL-1 $\beta$  production was well maintained in both the active and inactive aortitis syndrome groups, which seems to exclude the hypothesis of functional abnormalities in macrophages. Indeed, the fact that diminished IL-2 production was recognized in the active stage and in the inactive stage of aortitis syndrome, even years after the onset of the disease, indicates that, as reported with reference to patients with systemic lupus erythematosus (SLE) (14, 15) and other immunodeficiencies (16), decreased IL-2 production is actually due to qualitative abnormalities. This suggests the possibility of intrinsic defects of T cells in aortitis syndrome. Thus, one may infer that abnormalities in IL-2 production may be the phenomenon involved in the pathogenesis of aortitis syndrome.

IL-1 has  $\alpha$  and  $\beta$  forms, with different isoelectric points, but these two forms are thought to bind to the same receptors and to exhibit similar biological activity (17). IL-1 plays an important role in the acute-phase of the inflammatory response and in immunoregulatory functions; including endogenous pyrogenic activity, synthesis of acute inflammatory proteins such as haptoglobin (18), action upon vascular endothelial cells and fibroblasts to promote proliferation of capillaries and fibroblasts (19). The mononuclear cells in the peripheral blood of patients with Kawasaki's disease, characterized by inflammatory angiitis and similar to the aortitis syndrome, also show increased IL-1 $\beta$  production (20). The circulating level of IL-1 $\beta$  is significantly elevated in patients with Kawasaki's disease (21). IL-1 involvement in another inflammatory disorder has been reported, *e.g.*, strong IL-1 activity has been demonstrated in the synovial fluid and the supernatants of synovial membrane cell cultures obtained from patients with rheumatoid arthritis (22). Our results that IL-1 $\beta$  production in patients with active aortitis syndrome was significantly increased indicate that IL-1 may be somehow involved in the onset and further development of aortic inflammation. In particular, the local production of IL-1 $\beta$  infiltrated macrophages in the severely inflamed aortic media and adventitia may play an important role in the onset, progression, and continuation of aortic wall inflammation. Plasma concentrations of IL-1 $\beta$  were under the limit of detection of the kit in the all patients with aortitis syndrome in the present study. However, this does not exclude the possibility of IL-1 $\beta$  involvement on the pathogenesis of the aortitis syndrome. Since



IL-1 $\beta$  is locally produced, its concentrations in peripheral blood were extremely low, and below the detection sensitivity.

Pronounced infiltration of CD8<sup>+</sup> lymphocytes and cytotoxic T cells into the arterial wall of specimens excised from aortitis syndrome patients has been reported in immunohistological studies (23-25). Similarly, our results showed that the proportion of CD11b<sup>-</sup> CD8<sup>+</sup> cells (11) in the peripheral blood of aortitis syndrome patients was significantly elevated, indicating that cytotoxic T cells play a major role in the onset and progression of aortic inflammation.

NK cells, possessing cytotoxic activity, can be divided into three subtypes according to their reactivity with CD57 and CD16 antibodies; CD57<sup>-</sup> CD16<sup>+</sup> cells (strong activity) CD57<sup>+</sup> CD16<sup>+</sup> cells (intermediate activity), and CD57<sup>+</sup> CD16<sup>-</sup> cells (weak activity) (26). In the aortitis syndrome, the proportions of the cells with strong or intermediate activity, were the same as the controls', but the proportions of CD57<sup>+</sup> CD16<sup>-</sup> cells were significantly elevated. Furthermore, this proportion was higher in the inactive stage than the active stage. These findings may indicate that NK cells, rather than being involved in the development of cytotoxic expression during the acute inflammatory stage, are more closely related to the maintenance of inflammatory phenomena during the inactive stage of aortitis syndrome.

Further, the fact that the proportions of CD11b<sup>+</sup> CD8<sup>+</sup> cells, considered to be suppressor T cells (11), were significantly lower in the active aortitis syndrome group than the healthy subjects indicates the possibility that the decrease in IL-2 production impaired differentiation into and proliferation of suppressor T cells. This could create abnormalities in the inhibitory functions of immunoregulation, promote proliferation of cytotoxic T cells, and cause aortic and/or arterial inflammation through the mediation of IL-1 $\beta$ .

Acknowledgment. This study was partially supported by the Grant-in-Aid from the Ministry of Health and Welfare in Japan.

## References

- Ogino K: Immunologic studies of the aortitis syndrome. *J Jpn Coll Angiol* (1977) **17**, 283-293 (in Japanese).
- Ogino K, Chin K, Fujita K, Itagaki F, Uchida H and Teramoto S: Immunological study in aortitis syndrome; in Annual Report 1976, Vascular Lesion of Collagen Disease Research Committee, Ministry of Health and Welfare, Tokyo, Japan (1977) pp 151-155 (in Japanese).
- Inada K, Miyamoto R and Okada A: Immunological study on chronic occlusive arterial disease -cell-mediated immunity; in Annual Report 1978, Vascular Lesion of Collagen Disease Research Committee, Ministry of Health and Welfare, Tokyo, Japan (1979) pp 331-336 (in Japanese).
- Smith KA: Interleukin-2: Inception, impact, and implications. *Science* (1988) **240**, 1169-1176.
- Dinarelli CA: Interleukin-1 and interleukin-1 antagonism. *Blood* (1991) **77**, 1627-1652.
- Otsuka A: Hormonal study on hypothalamo-hypophyseal function in patients with aortitis syndrome. *J Jpn Coll Angiol* (1982) **22**, 253-264 (in Japanese).
- Judge RD, Currier RD, Gracie WA and Figley MM: Takayasu's arteritis and the aortic arch syndrome. *Am J Med* (1962) **32**, 379-386.
- Alcocer-Varela J, Reyes-Lopez PA, Sanchez-Torres G and Alarcon-Segovia D: Immunologic studies in patients with Takayasu's arteritis: III. Immunoregulatory changes. *Clin Exp Rheumatol* (1989) **7**, 345-350.
- Yu DTY, Winchester RJ, Fu SM, Gibofsky A, Ko HS and Kunkel HG: Peripheral blood Ia-positive T cell. *J Exp Med* (1980) **151**, 91-100.
- Morimoto C, Letvin NL, Distaso JA, Aldrich WR and Schlossman SF: The isolation and characterization of the human suppressor inducer T cell subset. *J Immunol* (1985) **134**, 1508-1515.
- Landay A, Gartland GL and Clement LT: Characterization of a phenotypically distinct subpopulations of Leu-2<sup>+</sup> cells that suppresses T cell proliferative responses. *J Immunol* (1983) **131**, 2757-2761.
- Uchiyama T, Broder S and Waldmann TA: A monoclonal antibody (anti-Tac) reactive with activated and functionally mature human T cells: I. Production of anti-Tac monoclonal antibody and distribution of Tac (+) cells. *J Immunol* (1981) **126**, 1393-1397.
- Kaye J, Gills S, Mizel S, Shevach EM, Malek TR, Dinarello CA, Lachman LB and Janeway CA: Growth of a cloned helper T cell line induced by a monoclonal antibody specific for the antigen receptor: Interleukin 1 is required for the expression of receptor for interleukin 2. *J Immunol* (1984) **133**, 1339-1345.
- Murakawa Y, Takada S, Ueda Y, Suzuki N, Hoshino T and Sakane T: Characterization of T lymphocyte subpopulations responsible for deficient interleukin 2 activity in patients with systemic lupus erythematosus. *J Immunol* (1985) **134**, 187-195.
- Alcocer-Varela J and Alarcon-Segovia D: Decreased production of and response to interleukin-2 by cultured lymphocytes from patients with systemic lupus erythematosus. *J Clin Invest* (1982) **69**, 1388-1392.
- Lopez-Botet, Fontan G, Rodriguez MCG and de Landazuri MO: Relationship between IL 2 synthesis and the proliferative response to PHA in different primary immunodeficiencies. *J Immunol* (1982) **128**, 879-885.
- Kilian PL, Kaffka KL, Stern AS, Woehle D, Benjamin WR, Dechiara TM, Gubler U, Farrar JJ, Mizel SB and Lomedico PT: Interleukin 1 $\alpha$  and interleukin 1 $\beta$  bind to the same receptor on T cell. *J Immunol* (1986) **136**, 4509-4514.
- Ramadori G, Sipe JD, Dinarello CA, Mizel SB and Colten HR: Pretranslational modulation of acute phase hepatic protein synthesis by murine recombinant interleukin 1 (IL-1) and purified human IL-1. *J Exp Med* (1985) **162**, 930-942.
- Oppenheim JJ, Kovacs EJ, Matsushima K and Durum SK: There is more than one interleukin 1. *Immunol Today* (1986) **7**, 45-56
- Leung DYM, Cotran RS, Kurt-Jones E, Burns JC, Newburger JW and Pober JS: Endothelial cell activation and high interleukin-1 secretion in the pathogenesis of acute Kawasaki disease. *Lancet* (1989) **2**,

- 1298-1302.
21. Maury CPJ, Salo E and Pelkonen P: Circulating interleukin-1 $\beta$  in patients with Kawasaki disease. *N Engl J Med* (1988) **319**, 1670-1671.
  22. Miyasaka N, Sato K, Goto M, Sasano M, Natsuyama M, Inoue K and Nishioka K: Augmented interleukin-1 production and HLA-DR expression in the synovium of rheumatoid arthritis. *Arthritis Rheum* (1988) **31**, 480-486.
  23. Scott DGI, Salmon M, Scott DL, Blann A, Bacon PA, Walton KW, Oakland CDH and Slaney GF: Takayasu's arteritis: A pathogenetic role for cytotoxic T lymphocytes? *Clin Rheumatol* (1986) **5**, 517-522.
  24. Seko Y, Yamasaki R, Yazaki Y, Tada Y and Okumura Y: Analysis of the immunological mechanism of vascular cellular impairment in Takayasu's arteritis -comparison with viral myocarditis; in Annual Report 1990, Intractable Angiitis Research Committee, Ministry of Health and Welfare, Tokyo, Japan (1991) pp 14-16 (in Japanese).
  25. Seko Y, Yamasaki R, Yazaki Y, Tada Y, Yagita H and Okumura Y: Analysis of the immunological mechanism of vascular cellular impairment in Takayasu's arteritis; in Annual Report 1991, Intractable Angiitis Research Committee, Ministry of Health and Welfare, Tokyo, Japan (1992) pp 32-34 (in Japanese).
  26. Lanier LL, Le AM, Phillips JH, Warner NL and Babcock GF: Subpopulations of human natural killer cells defined by expression of the Leu-7 (HNK-1) and Leu-11 (NK-15) antigen. *J Immunol* (1983) **131**, 1789-1796.

Received January 25, 1993; accepted March 11, 1993.