

Acta Medica Okayama

Volume 49, Issue 6

1995

Article 1

DECEMBER 1995

Antioxidant activity of Chinese ant extract preparations.

Yi Zhao*	Xiaoyan Wang [†]	Motoko Kawai [‡]
Jiankang Liu**	Miao Liu ^{††}	Akitane Mori ^{‡‡}

*Guangxi College of TCM,

[†]Okayama University,

[‡]Okayama University,

**Okayama University,

^{††}Okayama University,

^{‡‡}Okayama University,

Antioxidant activity of Chinese ant extract preparations.*

Yi Zhao, Xiaoyan Wang, Motoko Kawai, Jiankang Liu, Miao Liu, and Akitane Mori

Abstract

Chinese ant extract preparations (CAEP) are a Chinese traditional medicine which is mainly used as a health food or drink for the treatment of rheumatism, rheumatoid arthritis, chronic hepatitis, sexual hypofunction, and antiaging in China. The effects on free radicals were examined by electron spin resonance spectrometry using the spin trapping agent 5,5-dimethyl-1-pyrroline-1-oxide (DMPO). Superoxide radicals (3.35×10^{15} spins/ml) were quenched 50% by the extract at 0.5 mg/ml. The CAEP extract at 0.7 mg/ml inhibited 50% of hydroxyl radicals (52.0×10^{15} spins/ml) generated by the Fenton reaction. Against DPPH radical, the scavenging action of CAEP was observed at 1.8 mg/ml of the extract and 50% of the DPPH radicals (8.14×10^{15} spins/ml) were quenched. In vitro tests showed that CAEP inhibited the production of thiobarbituric acid-reactive substances, an index of lipid peroxidation, in rat brain homogenate.

KEYWORDS: free radical scavenger, antioxidant, hydroxyl radical, superoxide radical, lipid peroxidation

*PMID: 8770235 [PubMed - indexed for MEDLINE]

Copyright (C) OKAYAMA UNIVERSITY MEDICAL SCHOOL

Antioxidant Activity of Chinese Ant Extract Preparations

Yi ZHAO^a, Xiaoyan WANG^b, Motoko KAWAI^b, Jiankang LIU^b, Miao LIU^c, and Akitane MORI^{b,*}

^aInstitute of Traditional Chinese Medicine (TCM), Guangxi College of TCM, Nanning 530001, China, ^bDepartment of Neuroscience, Institute of Molecular and Cellular Medicine and ^cPublic Health, Okayama University Medical School, Okayama 700, Japan

Chinese ant extract preparations (CAEP) are a Chinese traditional medicine which is mainly used as a health food or drink for the treatment of rheumatism, rheumatoid arthritis, chronic hepatitis, sexual hypofunction, and antiaging in China. The effects on free radicals were examined by electron spin resonance spectrometry using the spin trapping agent 5,5-dimethyl-1-pyrroline-1-oxide (DMPO). Superoxide radicals (3.35×10^{15} spins/ml) were quenched 50% by the extract at 0.5 mg/ml. The CAEP extract at 0.7 mg/ml inhibited 50% of hydroxyl radicals (52.0×10^{15} spins/ml) generated by the Fenton reaction. Against DPPH radical, the scavenging action of CAEP was observed at 1.8 mg/ml of the extract and 50% of the DPPH radicals (8.14×10^{15} spins/ml) were quenched. *In vitro* tests showed that CAEP inhibited the production of thiobarbituric acid-reactive substances, an index of lipid peroxidation, in rat brain homogenate.

Key words; free radical scavenger, antioxidant, hydroxyl radical, superoxide radical, lipid peroxidation

Ants as a health food (drink) or medicine have a long history in China, dating back about 3,000 years. It was first recorded in Zhou-Li, a historical book which talks about the popularity of greetings in China, and ants were mentioned as an excellent food for the king (1). Li Shizhen, a famous pharmacologist during the Ming dynasty, edited Ben-Cao-Gang-Mu a Compendium of Materia medica in which ants were recorded as a food and as a medicine as well (2). At present, there are some areas in China such as Guangxi, where people eat ant powder or drink ant wine to maintain their health. In several other countries, such as Argentina, Burma, Colombia, Mexico, Africa, and America, ants are also used in folk medicine (3).

As a Chinese traditional medicine, Chinese ant extract preparation (CAEP) is mainly used as a health food or drink for the treatment of rheumatism, rheumatoid arthritis, chronic hepatitis, sexual hypofunction, and for antiaging in China (4). Some pharmacological effects and clinical uses of ants were reported (5, 6), but the results were preliminary.

Free radicals have been implicated in many disorders such as epilepsy (7), aging (8), ischemia (9), trauma (10), and rheumatoid arthritis (11), and even in cancer. Among the reactive oxygen species (ROS), hydroxyl radicals are the most reactive and they damage proteins, break DNA, and promote lipid peroxidation (7). Therefore, we investigated CAEP to determine whether or not it has the ability to quench free radicals.

Materials and Methods

CAEP

CAEP was prepared from *Polyrhachis vicina* R., a kind of black ant. Ant was rinsed with ethanol-water several times and powdered. Further ant powder was infused for 24 h with 5 volumes of ethanol 40% (w/v) (500 ml). After filtration with two layer gauzes, ant powder was centrifugated (3,000 rpm, 15 min) and filtrated with quantitative filter paper. After removal of ethanol, the remaining ant extracts water solution was evaporated and concentrated to 50% (Fig. 1). The extract solution (CAEP) prepared by this method was equivalent to 100 mg dried ants/ml.

Chemical

Hypoxanthine (HPX), 1,1-diphenyl-2-picrylhydrazyl (DPPH) and diethylenetriaminepentaacetic acid (DETAPAC) were obtained from Sigma Chemical Co.

* To whom correspondence should be addressed.

^aVisiting researcher of Okayama University Medical School, Okayama 700, Japan




Fig. 1 Method of Chinese ant extract preparations.

(St. Louis, MO); xanthine oxidase (XOD) was from Boehringer Mannheim (Mannheim, Germany); 5,5-dimethyl-1-pyrrolyne-1-oxide (DMPO) was from Labotec Co. (Tokyo, Japan), and ferrous sulfate, hydrogen peroxide and other ordinary reagents were of the highest quality available from Wako Pure Chemical Industries or Katayama Chemical Co. (Osaka, Japan).

Free Radical Analysis

Free radicals were analyzed by electron spin resonance (ESR) spectrometer (JES-FE1XG, JEOL Co., Japan) according to the method as described by Liu *et al.* (8, 9).

DPPH radical analysis. For the DPPH radical analysis, 100 μ l of 50 mM DPPH ethanol solution and 100 ml of CAEP solution were mixed for 5 sec and then transferred to a special flat cell in which the amount of DPPH radical was estimated by ESR spectrometry exactly 40 sec after the addition of CAEP. Signal intensities were evaluated by taking the height of the third peak of the five-peak DPPH radical signal. The ESR settings were as follows: microwave power 8 mW, modulation frequency 100 kHz, modulation width 0.08 mT, scan time 1 min/360 mm, response 0.1 s, magnetic field intensity 335.6 ± 10 mT and amplitude 4.0×10^2 .

Superoxide radical analysis. For the superoxide radical analysis, the following reagents were added in a test tube in the following order: 50 μ l of 2 mM HPX, 35 μ l of 11.0 mM DETAPAC, 50 μ l of the CAEP solution, 50 μ l of XOD (0.326 units/ml) and 15 μ l of 9.2 M DMPO. The solution was mixed and then placed in a

flat ESR cell (volume; 160 μ l) and DMPO-OH spin adduct was analyzed by ESR spectrometry. The ESR settings were as follows: microwave power 8 mW, modulation frequency 100 kHz, modulation width 0.08 mT, scan time 1 min/360 mm, response 0.1 s, magnetic field intensity 335.6 ± 5 mT and amplitude 3.2×10^3 .

Hydroxyl radical analysis. For the hydroxyl radical analysis, 75 μ l of 1 mM FeSO₄ and DETAPAC solution, 50 μ l of the extract solution, 20 μ l of 0.092 M DMPO and 75 μ l of 1 mM hydrogen peroxide were put into a test tube and mixed. The amount of DMPO-OH spin adducts formed was estimated exactly 40 sec after adding the DMPO. The ESR settings were as follows: microwave power 8 mW, modulation frequency 100 kHz, modulation width 0.08 mT, scan time 1 min/360 mm, response 0.1 s, magnetic field intensity 335.6 ± 5 mT and amplitude 1.0×10^2 .

Measurement of Lipid Peroxides

Tissue homogenate was prepared from the brains and livers of 15-week-old male Sprague-Dawley rats. Thiobarbituric acid-reactive substances (TBARS) was measured as described by Ohkawa *et al.* (10) using malondialdehyde as the standard. Briefly, 1.5 ml of 1% thiobarbituric acid (TBA) was added to each sample, followed by 1.5 ml of 20% acetic acid (pH 3.5) and 0.2 ml of sodium dodecyl sulfate. After heating for 60 min at 100 °C, the samples were extracted into 3 ml of n-butanol and measured spectrofluorimetrically.

Statistical Analysis

The statistical significance of the differences was examined by two-way analysis of variance.

Results

The scavenging action of CAEP on DPPH radicals, superoxide radicals and hydroxyl radicals were showed (Fig. 2-A, Fig. 3-A and Fig. 4-A). Against DPPH radical, the scavenging action of CAEP was observed at 1.8 mg/ml of the extract and 50% of the DPPH radicals (8.14×10^{15} spins/ml) were quenched (Fig. 2-B). Superoxide radicals (3.53×10^{15} spins/ml) were quenched 50% by the extract at 0.5 mg/ml (Fig. 3-B). The CAEP extract at 0.7 mg/ml inhibited 50% of hydroxyl radicals (52.0×10^{15} spins/ml) generated by the Fenton reaction (Fig. 4-B).

As shown in Fig. 5, at the concentrations of 0.3 and 0.63 mg/ml of CAEP, there was a decrease in the TBARS level compared to the control (7.26 ± 1.15 ; $n =$




Fig. 2 Effects of Chinese ant extract preparation (CAEP) on DPPH radicals. **A.** ESR spectra obtained in the reaction mixture containing 0–7.8 mg/ml concentrations of the extract. **B.** Dose-dependent effects on DPPH radicals. Each point is the mean \pm SD of 5–6 experiments.




Fig. 3 Effects of Chinese ant extract preparation (CAEP) on superoxide radicals, generated by the hypoxanthine-xanthine oxidase system, measured as DMPO-OOH spin adducts. **A.** ESR spectra obtained in the reaction mixture containing 0–31.25 mg/ml concentrations of the extract. **B.** Dose-dependent effects on superoxide radicals. Each point is the mean \pm SD of 5–6 experiments.




Fig. 4 Effects of Chinese ant extract preparation (CAEP) on hydroxyl radicals, generated by Fenton reagents, measured as DMPO-OH spin adducts.

A. ESR spectra obtained in the reaction mixture containing 0–28.8 mg/ml concentrations of the extract. B. Dose-dependent effects on hydroxyl radicals. Each point is the mean \pm SD of 5–6 experiments.




Fig. 5 TBARS level in brain homogenate incubated with Chinese ant extract preparation (CAEP).

6; $P < 0.001$ vs. control, 6.23 ± 1.55 ; $n = 7$; $P < 0.001$ vs. control)

Discussion

Although the antioxidants, or free radical scavengers, in plants have been studied extensively (11–13), the antioxidants in insects and animals have received little attention (14). In the present study, we studied the free radical scavenging effect of CAEP and found that it is a potent scavenger of superoxide, hydroxyl and DPPH radicals. Furthermore, the IC_{50} of superoxide radical scavenging activity by CAEP was almost the same as of TJ-960, which is a Japanese herbal medicine contained seven herbs (Table 1).

CAEP has been widely used as a detoxifying agent to eliminate swelling and facilitate recovery (17). It has been shown to have beneficial effects on both rheumatism and chronic hepatitis (18–20). Most recently, clinical and animal experiments showed that CAEP can regulate the immune system by activating certain enzymes, stimulating the division of immune cells, and promoting the growth of immune organs, and also that CAEP can strengthen resistance to disease by regulating the physiological equilibrium of the whole immune system (20 and

Table I Concentration of antioxidants required to quench 50% (IC₅₀) of the free radicals

Antioxidant/ scavenger	IC ₅₀ of antioxidant		Reference
	Hydroxyl radicals ($\mu\text{g/ml}$)	Superoxide radicals ($\mu\text{g/ml}$)	
Vitamin C	35	24	15
Vitamin E	37	—	15
TJ-960	—	700	16
Soybean paste miso	4000	5000	15
CAEP	700	500	—

unpublished data). Although the mechanisms are not clearly understood, it seems very likely that the effectiveness of CAEP is at least partly attributable to its free radical scavenging activity. Therefore, the beneficial effects of CAEP on free radical related diseases such as rheumatism, rheumatoid arthritis and chronic hepatitis, may be based, at least in a part, on this mechanism. Chemical analysis revealed that *Polyrhachis vicina* R. (PVR) contains aliphatic hydrocarbons in its stomach, formic acid in its toxic saliva, and more than 17 kinds of free amino acids such as, 2-amino-6-hydroxypteridine and bioppterin (14). Therefore, it is quite interesting and important to further identify the biologically active components in the extract and to study the relationship between their structure and activity. It is also important to identify the mechanism of the synergistic effects, by isolating the pure natural products from the extract and by experimenting with chemical substances of known structure.

References

- Anonymous (1066-256 BC): Courtesy of the Chinese Zhou Dynasty (A Chinese ancient book), Zhou Li ed., Vol 1, Shanghai Commercial Publishing Company, China (1955) p 26 (in Chinese).
- Li Shi-Zhen (AD 1518-1593): Compendium of Materia Medica, Ban Cao Gang Mu eds., Beijing People's Hygiene Publishing House, China (1981) Vol 40 p 2288 (in Chinese).
- Zhao Yi: Review of the ants as food or medicine. J Guangxi TCM (1989) 12, 35-36.
- Zhao Yi and Lian Wei: Pharmacological basis of therapeutics on the Chinese ant preparation; in ICBES Abstract Yamagata, Japan (1994) p 66.
- Zhao Yi: Pharmacological studies on the Chinese ant preparation. J Guangxi TCM (1983) 6, 39-42 (in Chinese).
- Zhao Yi: Review of clinical effect of 16015 cases rheumatoid arthritis treated by Chinese ant (*Polyrhachis vicina* Roger) preparation. J Guangxi TCM (1993) 16, 44-47 (in Chinese).
- Mori A, Hiramatsu M, Yokoi I and Edamatsu R: Biochemical pathogenesis of post-traumatic epilepsy. Pavlovian J Biol Sci (1990) 25, 54-62.
- Harman D: The aging process. Proc Natl Acad Sci USA (1981) 78, 7124-7128.
- Yoshikawa T, Naito Y and Kondo M: Free radical involvement in the aging process. Neurosciences (1990) 16, 603-612.
- Hallenback JM and Dutka J: Background review and current concepts of reperfusion injury. Arch Neurol. (1990) 47, 1245-1254.
- Braugher JM and Hall ED: Central nervous system trauma and stroke: Biochemical considerations for oxygen radical formation and lipid peroxidation. Free Radical Biol & Med (1989) 6, 289-301.
- Niwa Y, Sakane T, Singu M and Yokohama M: Effect of stimulated neutrophils from synovial fluid of patients with rheumatoid arthritis on lymphocytes: A possible role of increased oxygen radicals generated by the neutrophils. J Clin Immunol (1983) 3, 228-240.
- Sun Y: Free radicals, antioxidant enzymes, and carcinogenesis. Free Radical Biol & Med. (1990) 8, 583-599.
- Babbs CF: Free radicals and the etiology of colon cancer. Free Radical Biol & Med. (1990) 8, 191-200.
- Liu J and Mori A: Involvement of reactive oxygen species in emotional stress. Int J Stress Management (1994) 1, 249-263.
- Liu J, Edamatsu R, Kabuto H and Mori A: Antioxidant action of guilingji in the brain of rats with FeCl₂-induced epilepsy. Free Rad Biol & Med (1990) 9, 451-454.
- Mori A, Liu J, Wang X and Kawai M: Free radical scavenging by brain homogenate: Implication of free radical damage and antioxidant defense in brain. Neurochem Int (1994) 24, 201-207.
- Ohkawa H, Ohishi N and Yagi K: Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. Anal Biochem (1979) 95, 351-358.
- Larson AR: The antioxidants of higher plants. Phytochemistry (1988) 27, 969-974.
- Hiramatsu M and Mori A: Antioxidant herbal medicine; in Sozai-No-ESR Hyoka-Hou, Ohya H and Yamaguchi J, eds, IPC Press, Tokyo (1992) pp 470-485 (in Japanese).
- Okuda T and Yoshikawa T: Free Radicals to Wakanyaku. Kokusai Isho Shuppan, Tokyo (1991) (in Japanese).
- Wang X, Liu J and Mori A: Selective hydroxyl radical scavenging activity of injection Ligustrazine Phosphatid and anti embolism venin enzyme. Neurosciences (1992) 18, 27-35.
- Santiago LA, Hiramatsu M and Mori A: Japanese soybean paste miso scavenges free radicals and inhibits lipid peroxidation. J Nutr Sci Vitaminol (1992) 38, 297-304.
- Hiramatsu M, Edamatsu R, Kohno M and Mori A: Scavenging of free radicals by Sho-saiko-to-go-keishi-ka-shakuyaku-to (TJ-960); Recent Advances in the Pharmacology of KAMPO (Japanese herbal) Medicines, Excerpta Medica, Amsterdam (1988) pp 120-127.
- Jiang-su-xin-yi-xue-yuan: Zhongyao Dacidian, Shanghai People's Publishing House, Shanghai. (1978) pp 2387-2388 (in Chinese).
- Liu D: The ants powder treated rheumatoid arthritis in 808 patients. Convalescent Med (1989) 8, 6 (in Chinese).
- Wu Z: The ants treated rheumatoid arthritis in 4100 patients. Chin Med Res (1989) 2, 29 (in Chinese).
- Zhou Y, Lou Q, Wang C, Zhang X, Xie S and Chen D: The ants treated hepatitis in clinic observation. Chin Med Pharmacol Clin (1994) 6, 40-41 (in Chinese).
- Wang Z: The extract of black ants recovered immunological regulating effects and anti-aging effects. Gerontology (1987) 7, 41 (in Chinese).

Received June 8, 1995; accepted August 22, 1995.