

# *Acta Medica Okayama*

---

*Volume 28, Issue 5*

1974

*Article 1*

OCTOBER 1974

---

## Studies on ferrous ion-induced lipid peroxidation of rat liver mitochondria. I. Effect of inorganic phosphate

Goki Yamamoto\*

Masatada Tanabe<sup>†</sup>

Hiroshi Wakabayashi<sup>‡</sup>

Gonosuke Hashimoto\*\*

Michio Yamamoto<sup>††</sup>

\*Okayama University,

<sup>†</sup>Okayama University,

<sup>‡</sup>Okayama University,

\*\*Okayama University,

<sup>††</sup>Okayama University,

# Studies on ferrous ion-induced lipid peroxidation of rat liver mitochondria. I. Effect of inorganic phosphate\*

Goki Yamamoto, Masatada Tanabe, Hiroshi Wakabayashi, Gonosuke Hashimoto,  
and Michio Yamamoto

## Abstract

Effect of inorganic phosphate on ferrous ion- and ascorbate-induced lipid peroxidations of isolated rat liver mitochondria was investigated. As a result it has been shown that phosphate accelerates the ferrous ion-induced lipid peroxidation; namely, phosphate shortens the induction lag period of the lipid peroxidation reaction but the malondialdehyde after onset of its production is yielded at the same rate in various concentrations of phosphate. On the other hand, phosphate inhibits ascorbate-induced lipid peroxidation. There are stoichiometric interactions between the concentration of phosphate and the induction period. Oxygen uptake by mitochondria was observed in the presence of both ferrous ion and phosphate at initial step of the reaction without being accompanied by malondialdehyde production, and afterwards there occurred malondialdehyde production with rapid rate of the oxygen uptake. Possible mechanisms and interactions among ferrous ion, ascorbate and phosphate were discussed.

---

\*PMID: 4281994 [PubMed - indexed for MEDLINE] Copyright ©OKAYAMA UNIVERSITY  
MEDICAL SCHOOL

Acta Med. Okayama 28, 299—310 (1974)

**STUDIES ON FERROUS ION-INDUCED LIPID PEROXIDATION  
OF RAT LIVER MITOCHONDRIA,  
I. EFFECT OF INORGANIC PHOSPHATE**

Goki YAMAMOTO, Masatada TANABE, Hiroshi WAKABAYASHI,  
Gonosuke HASHIMOTO and Michio YAMAMOTO

*Department of Radiation Medicine, Okayama University Medical School,  
Okayama, Japan (Director: Prof. M. Yamamoto)*

*Received for publication, April 23, 1974*

*Abstract:* Effect of inorganic phosphate on ferrous ion- and ascorbate-induced lipid peroxidations of isolated rat liver mitochondria was investigated. As a result it has been shown that phosphate accelerates the ferrous ion-induced lipid peroxidation; namely, phosphate shortens the induction lag period of the lipid peroxidation reaction but the malondialdehyde after onset of its production is yielded at the same rate in various concentrations of phosphate. On the other hand, phosphate inhibits ascorbate-induced lipid peroxidation. There are stoichiometric interactions between the concentration of phosphate and the induction period. Oxygen uptake by mitochondria was observed in the presence of both ferrous ion and phosphate at initial step of the reaction without being accompanied by malondialdehyde production, and afterwards there occurred malondialdehyde production with rapid rate of the oxygen uptake. Possible mechanisms and interactions among ferrous ion, ascorbate and phosphate were discussed.

The presence of unsaturated fatty acids in the phospholipid of mitochondrial membrane appeared to be essential for energy transfer reactions (1). It is of importance to consider the possibility that some intermediates of oxidative phosphorylation are associated with the polyunsaturated fatty acids which lead to lipid peroxide formation or accumulation under certain conditions (2). It has been reported that lipid peroxides may be formed during normal activity of electron transport chain in vitamin E-deficient rat liver mitochondria (3). Phosphate induces mitochondrial swelling which is dependent on electron transport and generation of high energy intermediates in the process of oxidative phosphorylation (4, 5). It is of interest to know the interaction of phosphate with lipid peroxidation in mitochondria.

Mitochondrial lipid peroxidation has been demonstrated correlatively with mitochondrial membrane alterations (swelling, lysis and disintegration) (2, 6, 7). Ferrous ion seems to be specific to the initiation of lipid peroxidation in mitochondria (8-11). However, the mechanism of mitochondrial lipid peroxidation

300 G. YAMAMOTO, M. TANABE, H. WAKABAYASHI, G. HASHIMOTO and M. YAMAMOTO

is not understood quite well. Ferrous ion-induced lipid peroxidation of mitochondria has a lag of induction period (8, 12). The lag is observed far longer in high concentration of  $\text{Fe}^{++}$  (8), and is shortened by addition of  $\text{Fe}^{+++}$ , suggesting that oxidation-reduction rate in the incubation system is an important factor (8). HUNTER *et al.*(2) reported that phosphate, which induces no lipid peroxidation, inhibits the lipid peroxidation of mitochondria induced by the mixture of oxidized and reduced glutathion (GSSG + GSH) and by ascorbate. But many workers (9, 11, 13-16), who have observed lipid peroxide formation in mitochondria under various conditions, have overlooked the effect of phosphate used in their study. To clarify the interaction of phosphate in mitochondrial lipid peroxidation, effect of inorganic phosphate on lipid peroxidation of isolated rat liver mitochondria induced by  $\text{Fe}^{++}$  or ascorbate was studied in particular on the lag of induction period. Our report describes that inorganic phosphate shortens the lag of the induction period of  $\text{Fe}^{++}$ -induced lipid peroxidation, and lengthens that of ascorbate-induced one.

#### MATERIALS AND METHODS

*Preparation of mitochondria*: Albino rats (200 to 300g) derived from the Don-ryu strain were maintained on mouse-rat diet, MF type of Oriental Co.

Rat was killed by decapitation and the liver was immediately immersed in a chilled STE-solution (0.25M sucrose, 0.003M Tris-HCl and 0.0002M EDTA, pH 7.4). All procedures of preparation were carried out at 0-4°C by a modified method of HOGEBOM (17). The liver was cut with scissors to small pieces, washed in the STE solution, homogenized roughly by a glass homogenizer in ten volumes of the STE-solution, and gently homogenized by a teflon homogenizer. The homogenate was centrifuged at 100×g for 10 min. Nuclei and cell debris were removed and supernatant was layered on an equal volume of 0.34M sucrose containing 0.003M Tris-HCl, pH 7.4, and centrifuged at 700×g for 10min. The upper layers were gathered and centrifuged at 5,000×g for 10min to reduce contamination of microsomal materials. After cautious removal of fluffy layers, the precipitate was suspended in the STE-solution, centrifuged, and then washing was repeated in an ST-solution (0.25M sucrose and 0.003M Tris-HCl, pH 7.4). Final mitochondrial precipitate was resuspended in the ST-solution and maintained in a concentrated suspension in an ice-bath. Mitochondria were washed in a large volume of KT-medium (0.15M KCl and 0.01M Tris-HCl, pH 7.4), centrifuged, and suspended in the KT-medium by a teflon homogenizer before use.

*Incubation system and reagents*: Incubation system was composed of about 1 mg of mitochondrial protein per ml in KT-medium and/or in KP-medium (0.15M KCl and 0.01M Na-K-phosphate buffer). The medium was preincubated for 10min and the reaction was started by adding 0.25mM  $\text{Fe}^{++}$  and/or 1mM ascorbate in final concentration. Incubation at 4°C was carried out in an ice-water bath at 0-4°C.  $\text{Fe}^{++}$  used was ammonium ferrous sulfate pur-

chased from Katayama Kagaku Co., and other reagents were of special grade. Distilled and deionized water was used. Protein content was determined by Biuret method (18) with bovine serum albumin (Armour Laboratories, Fraction V) as a standard.

**TBA color reaction:** As a diagnostic test for lipid peroxidation, malondialdehyde (MA) produced was detected. The detection of MA by thiobarbituric acid (TBA) was carried out by a modified method of HUNTER *et al.* (8). After incubating for various lapses of time, 2ml of medium was mixed with the solution composed of 0.5ml of 40% trichloroacetic acid, 0.25ml of 5N HCl and 0.5ml of 2% TBA to stop the reaction, and then heated in a boiling-water bath for 10 min. The mixtures were chilled and centrifuged. The absorbance of clear supernatant was measured by spectrophotometer at 532 nm and 600 nm. TBA value was represented by  $OD_{532-600}/\text{mg}$  of protein. Peroxide is expressed in terms of MA, using the molar extinction coefficient,  $E_{532} = 1.56 \times 10^5 \text{ cm}^2 \text{ mmole}^{-1}$  (19).

**Oxymetric method:** Oxygen consumption was measured with a Bioxygraph (Kusui-Kagaku Co.) (20) with a 2ml-chamber kept at 25°C. Other details are described in the legend of figure.

## RESULTS

**Effect of phosphate on MA formation:** The characteristic curve for MA production in  $\text{Fe}^{++}$ -induced lipid peroxidation of mitochondria in KT-medium shows a lag of induction period followed by a rapid increase.  $\text{Fe}^{++}$  at a high concentration had a far longer lag but produced a larger amount of MA after




Fig. 1. Effect of inorganic phosphate (Pi) on the lipid peroxidation of rat liver mitochondria induced by  $\text{Fe}^{++}$  in KT (0.15M KCl, 0.01M, Tris-Cl, pH 7.4)-medium at 37°. Pi concentration in the medium is given in figure and reaction was started by addition of  $\text{Fe}^{++}$  (0.25mM).

302 G. YAMAMOTO, M. TANABE, H. WAKABAYASHI, G. HASHIMOTO and M. YAMAMOTO

the onset of the reaction than at a lower concentration of  $\text{Fe}^{++}$ . As shown in Fig. 1, in the presence of phosphate,  $\text{Fe}^{++}$ -induced lipid peroxidation at  $37^\circ\text{C}$  reflected the shortening of the lag period of induction. Increment of phosphate concentration further shortened the lag, but the velocity of MA production and the amount of MA produced were almost identical regardless of difference in the lag period. The changes of the induction period induced by phosphate in various concentrations of  $\text{Fe}^{++}$  were proportional to those in the control system without phosphate,




Fig. 2. Relation between the concentration of  $\text{Fe}^{++}$  and the induction period of mitochondrial lipid peroxidation induced by  $\text{Fe}^{++}$ . Incubation was carried out at  $37^\circ$  in KT-medium with or without 2mM Pi. Induction period was calculated from the intervals before the onset of rapid MA formation.

*Effect of pH on MA formation in phosphate buffer:* Effect of pH on the lipid peroxidations incubated at  $37^\circ\text{C}$  in KP-medium was tested. In  $\text{Fe}^{++}$ -induced lipid peroxidation, the onset of peroxidation was early in alkaline medium but MA was produced more abundantly at a later time in the range of pH 6 to 7 (Fig. 5). On the other hand, ascorbate-induced lipid peroxidation was optimum at pH 6, and hardly reacted in alkaline solution (Fig. 6).

indicating a linear correlation between the lags of induction and concentrations of  $\text{Fe}^{++}$  (Fig. 2).

On the contrary, phosphate prevented the appearance of ascorbate-induced MA formation at  $37^\circ\text{C}$  involving the lengthened lag period as shown in Fig. 3.

Judging from such divergent effects of phosphate on the mitochondrial lipid peroxidations induced by  $\text{Fe}^{++}$  and ascorbate, there seems to be a linear correlation between logarithmic values of the induction period and the molarity of phosphate in a certain range as illustrated in Fig. 4.




Fig. 3. Inhibitory effect of Pi on the lipid peroxidation induced by ascorbate (1 mM) in KT-medium at  $37^\circ$ .




Fig. 4. Relation between the concentration of Pi and the induction period of Fe<sup>2+</sup>- and ascorbate-induced lipid peroxidation of mitochondria. Experimental conditions were identical with those in Figs. 1 and 3.




Fig. 5. Effect of pH on Fe<sup>2+</sup>-induced lipid peroxidation of mitochondria. The incubation system was composed of KP (0.15M KCl, 0.01 M Na-K-phosphate buffer)-medium at 37°. Produced MAs at 2, 5 and 60 min after addition of 0.25 mM Fe<sup>2+</sup> are graphed respectively.




Fig. 6. Effect of pH on ascorbate-induced lipid peroxidation of mitochondria. The incubation system was composed of KP-medium at 37°. Produced MAs at 20, 30 and 60 min after addition of 1 mM ascorbate are graphed respectively.

304 G. YAMAMOTO, M. TANABE, H. WAKABAYASHI, G. HASHIMOTO and M. YAMAMOTO

*Effect of temperature on MA formation:* McKNIGHT *et al.* (21) reported that the inhibitory effect of  $\text{Fe}^{++}$  at high concentration in producing lag period before the onset of peroxidation in the KCl-malate medium was abolished at  $0^\circ\text{C}$ . This fact suggests non-enzymatic process of peroxidation in mitochondria. Peroxidation of mitochondria and heat-treated mitochondria was, therefore, compared at  $4^\circ\text{C}$  and  $37^\circ\text{C}$  in KP-medium, pH 7.4. The heat-treated mitochondria suspended in KT-medium were obtained by heating in a boiling-water bath for two min and resuspended by teflon homogenizer. It can be seen from Fig. 7 that  $\text{Fe}^{++}$ -induced peroxidation of non-treated mitochondria at  $37^\circ\text{C}$  almost allows no lag period, and at  $4^\circ\text{C}$  the reaction has a




Fig. 7.  $\text{Fe}^{++}$ -induced lipid peroxidations of mitochondria and heat-treated mitochondria. Incubations were carried out at  $37^\circ$  or in an ice-water bath, and the reaction was started by addition of  $0.25 \text{ mM Fe}^{++}$  in the system of  $0.15 \text{ M KCl}$  and  $0.01 \text{ M Na-K-phosphate buffer}$ , pH 7.4.

lag period. At both temperatures the peroxidation of heat-treated mitochondria occurs later than that of non-treated one. On the other hand, ascorbate-induced peroxidation occurs faster in the heat-treated mitochondria at both temperatures than in non-treated mitochondria, as shown in Fig. 8. Ascorbate-induced reaction of non-treated mitochondria appears to be not producing MA at  $4^\circ\text{C}$  in the tested period of time.

*Oxygen uptake on mitochondrial lipid peroxidation:* In an earlier report it has been demonstrated that there is an extra oxygen consumption during the






Fig. 8. Ascorbate-induced lipid peroxidations of mitochondria and heat-treated mitochondria. Ascorbate used was 1 mM and other conditions were identical as in Fig. 7.

Fe<sup>++</sup>-induced lipid peroxidation (12). Oxygen uptake of mitochondria following the lipid peroxidation in the present system was measured at 25°C as recorded in Fig. 9. Endogenous oxygen consumption of mitochondria in KT-




Fig. 9. Oxygen electrode traces involved in the Fe<sup>++</sup>-induced lipid peroxidation of mitochondria (2.1 mg of protein) in 2 ml of KT-medium at 25°. Concentrations of added reagents are shown in Fig. Respective records are summarized. Dotted lines show the traces after addition of EDTA, and the trace marked +EDTA was recorded in the medium containing 0.5 mM EDTA. The left under table represents the yields of MA at point described by capital letters with arrows in the recorded traces.

306 G. YAMAMOTO, M. TANABE, H. WAKABAYASHI, G. HASHIMOTO and M. YAMAMOTO

medium could hardly be observed. The addition of  $\text{Fe}^{++}$  or phosphate alone had almost no effect on the oxygen uptake in a short period after the addition. But the addition of  $\text{Fe}^{++}$  in the presence of phosphate induced immediately a rapid oxygen uptake (Fig. 9). Similar oxygen uptake was observed on the addition of phosphate in the presence of  $\text{Fe}^{++}$ . The velocity of this initial burst of oxygen uptake was dependent on the concentration of phosphate (Fig. 10), but MA was not produced at the periods (checked by arrow B in




Fig. 10. Changes of oxygen uptake in mitochondrial lipid peroxidation in the presence of Pi at various concentrations. Incubation system at 25° was composed of 2.1 mg protein of mitochondria, KT-medium (pH 7.4), and Pi (pH 7.4, concentration shown in Fig.), total volume 2 ml.  $\text{Fe}^{++}$  (0.25 mM) was added at point shown by arrow, and respective records are summarized. MAs produced at the end of reactions (shown in Fig. by arrow) are 16.6, 13.6, 12.8 and 13.2 nmoles in 10, 5, 2.5 and 1 mM Pi, respectively.

Fig. 9), revealing that the oxygen uptake occurs during the lag period of induction of peroxidation. After a certain period of such oxygen consumption, the oxygen uptake was again observed in association with the production of MA (arrows G & F). Peroxidation was inhibited rapidly on the addition of EDTA which decreased both oxygen uptake and MA production (arrow D), even in the midst of peroxidation reaction (arrow E). But even in the presence of EDTA, the oxygen uptake of mitochondria was induced on the addition of phosphate and  $\text{Fe}^{++}$ . The oxygen uptake in such cases was larger than in the absence of EDTA. Needless to say, the oxygen uptake in this instance was not accompanied by MA production (arrow C). The molar ratio of MA produced per oxygen calculated from the consumed oxygen in relation to the MA

production was approximately 0.05.

#### DISCUSSION

HUNTER *et al.* (2) state that phosphate at relatively low concentrations prevents the lipid peroxide production induced by GSSG+GSH in mitochondria, and point out some possible role of phosphate. The inhibitory behaviors of phosphate at high concentrations on ascorbate- and  $\text{Fe}^{++}$ -induced lipid peroxidation were also reported (2, 8). In the present study, the fact that phosphate prevents the ascorbate-induced lipid peroxidation in rat liver mitochondria coincides with other reports (2, 22). Since ascorbate-induced lipid peroxidation was facilitated at high temperature and in heat-treated mitochondria, the structural change of mitochondria could be preferential to lipid peroxidation reaction. However, such a lipid peroxidation was prevented by phosphate, so that phosphate might act as an antioxidant associated probably with oxidation of ascorbate. HUNTER *et al.* (2) suggested that phosphate inhibits GSSG+GSH-induced changes in mitochondria by a direct antioxidant action, possibly in a non-specific manner. It has been pointed out that phosphoric acid may belong to the electron-donating class of antioxidants (23).

The fact that phosphate eliminates the lag period of  $\text{Fe}^{++}$ -induced lipid peroxidation of mitochondria does not coincide with the other reports (2, 4). From the results of shortening the lag period and of pH dependency, the reaction of phosphate on lipid peroxidation induced by  $\text{Fe}^{++}$  might differ from that by ascorbate.

A number of interactions of phosphate with  $\text{Fe}^{++}$ -induced lipid peroxidation may be considered as follows:

(a) Since spontaneous swelling of mitochondria will permit  $\text{Fe}^{++}$  to penetrate into the site of certain unsaturated lipid (8), phosphate-induced swelling might affect the initiation of  $\text{Fe}^{++}$ -induced lipid peroxidation, but the accelerating action of phosphate can also be observed in heat-treated mitochondria.

(b) MA was formed earlier at a low concentration of  $\text{Fe}^{++}$  than at its high concentration (8). Practical change of  $\text{Fe}^{++}$  with phosphate toward insoluble salts might be considered for the lowering the concentration of  $\text{Fe}^{++}$ . However, yields of MA in a steady state after the onset of lipid peroxide formation in various concentrations of phosphate were fairly constant. Thus, a complex salt, if formed, might be of an active form. While, MA-producing velocity of mitochondria after onset of the reaction with or without phosphate is likely to be of the same rate. It is suggested that phosphate affects the initiation of lipid peroxidation reaction chain, and enhances the pre-reaction such as producing free radical intermediates.

(c) HUNTER *et al.* (8) have described that  $\text{Fe}^{++}$  has no effect at pH 8.0,

possibly because of its rapid conversion to  $\text{Fe}^{+++}$  and precipitation of ferric hydroxide.  $\text{Fe}^{++}$ -induced MA production in KP-medium occurred readily in an alkaline medium, but that at pH 6–7 was followed by the high rate of its production.  $\text{Fe}^{+++}$  added to the system eliminates the lag-producing effect in  $\text{Fe}^{++}$ -induced lipid peroxidation (8), then the rapid conversion of  $\text{Fe}^{++}$  to  $\text{Fe}^{+++}$  in alkaline might be considered to be responsible for the initiation of MA production. The mechanism of action of  $\text{Fe}^{+++}$  is not entirely clear, but it is possible that  $\text{Fe}^{+++}$  acts by rapidly releasing MA from oxidized fatty acid during incubation (24), and the oxidation-reduction rate in incubation system is an important role.

(d) The yield of MA (TBA chromogen) during peroxidation induced by  $\text{Fe}^{++}$  is far greater at  $0^\circ\text{C}$  than at  $25^\circ\text{C}$  (14). However, in KP-medium, the lag period was further lengthened at  $4^\circ\text{C}$  than at  $37^\circ\text{C}$ . The present experiment has not elucidated whether or not  $\text{Fe}^{++}$ -induced lipid peroxidation of mitochondria is involved in enzymatic reaction, but phosphate-induced alterations of structure and function in mitochondrial membrane dependent on temperature seem to be specific to the initial step of  $\text{Fe}^{++}$ -catalized peroxidation.

(e) Incorporation or binding of  $\text{Fe}^{++}$  in mitochondria is likely performed by both enzymatic and nonenzymatic processes, and the interaction between  $\text{Fe}^{++}$  and phosphoric base of phospholipid has been proposed in relation to the formation of metal-chelate complex (25, 26). Oxymetric experiments showed that the oxygen uptake in the presence of  $\text{Fe}^{++}$ , phosphate and mitochondria was observed without MA production, suggesting that it may reflect a metal-chelate complex formation with oxygen also being involved. TBA reaction (27), being employed to detect MA for judging lipid peroxidation, does not divulge peroxidation reaction probably occurring during the lag period (28). The metal complex formation with oxygen seems to be an important role in peroxidation reaction. These compounds may be a reflection of oxidation of  $\text{Fe}^{++}$  to  $\text{Fe}^{+++}$  for the production of free radical intermediates which initiate lipid peroxidation chain reactions.

The mechanisms of interaction of phosphate on lipid peroxidation in mitochondria induced by  $\text{Fe}^{++}$  and ascorbate seem to be variable as above discussed. Therefore, further detailed experiments are required to clarify a great possibility of cooperative and control mechanisms of lipid peroxidation among  $\text{Fe}^{++}$ , ascorbate and phosphate in mitochondria.

*Acknowledgements:* We express thanks to Associate Prof. Kozo UTSUMI, Department of Biochemistry, Cancer Institute of Okayama University Medical School, for helpful suggestions and painstaking proof reading of this manuscript, and to Mrs. Tomiko OHTSU for technical assistance.

## REFERENCES

1. KAGAWA, Y., KANDRACH, A. and RACKER, E.: Partial resolution of the enzymes catalyzing oxidative phosphorylation. XXVI. Specificity of phospholipids required for energy transfer reactions. *J. Biol. Chem.* **248**, 676-684, 1973
2. HUNTER, F. E. Jr., SCOTT, A., WEINSTEIN, J. and SCHNEIDER, A.: Effects of phosphate, arsenate, and other substances on swelling and lipid peroxide formation when mitochondria are treated with oxidized and reduced glutathione. *J. Biol. Chem.* **239**, 622-630, 1964
3. CORWIN, L. M. and SCHWARZ, K.: Prevention of decline of  $\alpha$ -ketoglutarate and succinate oxidation in vitamin E-deficient rat liver homogenates. *J. Biol. Chem.* **235**, 3387-3392, 1960
4. PACKER, L.: Metabolic and structural states of mitochondria. II. Regulation by phosphate. *J. Biol. Chem.* **236**, 214-220, 1961
5. UTSUMI, K.: Relation between mitochondrial swelling induced by inorganic phosphate and accumulation of  $^{32}\text{P}$  in mitochondrial Pi fraction. *Acta Med. Okayama* **17**, 259-271, 1963
6. HUNTER, F. E. Jr., SCOTT, A., HOFFSTEN, P. E., GUERRA, F., WEINSTEIN, J., SCHNEIDER, A., SCHUTZ, B., LILLIAN, F. and SMITH, E.: Studies on the mechanism of ascorbate-induced swelling and lysis of isolated liver mitochondria. *J. Biol. Chem.* **239**, 604-613, 1964
7. HUNTER, F. E. Jr., SCOTT, A., HOFFSTEN, P. E., GEBICKI, J. M., WEINSTEIN, J. and SCHNEIDER, A.: Studies on the mechanism of swelling, lysis, and disintegration of isolated liver mitochondria exposed to mixtures of oxidized and reduced glutathione. *J. Biol. Chem.* **239**, 614-621, 1964
8. HUNTER, F. E. J., GEBICKI, J. M., HOFFSTEN, P. E., WEINSTEIN, J. and SCOTT, A.: Swelling and lysis of rat liver mitochondria induced by ferrous ions. *J. Biol. Chem.* **238**, 828-835, 1963
9. OTTOLENGHI, A.: Interaction of ascorbic acid on mitochondrial lipids. *Arch. Biochem. Biophys.* **79**, 355-363, 1959
10. FORTNEY, S. R. and LYNN, W. S. Jr.: Role of ascorbate and cysteine on swelling and lipid peroxidation in rat liver mitochondria. *Arch. Biochem. Biophys.* **104**, 241-247, 1964
11. THIELE, E. H. and HUFF, J. W.: Quantitative measurements of lipid peroxide formation by normal liver mitochondria under various conditions. *Arch. Biochem. Biophys.* **66**, 203-207, 1960
12. UTSUMI, K. and YAMAMOTO, G.: Failure of  $\text{Fe}^{++}$ -induced lipid peroxidation and swelling in the mitochondria isolated from ascites tumor cells. *Biochim. Biophys. Acta* **105**, 368-371, 1965
13. TAPPEL, A. L. and ZALKIN, H.: Lipid peroxidation in isolated mitochondria. *Arch. Biochem. Biophys.* **80**, 326-332, 1959
14. ZALKIN, H. and TAPPEL, A. L.: Mechanism of vitamin E action. IV. Lipid peroxidation in the vitamin E-deficient rabbit. *Arch. Biochem. Biophys.* **88**, 113-117, 1960
15. HATEFI, Y. and HANSTEIN, W. G.: Lipid oxidation in biological membranes. I. Lipid oxidation in submitochondrial particles and microsomes induced by chaotropic agents. *Arch. Biochem. Biophys.* **138**, 73-86, 1970.
16. DILLARD, C. J. and TAPPEL, A. L.: Fluorescent products of lipid peroxidation of mitochondria and microsomes. *Lipids* **6**, 715-721, 1971
17. HOGEBOOM, G. H.: Fractionation of cell components of animal tissues. In *Methods in Enzymology*, eds. Colowick, S. P. and Kaplan, N. O., Vol. I, pp 16-19, Academic Press Inc., New York, 1955
18. GORNALL, A. G., BARDAWILL, C. J. and DAVID, M. M.: Determination of serum proteins by means of the Biuret reaction. *J. Biol. Chem.* **177**, 751-766, 1949

310 G. YAMAMOTO, M. TANABE, H. WAKABAYASHI, G. HASHIMOTO and M. YAMAMOTO

19. SINNHUBER, R. O. and YU, T. C.: 2-Thiobarbituric acid method for the measurement of rancidity in fishery products. II. The quantitative determination of malonaldehyde. *Food Technol.* **12**, 9-12, 1958
20. UTSUMI, K., ODA, T., KURAHASHI, K., MIYAHARA, M. and YASUDA, M.: Measurement of oxygen consumption by galvanic electrode. *Protein, Nucleic Acid and Enzyme* **14**, 621-624, 1969 (in Japanese)
21. MCKNIGHT, R. C., HUNTER, F. E. Jr. and OEHLERT, W. H.: Mitochondrial membrane ghosts produced by lipid peroxidation induced by ferrous ion. I. Production and general morphology. *J. Biol. Chem.* **240**, 3439-3446, 1965
22. CHRISTOPHERSEN, B. O.: The inhibitory effect of reduced glutathione on the lipid peroxidation of the microsomal fraction and mitochondria. *Biochem. J.* **106**, 515-522, 1968
23. STUCKEY, B. N.: Antioxidants. *Symposium on foods: lipids and their oxidation*, In eds. Schultz, H. W., Day, E. A. and Sinnhuber, R. O., AVI Publishing Co., Westport, Conn., pp 139-150, 1961
24. WILLS, E. D.: The effect of inorganic iron on the thiobarbituric acid method for the determination of lipid peroxides. *Biochim. Biophys. Acta* **84**, 475-477, 1964
25. MAZANOWSKA, A. M., NEUBERGER, A. and TAIT, G. H.: Effect of lipid and organic solvents on the enzymic fraction of zinc protoporphyrin and heam. *Biochem. J.* **98**, 117-127, 1966
26. ROMSLO, I. and FLATMARK, T.: Energy-dependent accumulation of iron by isolated rat liver mitochondria. I. General features. *Biochim. Biophys. Acta* **305**, 29-40, 1973
27. DAHLE, L. K., HILL, E. G. and HOLMAN, R. T.: The thiobarbituric acid reaction and the autoxidation of polyunsaturated fatty acid methyl esters. *Arch. Biochem. Biophys.* **98**, 253-261, 1962
28. TAM, B. K. and McCAY, P. B.: Reduced triphosphopyridine nucleotide oxidase-catalyzed alteration of membrane phospholipids. III. Transient formation of phospholipid peroxides. *J. Biol. Chem.* **245**, 2295-2300, 1970