

Acta Medica Okayama

Volume 61, Issue 4

2007

Article 5

AUGUST 2007

Seven Different Glucose-6-phosphate Dehydrogenase Variants Including a New Variant Distributed in Lam Dong Province in Southern Vietnam.

Hiroyuki Matsuoka*

Dang Thi Vinh Thuan†

Huynh van Thien‡

Toshio Kanbe**

Amadu Jalloh††

Makoto Hirai‡‡

Meiji Arai§

Nguyen The Dung¶

Fumihiko Kawamoto||

*Jichi Medical University,

†Bao Loc General Hospital,

‡Bao Loc General Hospital,

**Nagoya University,

††Nagoya University,

‡‡Jichi Medical University,

§Jichi Medical University,

¶Ho Chi Minh City Health Department,

||Oita University,

Seven Different Glucose-6-phosphate Dehydrogenase Variants Including a New Variant Distributed in Lam Dong Province in Southern Vietnam.*

Hiroyuki Matsuoka, Dang Thi Vinh Thuan, Huynh van Thien, Toshio Kanbe, Amadu Jalloh, Makoto Hirai, Meiji Arai, Nguyen The Dung, and Fumihiko Kawamoto

Abstract

We conducted a survey for glucose-6-phosphate dehydrogenase (G6PD) deficiency using blood samples from male outpatients of a local hospital in southern Vietnam. Most of the samples were from the Kinh (88.9%), the largest ethnic group in Vietnam, with a small number (11.1%) coming from the K'Ho, Chauma, Nung, and Tay minorities. We detected 25 G6PD-deficient cases among 1,104 samples (2.3%), and read the open reading frame of G6PD. A novel mutation (352T>C) predicting an aminoacid change of 118Tyr>His was found in a 1-year-old Kinh boy. His G6PD activity was estimated to be less than 10% residual activity, although he did not show chronic hemolytic anemia. Thus, we categorized this variant as Class II and named it G6PD Bao Loc. In the Kinh population, G6PD Viangchan (871G>A, 1311C>T, intron 11 nt93T>C), one of the most common variants in continental Southeast Asian populations, was the highest (6/19), followed by variants originating from the Chinese such as G6PD Canton (1376G>T) (5/19), G6PD Kaiping (1388G>A) (3/19), G6PD Gaohe (95A>G) (1/19), and G6PD Quing Yuan (392G>T) (1/19). In addition, G6PD Union (1360C>T) (2/19), which originated from the Oceania, was also detected. These findings suggest that the Kinh people are derived from various ancestries from continental Southeast Asia, China, and Oceania. In contrast, all of the 5 deficient cases in the K'Ho population were G6PD Viangchan, suggesting that they were very close to Southeast Asian populations such as the Khmer in Cambodia and the Lao in Laos. It is interesting that G6PD Mahidol (487G>A), another common variant in continental Southeast Asian populations in Myanmar, Thailand, and Malaysia, has not been detected from the Vietnamese.

KEYWORDS: Bao Loc, glucose-6-phosphate dehydrogenase deficiency, Kinh, malaria, Vietnam

Seven Different Glucose-6-phosphate Dehydrogenase Variants Including a New Variant Distributed in Lam Dong Province in Southern Vietnam

Hiroyuki Matsuoka^{a*}, Dang Thi Vinh Thuan^b, Huynh van Thien^b,
Toshio Kanbe^c, Amadu Jalloh^c, Makoto Hirai^a, Meiji Arai^a,
Nguyen The Dung^d, and Fumihiko Kawamoto^e

^aDivision of Medical Zoology, Department of Infection and Immunity, Jichi Medical University, Shimotsuke, Tochigi 329-0498, Japan,

^bBao Loc General Hospital, Lam Dong Province, Vietnam, ^cDepartment of Advanced Medical Science, Nagoya University Graduate School of Medicine, Nagoya 466-8560, Japan, ^dHo Chi Minh City Health Department, Ho Chi Minh, Vietnam, and

^eInstitute of Scientific Research, Faculty of Medicine, Oita University, Yubu, Oita 879-5593, Japan

We conducted a survey for glucose-6-phosphate dehydrogenase (G6PD) deficiency using blood samples from male outpatients of a local hospital in southern Vietnam. Most of the samples were from the Kinh (88.9%), the largest ethnic group in Vietnam, with a small number (11.1%) coming from the K'Ho, Chauma, Nung, and Tay minorities. We detected 25 G6PD-deficient cases among 1,104 samples (2.3%), and read the open reading frame of *G6PD*. A novel mutation (352T > C) predicting an amino-acid change of I18Tyr > His was found in a 1-year-old Kinh boy. His G6PD activity was estimated to be less than 10% residual activity, although he did not show chronic hemolytic anemia. Thus, we categorized this variant as Class II and named it G6PD Bao Loc. In the Kinh population, G6PD Viangchan (871G > A, 1311C > T, intron II nt93T > C), one of the most common variants in continental Southeast Asian populations, was the highest (6/19), followed by variants originating from the Chinese such as G6PD Canton (1376G > T) (5/19), G6PD Kaiping (1388G > A) (3/19), G6PD Gaohe (95A > G) (1/19), and G6PD Quing Yuan (392G > T) (1/19). In addition, G6PD Union (1360C > T) (2/19), which originated from the Oceania, was also detected. These findings suggest that the Kinh people are derived from various ancestries from continental Southeast Asia, China, and Oceania. In contrast, all of the 5 deficient cases in the K'Ho population were G6PD Viangchan, suggesting that they were very close to Southeast Asian populations such as the Khmer in Cambodia and the Lao in Laos. It is interesting that G6PD Mahidol (487G > A), another common variant in continental Southeast Asian populations in Myanmar, Thailand, and Malaysia, has not been detected from the Vietnamese.

Key words: Bao Loc, glucose-6-phosphate dehydrogenase deficiency, Kinh, malaria, Vietnam

Received November 22, 2006; accepted February 19, 2007.

*Corresponding author. Phone: +81-285-58-7339; Fax: +81-285-44-6489
E-mail: hiroyuki@jichi.ac.jp (H. Matsuoka)

We have introduced rapid diagnosis methods for malaria and the G6PD deficiency test in malaria endemic areas. With these methods, patients are notified of the results of the blood examination within

30 min and are able to receive anti-malaria medicine, including primaquine [1]. Primaquine can kill gametocytes, the sexual stage of malaria parasites, which are the cause of malaria transmission to mosquitoes. However, once G6PD-deficient persons take primaquine, a hemolysis attack can follow. Without G6PD, erythrocytes cannot prepare a sufficient amount of reduced pyridine nucleotide and reduced glutathione, and cannot prevent oxidant attack by primaquine. Thus, primaquine should not be administered to malaria patients before confirming their G6PD condition.

In the past, there have been many malaria endemic areas in Vietnam. Although the numbers of malaria cases and malaria endemic areas are decreasing, malaria is still a serious disease in Vietnam. It is therefore of value to introduce rapid diagnosis methods for malaria and G6PD deficiency in rural areas in Vietnam. We started the technical transfer to a general hospital in Bao Loc in 2001, and we have succeeded in transferring these methods. Local staffs can now carry out malaria diagnosis and G6PD deficiency tests by themselves in the hospital laboratory. In addition, we could obtain blood samples of malaria patients and G6PD-deficient persons with informed consent. We further conducted analysis of some genes of malaria parasites [2] and of *G6PD* of deficient persons. We here present the first report describing the genomic analysis of *G6PD* in Vietnamese.

Materials and Methods

This study was approved by the local government

of Lam Dong Province, Vietnam, and the Ethical Committee of Jichi Medical University, Japan. To test G6PD activity, we adopted Fujii's method [3]. This method is useful for testing hundreds of blood samples per day, although it is difficult to distinguish full activity from 50% residual activity in female samples [4]. We therefore concentrated only on males in the study.

Between 1997 and 2003 at Bao Loc General Hospital in Lam Dong Province, located 100 km from the border of Cambodia, blood samples (2 ml) were collected from malaria patients to investigate glucose metabolism [5] and genomic variations of *Plasmodium falciparum* [2]. Beginning in 2001, informed consent was obtained from each blood donor, allowing investigation of not only malaria DNA but also other hereditary abnormalities in each patient's own genes. In 2003, we began to expand the study population to outpatients who consulted the hospital for common diseases because the number of malaria patients was reduced. Blood samples (0.5 ml) were collected only from outpatients consenting to the test of G6PD activity followed by *G6PD* analyses.

Blood samples were stored at -20°C until the G6PD activity test. When more than 100 samples were collected, we conducted the G6PD activity test using Fujii's method [3]. When we found G6PD-deficient samples, we extracted genomic DNA from 0.1 ml of whole blood with a DNA purification kit (Amersham Pharmacia Biotech, Buckinghamshire, UK). Since genomic *G6PD* consists of 13 exons, we prepared primers for these exons [6] as shown in Table 1, amplified each exon by PCR, and read the

Table 1 Origonucleotide primers for PCR

| Exon | Name, Sequence and Location of SENSE primers | Name, Sequence and Location of ANTISENSE primers | Product length (bp) |
|-------|--|--|---------------------|
| 2 | 1a 5'-AGACCCAGAGGAACTCTCAAGAAA (Intron 1) | 2b 5'-TGCAACAATTAGTTGAAAAGCTGA (Intron 2) | 294 |
| 3 & 4 | B1a 5'-TGTCCCCAGCCACTTCT (Intron 2) | B2 5'-CCGAAGCTGGCCATGCTGGG (Intron 4) | 358 |
| 5 | B3 5'-ACACACGGACTCAAAGAGAG (Intron 4) | B2a 5'-TGGTGGGAGCACTGCCTG (Intron 5) | 343 |
| 6 | B5 5'-AGCTCTGATCCTCACTCCCC (Intron 5) | 8b 5'-GGCCAGGTGAGGCTCCTGAGTA (Intron 6) | 285 |
| 7 | 7a 5'-ACATGTGGCCCCTGCACCACA (Intron 6) | B6 5'-GTGACTGCTCTGCCACCCTG (Intron 7) | 242 |
| 8 | 9a 5'-TTGGGGTCCCCATGCCCTTG (Intron 7) | B8 5'-TGCCTCGTCACAGATGGGCC (Intron 8) | 231 |
| 9 | B9a 5'-ACCCAAGGAGCCCATTC (Intron 8) | R2 5'-ACACAGGGCATGCCAGTTCTG (Intron 9) | 276 |
| 10 | E 5'-CTGAGAGAGCTGGTGCT (Intron 9) | B10 5'-CACCATGTGGAGTCCCCCGG (Intron 10) | 342 |
| 11&12 | B11 5'-ACTCCACATGGTGGCAGGCAG (Intron 10) | B12 5'-ATGAGGTAGCTCCACCCTCA (Intron 12) | 397 |
| 13 | RB12 5'-TGAGGGTGGAGCTACCTCAT (Intron 12) | 18a 5'-CGGGGTGGAGGTGGGTGCCCA (Intron 13) | 164 |

DNA sequence (ABI PRISM 310; Applied Biosystems, Foster City, CA, USA). Both strands of each exon were sequenced. To indicate the mutation point, the nucleotide number of the cDNA sequence was used. We also read some introns of *G6PD* because silent mutations had been found on introns of genomic *G6PD*; e.g. nt175C > T on intron 7, nt163C > T on intron 8 [7] and nt93T > C on intron 11 [8].

Results

From 1997 to 2003, we collected 362 blood samples from malaria patients. We added 742 blood samples from outpatients in 2003 and 2004.

Among the samples, 88.9% (981/1104) were from the Kinh. The K'Ho (10.3%, 114/1104), Chauma (0.5%, 5/1104), Nung (0.3%, 3/1104), and Tay (0.1%, 1/1104) were minorities. The incident rates of G6PD deficiency in the Kinh and the K'Ho were 1.9% and 4.4%, respectively (Table 2). Among malaria patients, 12 samples showed G6PD deficiency (3.3%, 12/362). Among samples from outpatients with common diseases, 13 samples showed G6PD deficiency (1.8%, 13/742). There was no significant difference between these 2 incidence rates. Together with those 2 groups, the incidence rate of G6PD deficiency was 2.3% (25/1104) in Lam Dong Province, Vietnam (Table 2).

We analyzed the *G6PD* genes of G6PD-deficient persons. Table 3 shows the frequency of G6PD mutations in 25 samples. G6PD Viangchan (871G > A, 1311C > T and intron 11 nt93T > C) was the most frequent mutation, but 6 other mutations were also detected. In particular, we found a novel G6PD mutation with a single nucleotide change at 352T > C (Fig. 1). Amino-acid change was predicted as 118

Table 2 The incidence of glucose-6-phosphate dehydrogenase (G6PD) deficiency in 5 ethnic groups in Lam Dong Province, Vietnam

| | Ethnic group | | | | | Total |
|-------------------|--------------|------|--------|------|-----|-------|
| | Kinh | K'Ho | Chauma | Nung | Tay | |
| No. of sample | 981 | 114 | 5 | 3 | 1 | 1,104 |
| No. of deficiency | 19 | 5 | 0 | 1 | 0 | 25 |
| Incidence (%) | 1.9% | 4.4% | | | | 2.3% |

Table 3 Glucose-6-phosphate dehydrogenase (G6PD) variants in 3 ethnic groups in Lam Dong Province, Vietnam

| Nucleotide change | Amino acid change | Name of variant | Ethnic group | | | |
|-------------------|-------------------|-----------------|--------------|------|------|-------|
| | | | Kinh | K'Ho | Nung | Total |
| 95A > G | 32H > R | Gaohe | 1 | 0 | 1 | 2 |
| 352T > C | 118Y > H | Bao Loc* | 1 | 0 | 0 | 1 |
| 392G > T | 131G > V | Quing Yuan | 1 | 0 | 0 | 1 |
| 871G > A | 291V > M | Viangchan** | 6 | 5 | 0 | 11 |
| 1360C > T | 454R > C | Union*** | 2 | 0 | 0 | 2 |
| 1376G > T | 459R > L | Canton | 5 | 0 | 0 | 5 |
| 1388G > A | 463R > H | Kaiping | 3 | 0 | 0 | 3 |
| Total | | | 19 | 5 | 1 | 25 |

* New variant being reported in the present study.

** In each case, 2 more nucleotide changes, a nonsense mutation of 1311C > T on exon 11 and a mutation of nt93T > C on intron 11, were confirmed. Note: There is a variant named G6PD Jammu, which has a change of 871G > A, but 1311C on exon 11 and nt93T on intron 11 are the wild type [33].

*** In each case, one more nucleotide, 99A in the exon 2, was confirmed. Note: There is a variant named G6PD Honiara, which has double mutations of 99A > G and 1360C > T [29].




Fig. 1 A, A part of the DNA sequence on exon 5 of a male with glucose-6-phosphate dehydrogenase (G6PD) Bao Loc, hemizygote; B, A part of the DNA sequence on exon 5 of a normal control. Arrows show 352T > C and 352T.

Tyr > His. G6PD activity was estimated at less than 10% residual activity. There was no other mutation detected in his *G6PD*. The subject was 14 months old and had been brought to the hospital with an acute pharyngitis. The hemoglobin level was relatively low (9.3 g/dl), but he did not have chronic hemolytic anemia. Thus, we classified his condition as Class II, less than 10% residual activity without chronic hemolytic anemia, as defined by the WHO working group [9]. We named this new mutation G6PD Bao Loc (GenBank database accession number: DQ839546). All of the other six G6PD mutations found during this work have also been categorized as Class II.

Discussion

Glucose-6-phosphate dehydrogenase (G6PD) deficiency is one of the most frequent hereditary disorders in the world. Since the G6PD gene is located on the X chromosome, G6PD activity is drastically reduced (less than 10% residual activity) in males with G6PD abnormality. In females, only homozygous persons show complete G6PD deficiency, and G6PD activities vary greatly in heterozygous persons. The G6PD gene consists of 13 exons. The size of the open reading frame is 1,545 base pairs [10]. Most G6PD deficiency is caused by a single nucleotide change, resulting in 1 amino-acid change. One hundred forty mutations causing G6PD deficiency have been reported thus far [11].

The distribution of G6PD mutations in Southeast Asian countries differs in each country. For example, in Myanmar, more than 90% of G6PD-deficient cases are G6PD Mahidol (487G > A), and there are no cases of G6PD Viangchan (871G > A, 1311C > T, and intron 11 nt93T > C) [12]. In contrast with Myanmar, more than 90% of G6PD-deficient cases in Cambodia are G6PD Viangchan, and there are no cases of G6PD Mahidol [13, 14]. When we investigated G6PD deficiency in Indonesia, we found 7 different G6PD mutations even on a small island [15, 16]. Other groups have reported finding several G6PD variants in the Thai population [17], Malaysian population [18], and Chinese population [19]. In terms of G6PD-deficient people, Myanmar and Cambodian people are derived from homogenous ancestries, but Indonesian, Thai, Malaysian, and Chinese people are from various

ancestries.

In Vietnam, the frequency of G6PD deficiency is reported to be 1.4-4.1% [20]. Thus, it is likely that the incidence of G6PD deficiency was approximately 2.3% in our present study. However, if we compare the incidence rates in terms of malaria endemic areas, the rate (2.3%) is lower than those in other Southeast Asian countries. According to our previous observation, the rates of G6PD deficiency are 10.5% (45/430) in Myanmar, 3.9% (25/648) in Indonesia [21], and 8.1% (29/360) in Cambodia [14]. The incidence of G6PD deficiency and endemicity of malaria are thought to correlate with each other [22]. There is a hypothesis that G6PD-deficient red blood cells are more tolerant of malaria parasites of *Plasmodium falciparum* [23]. Places in Myanmar, Indonesia, and Cambodia where we examined G6PD activity are all malaria endemic areas. For instance, positive rates of malaria are 45.8% (298/650) in Myanmar, 43.4% (308/709) in Indonesia [21], and 24.9% (167/670) in Cambodia [14]. Compared to these high rates, those in our observation area in Lam Dong Province are not as high. We have found similar results in North Sumatra, Indonesia. The incidence of G6PD deficiency was 4.6% (21/458) in malaria endemic areas, but was 0.9% (1/110) in non-malaria endemic areas [24].

Vietnamese are composed of more than 50 ethnic groups. The Kinh is the largest ethnic group, accounting for approximately 90% of the Vietnamese population [25]. In the present study, the tendency was similar because 88.9% of the blood samples were from ethnic Kinh. Even in the same ethnic group, we found 7 different mutations in 19 persons with G6PD deficiency. G6PD Viangchan was the most frequent mutation in the Kinh population. As shown in Fig. 2, this variant is very common in Laos [26] and Cambodia [13, 14], and it may be from the Khmer, which is the largest ethnic group in Southeast Asian countries.

Furthermore, G6PD Gaohe (95A > G), G6PD Quing Yuan (392G > T), G6PD Canton (1376G > T), and G6PD Kaiping (1388G > A) were also detected in the Kinh population, and their total frequency reached 52.6% (10/19). These variants are commonly found in Chinese [19]. It is well known that Vietnam has had a strong historical connection with China for 2,000 years. Until recently, marriages

between the Kinh and Chinese have not been rare. Thus, it is easily estimated that the G6PD variant of Chinese origins were introduced into the Kinh population and shared with G6PD Viangchan, reflecting another old ancestral source, for Southeast Asian populations.

G6PD Union is dominantly found in countries in the Pacific Ocean such as the Philippines [27], Papua New Guinea [28], Solomon Islands [29], and the Vanuatu archipelago [30]. Two cases of this variant were also detected among the Kinh. These results suggest that the Kinh ethnic group has several different ancestral sources, including the Khmer, Chinese, and people from the Pacific Ocean.

In contrast, all 5 cases detected in the K'Ho were G6PD Viangchan. In Laos [26] and Cambodia [13], more than 90% of G6PD-deficient cases are G6PD Viangchan, and this tendency is the same in the 2 minorities (the Tum Pun and the Cha Ray) in Cambodia [14]. In Vietnam and Cambodia, these minorities have remained isolated from each other and

have not until recently intermarried with other tribes. Thus we suspect that G6PD Viangchan was introduced in Southeast Asian countries in an old era, more than 1,000 years ago, and the K'Ho, the Khmer, and the Lao were separated after this variant was introduced in their original tribe.

The Kinh people might obtain deferent genes by intermarrying with other populations from China and Oceania. It is interesting to note that there were no Kinh people showing G6PD Mahidol, which is another common G6PD variant distributed in continental Southeast Asian countries of Myanmar, Thailand, and Malaysia. This variant has not yet reached Cambodia, Laos, and Vietnam.

In the present study, we adopted Fujii's method, which makes it difficult to distinguish full activity from 50% residual activity in female samples. However, new rapid G6PD tests are currently being developed to detect samples with 50% residual activity [31, 32]. With these methods, we may detect heterozygous G6PD deficiency among female blood




Fig. 2 Distribution and frequencies of glucose-6-phosphate dehydrogenase (G6PD) variants in Southeast Asian countries. Each number indicates the number of G6PD-deficient cases confirmed by sequence analysis. Data for Myanmar, Laos, and Cambodia are from our previous reports [12, 14, 26]. Data for Thailand [17], Malaysia [18], and China [19] are from other group's reports.

samples. In further studies, we should carry out genetic analysis of female cases of G6PD deficiency.

Acknowledgments. We thank Mrs. Midori Sato for technical assistance. This work was supported by a grant-in-aid for Scientific Research from the Ministry of Education, Culture, Sports, Science, and Technology of Japan (14406026/17406025 to HM and 16590341/17406010 to FK). We also thank the staff of Bao Loc General Hospital for support and good collaboration during the study.

References

- Tantular IS, Iwai K, Lin K, Basuki S, Horie T, Htay HH, Matsuoka H, Marwoto H, Wongsrichanalai C, Dachlan YP, Kojima S, Ishii A and Kawamoto F: Field trials of a rapid test for G6PD deficiency in combination with a rapid diagnosis of malaria. *Trop Med Int Health* (1999) 4: 245-250.
- Jalloh A, van Thien H, Ferreira MU, Ohashi J, Matsuoka H, Kanbe T, Kikuchi A and Kawamoto F: Sequence variation in the T-cell epitopes of the *Plasmodium falciparum* circumsporozoite protein (CSP) among field isolates is temporally stable: a five-year longitudinal study in Southern Vietnam. *J Clin Microbiol* (2006) 44: 1229-1235.
- Fujii H, Takahashi K and Miwa S: A new simple screening method for glucose 6-phosphate dehydrogenase deficiency. *Nippon Ketsueki Gakkai Zasshi (Acta Haematol Jpn)* (1984) 47: 185-188.
- Pujades A, Lewis M, Salvati AM, Miwa S, Fujii H, Zarza R, Alvarez R, Rull E and Corrons JL: Evaluation of the blue formazan spot test for screening glucose 6 phosphate dehydrogenase deficiency. *Int J Hematol* (1999) 69: 234-236.
- van Thien H, Weverling GJ, Ackermans MT, canh Hung N, Ender T, Kager PA and Sauerwein HP: FFAs are not involved in regulation of gluconeogenesis and glycogenolysis in adults with uncomplicated *P. falciparum* malaria. *Am J Physiol Endocrinol Metab* (2004) 287: E609-615.
- Hirono A, Miwa S, Fujii H, Ishida F, Yamada K and Kubota K: Molecular study of eight Japanese cases of glucose-6-phosphate dehydrogenase deficiency by nonradioisotopic single-strand conformation polymorphism analysis. *Blood* (1994) 83: 3363-3368.
- Vulliamy TJ, Othman A, Town M, Nathwani A, Falusi AG, Mason PJ and Luzzatto L: Polymorphic sites in the African population detected by sequence analysis of the glucose-6-phosphate dehydrogenase gene outline the evolution of the variants A and A-. *Proc Natl Acad Sci USA* (1991) 88: 8568-8571.
- Beutler E, Westwood B and Sipe B: A new polymorphic site in the G6PD gene. *Hum Genet* (1992) 89: 485-486.
- WHO working group: Glucose-6-phosphate dehydrogenase deficiency. *Bull World Health Organ* (1989) 67: 601-611.
- Chen EY, Cheng A, Lee A, Kuang WJ, Hillier L, Green P, Schlessinger D, Ciccodicola A and D'Urso M: Sequence of human glucose-6-phosphate dehydrogenase cloned in plasmids and a yeast artificial chromosome. *Genomics* (1991) 10: 792-800.
- Beutler E and Vulliamy TJ: Hematologically important mutations: glucose-6-phosphate dehydrogenase. *Blood Cells Mol Dis* (2002) 28: 93-103.
- Matsuoka H, Wang J, Hirai M, Arai M, Yoshida S, Kobayashi T, Jalloh A, Lin K and Kawamoto F: Glucose-6-phosphate dehydrogenase (G6PD) mutations in Myanmar: G6PD Mahidol (487G > A) is the most common variant in the Myanmar population. *J Hum Genet* (2004) 49: 544-547.
- Louicharoen C and Nuchprayoon I: *G6PD Viangchan* (871G > A) is the most common variant in the Cambodian population. *J Hum Genet* (2005) 50: 448-452.
- Matsuoka H, Nguon C, Kanbe T, Jalloh A, Sato H, Yoshida S, Hirai H, Arai M, Socheat D and Kawamoto F: Glucose-6-phosphate dehydrogenase (G6PD) mutations in Cambodia: *G6PD Viangchan* (871G > A) is the most common variant in the Cambodian population. *J Hum Genet* (2005) 50: 468-472.
- Matsuoka H, Arai M, Yoshida S, Tantular IS, Pusarawati S, Kerong H and Kawamoto F: Five different glucose-6-phosphate dehydrogenase (G6PD) variants found among 11 G6PD-deficient persons in Flores Island, Indonesia. *J Hum Genet* (2003) 48: 541-544.
- Kawamoto F, Matsuoka H, Kanbe T, Tantular IS, Pusarawati S, Kerong HI, Damianus W, Mere D and Dachlan YP: Further investigations of glucose-6-phosphate dehydrogenase variants in Flores Island, eastern Indonesia. *J Hum Genet* (2006) 51: 952-957.
- Nuchprayoon I, Sanpavat S and Nuchprayoon S: Glucose-6-phosphate dehydrogenase (G6PD) mutations in Thailand: *G6PD Viangchan* (871G > A) is the most common deficiency variant in the Thai population. *Hum Mutat* (2002) 19: 185.
- Ainon O, Yu YH, Amir Muhriz AL, Boo NY, Cheong SK and Hamidah NH: Glucose-6-phosphate dehydrogenase (G6PD) variants in Malaysian Malays. *Hum Mutat* (2003) 21: 101.
- Jiang W, Yu G, Liu P, Geng Q, Chen L, Lin Q, Ren X, Ye W, He Y, Guo Y, Duan S, Wen J, Li H, Qi Y, Jiang C, Zheng Y, Liu C, Si E, Zhang Q, Tian Q and Du C: Structure and function of glucose-6-phosphate dehydrogenase-deficient variants in Chinese population. *Hum Genet* (2006) 119: 463-478.
- Panich V: *G6PD variants in Southern Asian populations; in Glucose 6-phosphate dehydrogenase*, Yoshida A and Beutler E eds, Academic Press, New York (1986) pp 195-241.
- Jalloh A, Tantular IS, Pusarawati S, Kawilarang AP, Kerong H, Lin K, Ferreira MU, Matsuoka H, Arai M, Kita K and Kawamoto F: Rapid epidemiologic assessment of glucose-6-phosphate dehydrogenase (G6PD) deficiency in malaria-endemic areas in Southeast Asia using a novel diagnostic kit. *Trop Med Int Health* (2004) 9: 615-623.
- Ruwende C, Khoo SC, Snow RW, Yates SN, Kwiatkowski D, Gupta S, Warn P, Allsopp CE, Gilbert SC, Peschu N, Newbold CI, Greenwood BM, Marsh K and Hill AVS: Natural selection of hemi- and heterozygotes for G6PD deficiency in Africa by resistance to severe malaria. *Nature* (1995) 376: 246-249.
- Luzzatto L, Usanga FA and Reddy S: Glucose-6-phosphate dehydrogenase deficient red cells: resistance to infection by malarial parasites. *Science* (1969) 164: 839-842.
- Matsuoka H, Ishii A, Panjaitan W and Sudiranto R: Malaria and glucose-6-phosphate dehydrogenase deficiency in North Sumatra, Indonesia. *Southeast Asian J Trop Med Public Health* (1986) 17: 530-536.
- Dien K: *Population and ethno-demography in Vietnam*. Silksworm Books, Chiang Mai, Thailand (2002) pp 1-207.
- Iwai K, Hirono A, Matsuoka H, Kawamoto F, Horie T, Lin K, Tantular IS, Dachlan YP, Notopuro H, Hidayah NI, Salim A M, Fujii H, Miwa S and Ishii A: Distribution of glucose-6-phosphate dehydrogenase mutations in Southeast Asia. *Human Genetics* (2001) 108: 445-449.
- Silao CL, Shirakawa T, Nishiyama K, Padilla C and Matsuo M: Molecular basis of glucose-6-phosphate dehydrogenase deficiency among Filipinos. *Pediatr Int* (1999) 41: 138-141.

August 2007

G6PD Variants in Vietnamese 219

28. Wagner G, Bhatia K and Board P: Glucose-6-phosphate dehydrogenase deficiency mutations in Papua New Guinea. *Hum Biol* (1996) 68: 383–394.
29. Hirono A, Ishii A, Kere N, Fujii H, Hirono K and Miwa S: Molecular analysis of glucose-6-phosphate dehydrogenase variants in the Solomon Islands. *Am J Hum Genet* (1995) 56: 1243–1245.
30. Ganczakowski M, Town M, Bowden DK, Vulliamy TJ, Kaneko A, Clegg JB, Weatherall DJ and Luzzatto L: Multiple glucose 6-phosphate dehydrogenase-deficient variants correlate with malaria endemicity in the Vanuatu archipelago (southwestern Pacific). *Am J Hum Genet* (1995) 56: 294–301.
31. Tantular IS and Kawamoto F: An improved, simple screening method for detection of glucose-6-phosphate dehydrogenase deficiency. *Trop Med Int Health* (2003) 8: 569–574.
32. Arai M, Kosuge K, Kawamoto F and Matsuoka H: Reactivity of blood samples spotted onto filter papers in the WST-8 method for screening of G6PD deficiency. *Acta Med Okayama* (2006) 60: 127–134.
33. Beutler E, Westwood B and Kuhl W: Definition of the mutations of G6PD Wayne, G6PD Viangchan, G6PD Jammu, and G6PD 'LeJeune'. *Acta Haematol* (1991) 86: 179–182.