

Acta Medica Okayama

Volume 61, Issue 1

2007

Article 3

FEBRUARY 2007

Biochemical characterization of reactive nitrogen species by eosinophil peroxidase in tyrosine nitration

Kei Takemoto*

Keiki Ogino[†]

Da-Hong Wang[‡]

Tomoko Takigawa**

Carmen M. Kurosawa^{††}

Yasuhiro Kambayashi^{‡‡}

Yuri Hibino[§]

Yoshiaki Hitomi[¶]

Hiroshi Ichimura^{||}

* Kanazawa University,

[†] Okayama University,

[‡] Okayama University,

** Okayama University,

^{††} Okayama University,

^{‡‡} Kanazawa University,

[§] Kanazawa University,

[¶] Kanazawa University,

^{||} Kanazawa University,

Biochemical characterization of reactive nitrogen species by eosinophil peroxidase in tyrosine nitration*

Kei Takemoto, Keiki Ogino, Da-Hong Wang, Tomoko Takigawa, Carmen M. Kurosawa, Yasuhiro Kambayashi, Yuri Hibino, Yoshiaki Hitomi, and Hiroshi Ichimura

Abstract

It is well known that eosinophils are involved in tyrosine nitration. In this study, we evaluated tyrosine nitration by rat eosinophils isolated from peritoneal fluid and constituent eosinophils in the stomach. Rat peritoneal eosinophils activated with 1 μ M phorbol myristate acetate (PMA) and 50 μ M NO_2 showed immunostaining for nitrotyrosine only in smaller cells, despite the fact that eosinophils are capable of producing superoxide ($\text{O}_2^{\cdot-}$). Free tyrosine nitrating capacity after incubation with PMA and NO_2 was 4-fold higher in eosinophils than in neutrophils. Catalase and α -tocopherol inhibited free tyrosine nitration by reactive nitrogen species from eosinophils but not that by peroxynitrite. Superoxide dismutase augmented free tyrosine nitration by activated eosinophils and peroxynitrite. The concentration of nitric oxide released from eosinophils was relatively low (0.32 μ M/106 cells/h) and did not contribute to the formation of nitrotyrosine. On the other hand, most constituent eosinophils in the rat stomach stimulated by PMA and NO_2 showed tyrosine nitration capacity. These results suggest that intact cells other than apoptotic-like eosinophils eluted in the intraperitoneal cavity could not generate reactive species responsible for nitration by a peroxidase-dependent mechanism. In contrast, normal eosinophils in the stomach were capable of nitration, suggesting that the characteristics of eosinophils in gastric mucosa are different from those eluted in the peritoneal cavity.

KEYWORDS: eosinophil peroxidase, reactive nitrogen species, nitrotyrosine

*PMID: 17332838 [PubMed - in process]

Original Article

Biochemical Characterization of Reactive Nitrogen Species by Eosinophil Peroxidase in Tyrosine Nitration

Kei Takemoto^a, Keiki Ogino^{b*}, Da-Hong Wang^b,
Tomoko Takigawa^b, Carmen M. Kurosawa^b, Yasuhiro Kambayashi^c,
Yuri Hibino^c, Yoshiaki Hitomi^c, and Hiroshi Ichimura^a

Departments of ^aViral Infection and International Health, ^cEnvironmental and Preventive Medicine,
Kanazawa University Graduate School of Medical Science, Kanazawa 920-0934, Japan,
^bDepartment of Public Health, Okayama University Graduate School of Medicine,
Dentistry and Pharmaceutical Sciences, Okayama 700-8558, Japan

It is well known that eosinophils are involved in tyrosine nitration. In this study, we evaluated tyrosine nitration by rat eosinophils isolated from peritoneal fluid and constituent eosinophils in the stomach. Rat peritoneal eosinophils activated with 1 μM phorbol myristate acetate (PMA) and 50 μM NO_2^- showed immunostaining for nitrotyrosine only in smaller cells, despite the fact that eosinophils are capable of producing superoxide (O_2^-). Free tyrosine nitrating capacity after incubation with PMA and NO_2^- was 4-fold higher in eosinophils than in neutrophils. Catalase and α - and γ -tocopherol inhibited free tyrosine nitration by reactive nitrogen species from eosinophils but not that by peroxynitrite. Superoxide dismutase augmented free tyrosine nitration by activated eosinophils and peroxynitrite. The concentration of nitric oxide released from eosinophils was relatively low (0.32 $\mu\text{M}/10^6$ cells/h) and did not contribute to the formation of nitrotyrosine. On the other hand, most constituent eosinophils constituent in the rat stomach stimulated by PMA and NO_2^- showed tyrosine nitration capacity. These results suggest that intact cells other than apoptotic-like eosinophils eluted in the intraperitoneal cavity could not generate reactive species responsible for nitration by a peroxidase-dependent mechanism. In contrast, normal eosinophils in the stomach were capable of nitration, suggesting that the characteristics of eosinophils in gastric mucosa are different from those eluted in the peritoneal cavity.

Key words: eosinophil peroxidase, reactive nitrogen species, nitrotyrosine

Eosinophils in biological systems have been suggested to be important in the defense against parasitic helminths [1, 2]. Eosinophil recruitment is a characteristic feature of asthma and other allergic inflammatory diseases [3-5]. Inappropriate accumu-

lation of eosinophils and eosinophil-derived granules in the tissue and sputum are not only considered as biomarkers for monitoring the severity of asthmatic events [3-8] but also suggested to evoke cellular and tissue damage [3-6]. Eosinophils possess numerous highly basic and toxic granules that are released upon activation or during cell necrosis. They are also equipped with the enzymes that inflict oxidative damage upon biological targets [3, 6]. The respira-

Received November 29, 2006; accepted December 19, 2006.
*Corresponding author. Phone: +81-86-235-7184; Fax: +81-86-226-0715
E-mail: kogino@md.okayama-u.ac.jp (K. Ogino)

tory burst occurs in activated eosinophils where O_2^- and its dismutation product, H_2O_2 , are formed [1, 9].

The formation of peroxynitrite ($ONOO^-$) and peroxidase-catalyzed reactive nitrogen species is implicated in eosinophils of the lung in asthma patients [10, 11]. $ONOO^-$ is a potent oxidant formed by a rapid reaction of NO and O_2^- [12]. Inducible nitric oxide synthase (iNOS), which generates NO from L-arginine, is expressed in human circulating eosinophils [13] and rat peritoneal eosinophils [14]. Simultaneous production of O_2^- and NO in eosinophils enables the production of $ONOO^-$. Peroxynitrite can react with tyrosine residues at a physiologically relevant pH to yield nitrotyrosine [15, 16]. On the basis of nitrotyrosine formation, $ONOO^-$ has been implicated in the pathogenesis of a variety of diseases including neurodegenerative disease [17], atherosclerosis [18], allergic asthma [10, 11], and inflammatory gastro-intestinal diseases [19, 20].

Peroxidase-catalyzed formation of reactive nitrogen species using NO_2^- and H_2O_2 as substrates has been demonstrated in purified myeloperoxidase (MPO) [21, 22] and eosinophil peroxidase (EPO) [23]. In healthy human subjects, NO_2^- can be detected at levels of 0.5–3.6 μM in plasma [24]. Increased levels of NO_2^- have been reported in a variety of pathological conditions [25, 26]. It is likely that eosinophils generate reactive nitrogen species, which contributes to tyrosine nitration, if EPO utilizes H_2O_2 as a cosubstrate to oxidize NO_2^- . However, there is little evidence for the production of reactive nitrogen species from eosinophils other than $ONOO^-$ that contributes to tyrosine nitration. Therefore, in the present study we aimed at clarifying the precise mechanisms for generation of nitrotyrosine by rat eosinophils.

Materials and Methods

Animals. Male Wistar (Kwl) rats weighing 250–300 g were purchased from Kiwa Laboratory Animals. The animal experiments were carried out according to protocols approved by the Kanazawa University Committee on Laboratory Animals.

Isolation of rat peritoneal eosinophils. To elicit an eosinophil rich exudate, we injected rats with 1 ml of horse serum i.p. 3 times at 2-day intervals and performed a peritoneal lavage 24 h after the final injection. After centrifugation at $200 \times g$ for

5 min, cells were suspended in 3 ml of Ca^{2+} Mg^{2+} -free Hanks' balanced salt solution (HBSS) containing 10 % rat plasma and 0.01 % ethylenediamine tetraacetic acid (EDTA), and layered on a discontinuous Percoll (Amersham Pharmacia Biotech, Uppsala, Sweden) gradient, consisting of 45%, 55%, 60%, 65% Percoll (based on 100% = isotonic Percoll, made up with $10 \times$ concentrated HBSS), each layer containing 10% rat plasma and 0.01% EDTA. The gradient was centrifuged at $400 \times g$ for 30 min. The purest eosinophil fraction was recovered at 65% Percoll interphase, yielding $2-4 \times 10^6$ eosinophils per rat. The purity of the cell population $> 95\%$ was confirmed by cytocentrifuge smears stained with hematoxylin and eosin and Luxol-fast blue (LFB) [27]. Contaminating cells were mainly macrophages with $< 1\%$ neutrophils. Viability of the cells by trypan blue exclusion was $> 99\%$.

Isolation of rat peritoneal neutrophils. Rat neutrophils were prepared from intraperitoneal fluid [28]. The rats were injected i.p. with 20 ml of sterile 12% sodium caseinate in physiological saline. Twelve h later, the peritoneal lavage was performed. After centrifugation at $200 \times g$ for 5 min, cells were suspended in 1 ml of physiological saline, and we then induced hypotonic lysis of contaminated erythrocytes. The cell suspension in Ca^{2+} Mg^{2+} -free HBSS was layered on sodium metrizoate Ficoll ($d = 1.09$) to separate neutrophils from macrophages and lymphocytes, and centrifuged at $1,000 \times g$ for 30 min. The purity of the cell population $> 99\%$ was determined by Wright's staining of cytocentrifuge smears. Cell viability by Trypan blue exclusion was $> 97\%$.

Nitration of endogenous protein in eosinophils and neutrophils. We used both unfixed and fixed cells to detect nitrotyrosine formation by eosinophil peroxidase and myeloperoxidase. In the study of unfixed cells, eosinophils (5×10^5) and neutrophils (5×10^5) were incubated with 50 μM NO_2^- and 1 μM phorbol myristate acetate (PMA) in 250 μl of HBSS containing 0.1 mM diethylenetriamine pentaacetic acid (DTPA) at 37 °C for 1 h. After incubation, cells were cytocentrifuged onto the slides and fixed in 10% neutral-buffered formalin for 20 min.

In the study of fixed cells, cells were cytocentrifuged onto the slides first and then fixed with 10% neutral-buffered formalin for 5 min and washed with PBS 3 times for 5 min. Next, slides were incubated

with the combination of NO_2^- (1, 5, and 1,000 μM) and H_2O_2 (1, 5, and 1,000 μM) in 250 μl of phosphate buffered saline (PBS) commonly containing 0.1 mM DTPA, 1 mM phenylmethylsulfonyl fluoride (Wako Pure Chemical Industries, Osaka, Japan), 1 $\mu\text{g}/\text{ml}$ leupeptine (Sigma-Aldrich, St. Louis, MO, USA), and 1 $\mu\text{g}/\text{ml}$ aprotinin (Roche Diagnostics, Basel, Switzerland) at 37 °C for 30 min. After incubation, the slides were refixed in 10% neutral-buffered formalin for 20 min.

Immunocytochemical study for nitrotyrosine. The slides of fixed eosinophils and neutrophils were treated with 5% H_2O_2 in methanol for 20 min to inactivate endogenous peroxidase. After washing with PBS for 10 min 3 times, slides were mounted with 2% normal goat serum for 1 h in PBS. Polyclonal anti-nitrotyrosine antibodies (rabbit immunofluorescence purified IgG, Upstate Biotechnology, Inc., Lake Placid, NY, USA) or monoclonal anti-nitrotyrosine antibodies (mouse IgG, Upstate Biotechnology) was diluted 800- or 100-fold with PBS, respectively, and layered on the slides overnight at 4 °C. Slides were washed 3 times with PBS and incubated with a second antibody, horseradish peroxidase polymer-conjugated Envision plus (DAKO, Denmark) for 30 min at room temperature. After washing 3 times with PBS, peroxidase activity was localized by 3,3'-diaminobenzidine tetrahydrochloride and 0.01% H_2O_2 , and slides were counterstained with hematoxylin or LFB.

The specificity of immunocytochemistry for nitrotyrosine was confirmed by preincubation of antibodies with 10 mM 3-nitrotyrosine for 2 h or by pretreatment of the slides with 1 mg/ml dithionite in PBS for 1 h. A negative control study was performed using non-immune rabbit or mouse immunoglobulin. Light microscopic observations were carried out with an Olympus IX-70.

Superoxide generation from eosinophils and neutrophils. Superoxide release was measured following the superoxide dismutase (SOD)-inhibitable reduction of ferricytochrome c at 550 nm in a single spectrophotometer by a modification of the procedure of Babior *et al.* [29]. Eosinophils or neutrophils (1×10^6) were incubated at 37 °C for 1 h in 1 ml of HBSS with 0.1 mM ferricytochrome c, 1 μM PMA, and 50 μM NO_2^- . Before incubation, 50 $\mu\text{g}/\text{ml}$ of SOD was added as a reference.

Histochemical detection of superoxide generation from eosinophils. Superoxide generation was detected histochemically by a modification of the nitro blue tetrazolium (NBT) reduction method [30]. Eosinophils (1×10^6) were incubated at 37 °C for 1 h in 1 ml of HBSS with 500 $\mu\text{g}/\text{ml}$ NBT, and the addition of 1 μM PMA or 1 μM PMA and 50 μM NO_2^- . After incubation, cells were cytocentrifuged on slides and fixed with 10% neutral-buffered formalin, and we then performed an immunohistochemical study for nitrotyrosine.

Western blot analysis of nitrated proteins. Eosinophils (1×10^6) or neutrophils (1×10^6) were incubated at 37 °C for 1 h in HBSS (500 μl) with PMA (1 μM), NO_2^- (0–50 μM), H_2O_2 (50 μM), SOD (50 $\mu\text{g}/\text{ml}$), catalase (30 $\mu\text{g}/\text{ml}$), 1 mg/ml bovine serum albumin (BSA), or the combination of these reagents. After incubation, cells were immediately centrifuged and the pellets were solubilized by the addition of 50 μl non-reducing Laemmli sample buffer and then boiled for 5 min. Samples (40 μg protein) were subjected to 4–20% sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and transferred onto nitrocellulose membranes.

The membranes were blocked by overnight incubation at 4 °C with 3% (w/v) nonfat dried milk in 20 mM Tris-HCl, pH 7.7, containing 137 mM NaCl and 0.1% (w/v) Tween-20 (buffer A). The membranes were washed twice for 5 min with buffer A, following incubation for 2 h with a polyclonal anti-nitrotyrosine antibody (IgG, Upstate Biotechnology, Lake Placid, NY, USA) diluted 1,000-fold in buffer A containing 1.5% (w/v) nonfat dried milk. After washing 3 times for 20 min with buffer A, the membranes were incubated for 1 h with a horseradish peroxidase (HRP)-labeled anti-mouse IgG antibody, which was diluted 2,000-fold in buffer A containing 1.5% nonfat dried milk. The membranes were washed three times for 20 min with buffer A, and peroxidase activity was detected on X-ray film with enhanced chemiluminescence (NEN Life Science Products, Boston, MA, USA). The specificity of nitrated proteins was clarified by pre-incubation of anti-nitrotyrosine antibody with 10 mM 3-nitrotyrosine (Aldrich Chemical Co., Milwaukee, WI, USA) for 2 h.

Nitration of free tyrosine. We investigated

free tyrosine nitration by eosinophils and neutrophils using the following incubation mixture. Eosinophils or neutrophils (2.5×10^5) were incubated at 37 °C for 1 h in 250 μ l of HBSS with 1 mM DL-tyrosine, and the addition of 1 μ M PMA, 0–50 μ M sodium nitrite (NO_2^-), 0–50 μ M H_2O_2 , or the combination of these reagents.

To clarify a characteristic of reactive nitrogen species concerning tyrosine nitration from eosinophils, we added one of various antioxidants, 50 μ g/ml superoxide dismutase (SOD), 30 μ g/ml catalase, 1 mM methionine, 50 mM mannitol, 50 mM dimethylsulfoxide (DMSO), 1 mM urate, 1 mM cysteine, 69 mM ethanol, 0.1 mM α -tocopherol (in 69 mM ethanol), or 0.1 mM γ -tocopherol (in 69 mM ethanol), to the incubation mixtures described above. The concentrations of PMA and NO_2^- in the mixture used for the antioxidative study were 1 μ M and 50 μ M, respectively. In some experiments, 0.1 mM peroxytrinitrite (synthesized from NO_2^- and acidified H_2O_2 [31], Dojindo Laboratories, Kumamoto, Japan) was used instead of the combination of PMA and NO_2^- .

The incubation was terminated by centrifugation (5,000 rpm) on ULTRAFREE-MC (10 kDa molecular weight cut-off) (Millipore Co., Billerica, MA, USA) to remove proteins. Tyrosine and its nitration products in filtrates were analyzed by high-performance liquid chromatography (HPLC) using a 5- μ m Spherisorb ODS-2RP-18 column, with 93% 50 mM potassium phosphate (pH 3.0), 7% methanol as the mobile phase at 1 ml/min, and UV detection at 274 nm [32]. Identification and quantitation of nitrotyrosine were performed with excess sodium hydrogensulfite, which reduces nitrotyrosine to aminotyrosine [33], and by comparison with external standards.

Gastric mucosal slice preparation and incubation. Rat gastric mucosal slices were prepared and incubated by a modification of the method described previously [34]. Briefly, rats were killed by cervical dislocation after a slight ether anesthesia. The stomach was then removed and placed in ice-cold oxygenated (95% $\text{O}_2/5\%$ CO_2) Krebs-Henseleit buffer. Gastric mucosal slices of about 0.3-mm in thickness were cut with a microslicer and put into 5-ml vials containing 3 ml of Waymouth's culture medium (pH 7.4) supplemented with 10% heat-inactivated fetal bovine serum, 20 mM HEPES, 1 μ g insulin/ml, 84 μ g gentamycin/ml, and 25 mM glucose. The vials

were closed with caps and gassed with a mixture of 95% $\text{O}_2/5\%$ CO_2 . The slices were incubated at 37 °C with shaking (80 strokes/min) for 1 h prior to experiments. Then, sections were incubated with or without 1 μ M of PMA and 1 μ M NO_2^- in Krebs-Henseleit buffer at 37 °C for 1 h. The resulting sections were fixed with 10% neutral buffered formalin and then embedded in paraffin, and serial sections were retrieved to investigate the localization of eosinophils and nitrotyrosine.

Nitric oxide production from eosinophils and neutrophils. Nitric oxide (NO) was measured as NO_2^- by the reaction of NO_2^- and 2,3-diaminonaphthalene [35]. Eosinophils or neutrophils (1×10^6) were incubated with 0.5 mM L-arginine, 12 μ M S-methylisothiourea, or a combination of each reagent at 37 °C for 3 h in 500 μ l of HBSS. After incubation, cells were centrifuged at 3,000 rpm for 5 min, and then the supernatants were incubated with 50 μ g 2,3-diaminonaphthalene dissolved in 1 ml of 0.62 N hydrochloric acid for 10 min, followed by the addition of 5 ml of 2.8 N sodium hydroxide. Incubation mixtures (100 μ l) were dissolved with 4 ml of distilled water and measured for fluorescence detection (excitation, 365 nm and emission, 450 nm).

Immunocytochemical study for iNOS. The slides of cytocentrifuged eosinophils and neutrophils were fixed with ice-cold methanol for 5 min and incubated with non-immune mouse IgG2a to prevent non-specific binding for 20 min, and then incubated with fluorescein isothiocyanate (FITC)-conjugated mouse anti-iNOS monoclonal antibodies (IgG2a, Transduction Laboratories, Lexington, KY, USA) diluted 25-fold with PBS for 2 h. After 3 times washing with PBS for 5 min, slides were mounted in Immunon™ (Shandon, Pittsburgh, PA, USA) and sealed with clear nail varnish. A control study was performed using FITC-conjugated non-immune IgG2a (DAKO). Fluorescence microscopic observations were analyzed using an Olympus IX-70 equipped with an IX-FLA fluorescence observation attachment.

Reverse transcription polymerase chain reaction (RT-PCR). Total RNA was isolated from eosinophils or neutrophils (1×10^6) using an Isogen RNA reagent (Wako), and 0.5 μ g of the resulting RNA was treated with DNase I (Takara, Kyoto, Japan) for 1 h at 37 °C. DNase I was inactivated for 5 min at 90 °C. First-strand cDNA synthe-

sis was performed using 0.5 μg of each RNA sample, 0.125 μM random primers, and 0.25 U avian myeloblastosis viral reverse transcriptase (RT) (Takara) in a 20- μl reaction volume. Contamination with DNA was excluded by performing polymerase chain reaction (PCR) from templates incubated without RT. The PCR primers used were as follows: iNOS (5'-GATCAATAACCTGAACCCG-3', and antisense primer 5'-GCCCTTTTTTGCTCCATAGG-3', 578-bp fragment), glyceraldehyde-3-phosphate dehydrogenase (G3PDH) (5'-ACCACAGTCCATGCCATCAC-3', and antisense primer 5'-TCCACCACCCTGTTGCTGTA-3', 452-bp fragment).

The PCR reaction mixture (100 μl) contained 2.5 mM MgCl_2 , 0.2 mM dNTP, 1 μM primers, and 2.5 U Taq DNA polymerase. Samples were amplified for 30 or 40 cycles for iNOS and 30 cycles for G3PDH by 30 sec denaturation at 94 $^\circ\text{C}$, 30 sec annealing at 56 $^\circ\text{C}$, and 1.5 min elongation at 72 $^\circ\text{C}$ in a Takara PCR thermal cycler (Takara). The PCR reactions were then visualized on a 2% agarose gel (Takara) containing 5 $\mu\text{g}/\text{ml}$ of ethidium bromide. Steady-state expression of G3PDH mRNA was used to control for equal product loading.

Statistical analysis. All values are expressed as means \pm SD of several experiments. The significance of the difference among groups was analyzed by one-way analysis of variance following *post hoc* multiple-comparisons, and $p < 0.05$ was considered significant.

Results

Immunocytochemistry for nitrotyrosine in peritoneal eosinophils and neutrophils. The specificity of eosinophils was confirmed by staining with hematoxylin and eosin (Fig. 1A). Incubation of eosinophils with 1 μM of PMA and 50 μM of NO_2^- showed prominent immunostaining for nitrotyrosine in a few smaller cells or cells containing a round nucleus (Fig. 1B). Almost all eosinophils changed the morphology to small nuclei within 5 h of being isolated (data not shown). Incubation with 5 μM NO_2^- and 5 μM H_2O_2 showed diffuse cytoplasmic staining (Fig. 1C), while incubation with 1 μM NO_2^- and 1 μM H_2O_2 showed focal staining in the site of exocytosis (Fig. 1D, indicated as arrows). Significant disruption of eosinophils was observed following expo-

sure to 1 mM of NO_2^- and 1 mM of H_2O_2 (Fig. 1E). Pretreatment of anti-nitrotyrosine antibody with 10 mM of nitrotyrosine abolished the immunostaining for nitrotyrosine by PMA and NO_2^- (data not shown). Incubation of neutrophils with 1–5 μM NO_2^- and 1–5 μM H_2O_2 did not show immunoreactivity for nitrotyrosine (Fig. 1F).

Superoxide generation from activated eosinophils and neutrophils. Superoxide generation from eosinophils or neutrophils activated by PMA was 60.4 ± 1.3 or 57.4 ± 1.3 nmol/ 1×10^6 /h, respectively.

Histochemical detection of superoxide generation. Superoxide generation from activated eosinophils by PMA was detected using the histochemical NBT reduction method. After incubation of eosinophils with 1 μM of PMA, blue granules were observed in the eosinophil fraction (Fig. 2A). A phase-contrast study of the same slides revealed that NBT-reducible granules were located in most eosinophils (Fig. 2B). The staining of NBT in eosinophils was not changed by the incubation of cells with 1 μM PMA and 50 μM NO_2^- .

Western blot analysis. Protein tyrosine nitration in eosinophils and neutrophils was analyzed by SDS-PAGE and Western blotting after incubation of cells with a combination of agents (Fig. 3A). Incubation of eosinophils with PMA and 50 μM NO_2^- or 50 μM NO_2^- and 50 μM H_2O_2 showed many nitrated proteins (lane 3, lane 7). The staining intensity of nitrated protein bands was augmented by SOD (lane 5) and diminished by catalase (lane 6). Bovine serum albumin in the incubation of eosinophils with PMA and 50 μM NO_2^- did not show the staining band for nitrotyrosine (lane 8). Neutrophils incubated with PMA and 50 μM NO_2^- or 50 μM NO_2^- and 50 μM H_2O_2 did not show marked bands of nitration proteins (lane 9, lane 10). The nitrated proteins were abolished by pretreatment of antibodies with 3-nitrotyrosine (Fig. 3B).

Free tyrosine nitration. We investigated free tyrosine nitration by eosinophils or neutrophils by incubating cells with a combination of PMA and various concentrations of NO_2^- . Nitration of free tyrosine by eosinophils was 3.5-, 4.8-, and 7.3-fold higher than that by neutrophils when these cells were incubated with PMA and 5 μM NO_2^- , PMA and 50 μM NO_2^- , or 50 μM NO_2^- and 50 μM H_2O_2 ,




Fig. 1 Histocytochemical purity of rat peritoneal eosinophils and immunocytochemistry of eosinophils for nitrotyrosine. The purity of peritoneal eosinophils is demonstrated with hematoxylin and eosin staining (A) and Luxol-fast blue staining (B). Staining for nitrotyrosine was detected in small-sized eosinophils incubated with $1\ \mu\text{M}$ PMA and $50\ \mu\text{M}$ NO_2^- (B). Incubation of fixed eosinophils with $5\ \mu\text{M}$ H_2O_2 and $5\ \mu\text{M}$ NO_2^- shows immunostaining of the entire cytoplasm of almost all eosinophils (C). When the slide was incubated with $1\ \mu\text{M}$ H_2O_2 and $1\ \mu\text{M}$ NO_2^- after fixation with neutral-buffered formalin, focal immunoperoxidase staining was detected in the marginal space of almost all eosinophils (D). Eosinophils were damaged by a higher concentration ($1\ \text{mM}$) of H_2O_2 and NO_2^- (E). Under this condition, almost no neutrophils were stained (F). Bars indicate $10\ \mu\text{m}$.




Fig. 2 Histocytochemical detection of superoxide generation by nitroblue tetrazolium (NBT) in activated rat peritoneal eosinophils. Eosinophils were incubated with 1 μM of PMA (A). A phase contrast of the same slide (B).




Fig. 3 Immunoblot analysis of protein nitration by rat peritoneal eosinophils and neutrophils. Samples of eosinophils are shown in lanes 1–8, and those of neutrophils are shown in lanes 9–10. Lane 1, cell (eosinophils); lane 2, cell + 1 μM PMA; lane 3, cell + 1 μM PMA + 50 μM NO_2^- ; lane 4, cell + 50 μM NO_2^- ; lane 5, cell + 1 μM PMA + 50 μM NO_2^- + 50 $\mu\text{g}/\text{ml}$ SOD; lane 6, cell + 1 μM PMA + 50 μM NO_2^- + 30 $\mu\text{g}/\text{ml}$ catalase; lane 7, cell + 50 μM H_2O_2 + 50 μM NO_2^- ; lane 8, supernatant of cell + 1 μM PMA + 50 μM NO_2^- + 1 mg/ml bovine serum albumin; lane 9, cell (neutrophils) + 1 μM PMA + 50 μM NO_2^- ; lane 10, cell + 50 μM H_2O_2 + 50 μM NO_2^- . Tyrosine-nitrated proteins immunostained with anti-nitrotyrosine antibodies (A) and immunostained with anti-nitrotyrosine antibodies preincubated with 10 mM 3-nitrotyrosine for 2 h (B).

respectively (Table 1). Nitration of free tyrosine by eosinophils after incubation with PMA and 50 μM NO_2^- was significantly augmented by SOD, methionine, and mannitol but inhibited by catalase, DMSO, uric acid, cysteine, α -tocopherol, and γ -tocopherol. Tyrosine nitration by ONOO^- was augmented by SOD but inhibited by methionine, mannitol, DMSO, uric acid, cysteine, and ethanol (Fig. 4).

Nitrotyrosine formation in eosinophils in gastric mucosa. Immunostaining for nitrotyro-

sine was seen on infiltrating multinucleate cells (Fig. 5A), while the control study using non-immune rabbit IgG showed negative staining for nitrotyrosine (Fig. 5B). Counterstaining with LFB revealed concordance between the immunoreactive cells and LFB-positive cells (Fig. 5C). Pre-treatment of antibodies with 3-nitrotyrosine abolished the immunoreactivity (Fig. 5D).

Serial sections were recruited to investigate the status and localization of eosinophils further.

Table 1 Free-tyrosine nitration by rat eosinophils and neutrophils

Chemicals	Nitrotyrosine (μM)	
	Eosinophils	Neutrophils
Control	ND	ND
PMA	ND	ND
PMA + 1 μM NO_2^-	ND	ND
PMA + 5 μM NO_2^-	0.25 \pm 0.08 **	0.06 \pm 0.02
PMA + 50 μM NO_2^-	1.15 \pm 0.10 **	0.24 \pm 0.01
1 μM H_2O_2 + 1 μM NO_2^-	0.03 \pm 0.03	ND
5 μM H_2O_2 + 5 μM NO_2^-	0.10 \pm 0.03	ND
50 μM H_2O_2 + 50 μM NO_2^-	2.18 \pm 0.12 **	0.46 \pm 0.02
50 μM H_2O_2	ND	ND
50 μM NO_2^-	ND	ND

Eosinophils or neutrophils (2.5×10^5 cells/250 μl) were incubated in HBSS at 37 $^\circ\text{C}$ with 1 mM DL-tyrosine, with the addition of 1 μM PMA, 1–50 μM NO_2^- , 1–50 μM H_2O_2 or a combination of these reagents. After incubation for 1 h, the reactions were stopped by pelleting the cells. Concentrations of nitrated tyrosine were determined by HPLC as described in Materials and Methods. Values are expressed as the mean \pm SD of 5 experiments. ND = not detected. ** $p < 0.01$ versus neutrophils (unpaired t-test).




Fig. 4 Effect of various scavengers on tyrosine nitration by rat peritoneal eosinophils and peroxynitrite. SOD (50 $\mu\text{g}/\text{ml}$), 30 $\mu\text{g}/\text{ml}$ catalase, 1 mM methionine, 50 mM mannitol, 50 mM DMSO, 1 mM urate, 1 mM cysteine, 69 mM ethanol, 0.1 mM α -tocopherol (in 69 mM ethanol), and 0.1 mM γ -tocopherol (in 69 mM ethanol) were added to the incubation mixtures containing eosinophils (2.5×10^5), 1 mM DL-tyrosine, 1 μM PMA, and 50 μM NO_2^- in 250 μl of HBSS or 0.1 mM ONOO^- and 1 mM DL-tyrosine in 250 μl of HBSS. Each value represents the mean \pm SD of 5 determinations. ** $p < 0.01$, * $p < 0.05$ versus eosinophil control; ## $p < 0.01$ versus ONOO^- control; @@ $p < 0.01$ versus ethanol control.




Fig. 5 The presence of eosinophils and immunohistochemistry for nitrotyrosine in rat gastric wall. Gastric wall was incubated with $1\ \mu\text{M}$ of H_2O_2 and $1\ \mu\text{M}$ of PMA at $37\ ^\circ\text{C}$ for 1 h. Paraffin-embedded sections were immunostained for nitrotyrosine. We used $10\ \text{mM}$ of nitrotyrosine to block the reaction of anti-nitrotyrosine antibody (**D**). Counterstaining was held with hematoxylin (**A**, **B**, **D**) or LFB (**C**). Bars indicate $50\ \mu\text{m}$.




Fig. 6 Co-localization between nitrotyrosine and eosinophil granules in rat gastric wall. Gastric wall slices were incubated with $1\ \mu\text{M}$ of H_2O_2 and $1\ \mu\text{M}$ of PMA at $37\ ^\circ\text{C}$ for 1 h before being embedded in paraffin. Serial sections were stained with LFB (**A**), or immunostained for nitrotyrosine with hematoxylin counterstaining (**B**). Bars indicate $50\ \mu\text{m}$.

Staining with LFB revealed that eosinophils were localized in lamina muscularis mucosa in the gastric mucosa (Fig. 6A). Almost all of the eosinophils were 3-nitrotyrosine-positive (Fig. 6B).

Nitric oxide production from eosinophils and neutrophils. The release of NO from neutrophils was elevated by the addition of L-arginine and suppressed by an inhibitor of iNOS, S-methylisothiourea. The basal level of NO produced from eosinophils was not changed by the addition of L-arginine and S-methylisothiourea (Fig. 7).

Immunocytochemical detection of iNOS.

Eosinophils did not show immunofluorescence for iNOS (Fig. 8A), but neutrophils and contaminating mononuclear cells did show such immunofluorescence (Fig. 8B). FITC-conjugated non-immune mouse IgG2a did not show the specific fluorescence for nitrotyrosine.

Expression of iNOS mRNA in eosinophils and neutrophils. We evaluated the iNOS mRNA expression in rat eosinophils. iNOS mRNA expression in eosinophils was not detected by 30 PCR cycles, but it was faintly detectable by 40 PCR cycles. iNOS mRNA in neutrophils was detected by 30 PCR cycles (Fig. 9).

Discussion

Eosinophils are considered to be an important source of protein nitration [11], but details of eosinophils involving nitration have not been investigated. In the present study we demonstrated the nitration of protein-bound or free tyrosine by the peroxidase system of eosinophils. We found a marked difference in the immunocytochemical appearance of nitrotyrosine between eosinophils incubated with PMA in the presence of NO_2^- and eosinophils incubated with NO_2^- and H_2O_2 . The NBT test for activated eosinophils showed that almost all eosinophils generated $\text{O}_2^{\cdot-}$ by PMA. Therefore, theoretically, $\text{O}_2^{\cdot-}$ generated in the outside of eosinophils should be converted to H_2O_2 . Hydrogen peroxide easily penetrates the cell wall and acts on eosinophil peroxidase to produce immunostainable nitrotyrosine in the presence of $50 \mu\text{M NO}_2^-$. Incubation with H_2O_2 and NO_2^- indicated that all eosinophils possess the capacity for nitration (Fig. 1C, D). However, it seems only eosinophils with a small round nucleus (Fig. 1B) con-

tributed to tyrosine nitration when stimulated with PMA and $50 \mu\text{M NO}_2^-$.

Generally, eosinophils collected from blood easily fall into apoptosis [36]. We observed that almost all eosinophils had small nuclei within 5 h of being isolated (data not shown). Thus, small eosinophils, probably apoptotic-like cells, and eosinophilic granules released from these cells are concerned in local nitration of eosinophils. Since most intact eosinophils were not concerned in nitration, we can say that tyrosine nitration couldn't be raised without apoptotic-like cells in peritoneal fluid. Further investigation is needed to clarify whether small nuclei cells are apoptotic.

Why were the intact cells prevented from nitration? The higher extracellular excretion potential of NO_2^- and enzymatic activity of catalase in intact cells might be a cause. Further investigation to elucidate the nitration mechanism by peripheral eosinophils will be important in clarifying the cause.

Cyto-centrifuged eosinophils showed immunostaining for nitrotyrosine when they were incubated with $1 \mu\text{M H}_2\text{O}_2$ and $1 \mu\text{M NO}_2^-$ (Fig. 1D), implying that fixation before the stimulation may cause the disruption of the cytoplasmic membrane and lead to remarkable nitration. Moreover, freezing for storage and




Fig. 7 Nitric oxide (NO) production from rat peritoneal eosinophils and neutrophils. NO production from eosinophils and neutrophils was measured after incubation with 0.5 mM L-arginine or 0.5 mM L-arginine and 12 μM S-methylisothiourea (S-Methyl-ITU). Each value represents the mean \pm SD of 5 determinations. @@ $p < 0.01$ versus eosinophil control, ** $p < 0.01$ versus neutrophil control, and ## $p < 0.01$ versus arginine.




Fig. 8 Immunofluorescence for iNOS in rat peritoneal eosinophils and neutrophils. Immunofluorescence for iNOS was not detected in eosinophils (A). A phase contrast of the slide shown in (A) (C). Immunofluorescence was detected in neutrophils and contaminating mononuclear cells (B). A phase contrast of the same slide shown in (B) (D).




Fig. 9 Expression of iNOS mRNA in rat peritoneal eosinophils (E) and neutrophils (N). "30" or "40" indicate the PCR cycles run.

drying after cytocentrifugation may inactivate catalase and other enzymes that suppress nitration.

We also investigated the status of constituent eosinophils in the stomach and found many eosinophils existed in lamina muscularis mucosa, which was consistent with a previous report [37]. More importantly, very high potency of nitration was observed in almost all cells. This suggests that the characteristics of eosinophils in gastric mucosa are different

from those eluted in the peritoneal cavity. Eosinophils in the tissue are resistant to apoptosis, and interleukin-5 may be associated with this phenomenon [38]. However, the mechanisms of expression of nitration capacity without apoptosis are not yet known.

It is unclear what reactive nitrogen species contribute to the nitration of tyrosine by EPO-catalyzed oxidation of NO_2^- . There is no direct evidence that a potential nitrogen species contributes to tyrosine nitration by peroxidase-catalyzed oxidation of NO_2^- . However, 2 reactive nitrogen species from the results of MPO are postulated. One is nitrogen dioxide (NO_2) formed by one-electron oxidation [20], and the other is peroxyntrous acid (ONOOH) formed by two-electron oxidation [39]. The difference in the inhibition of free-tyrosine nitration by both EPO-catalyzed oxidation of NO_2^- and ONOO^- with catalase, methionine, mannitol, ethanol, α -tocopherol,

and γ -tocopherol suggests that reactive nitrogen species contributing to the nitration of tyrosine in EPO-catalyzed oxidation of NO_2^- may be different from ONOO^- (Fig. 4). The inhibitory effect of catalase and the augmented effect of SOD (Fig. 3) on the generation of nitrotyrosine support this idea. α -Tocopherol and γ -tocopherol reacted with ONOO^- to become α -tocopheryl quinone and 5-nitro- γ -tocopherol, respectively [40]. SOD1/ H_2O_2 -dependent oxidation of NO_2^- depleted α - and γ -tocopherol [41]. NO inhibition of α - and γ -tocopherol in ONOO^- -dependent tyrosine nitration may have contributed to the high reaction rate constants of ethanol with hydroxyl radical-like oxidants from ONOO^- [42] because ethanol was used as a solvent of α - and γ -tocopherol. However, although oxidation and nitration products from the reaction of α - and γ -tocopherol with reactive nitrogen species from EPO-catalyzed oxidation of NO_2^- were not presented, tyrosine-nitrating species from EPO-catalyzed oxidation of NO_2^- may be similar to that from SOD1/ H_2O_2 -dependent oxidation of NO_2^- [41].

A previous report indicated that the tyrosine-nitrating capacity of porcine EPO exceeded that of MPO by at least 4-fold at physiological concentrations of halides, and every concentration of NO_2^- examined [23]. In this study, similar results were obtained in the difference of the tyrosine-nitrating capacity between rat eosinophils and neutrophils activated by PMA and NO_2^- . The enzymological difference in the capacity of tyrosine nitration between EPO and MPO and the difference in the total contents of EPO in eosinophils and MPO in neutrophils may have contributed to the difference in tyrosine-nitrating activity between eosinophils and neutrophils [23, 43, 44].

Previous reports have described another nitrogen species generated from the reaction of O_2^- with NO_2^- , namely peroxyxynitrate (O_2NOO^-) (Fig. 10) [45, 46]. O_2NOO^- is reported to be unreactive to tyrosine and to act as an inactivator of NO_2 [47], and it theoretically forms nitrite and singlet oxygen ($^1\text{O}_2$) [48]. A recent report suggests that singlet oxygen is capable of forming nitrotyrosine under photosensitization [49]. In this study, we examined free tyrosine nitration by exposing eosinophils to PMA and H_2O_2 . By our calculation, 10^6 eosinophils can produce 30.2 nmol/ml/hr of H_2O_2 when stimulated with 1 μM PMA. The

capacity for tyrosine nitration of eosinophils stimulated with 1 μM PMA (= 30.2 μM H_2O_2) and 50 μM NO_2^- was proportional to that of 50 μM H_2O_2 and 50 μM NO_2^- , as indicated in Table 1. Therefore, the inhibitory effect of O_2NOO^- on tyrosine nitration may be very limited, but such an effect cannot be ruled out because we did not investigate the formation of O_2NOO^- and singlet oxygen in this study. Further investigation is needed to reveal the contribution of O_2NOO^- toward nitration.

In conclusion, the present study demonstrated that smaller, possibly apoptotic eosinophils have a potent capacity to generate reactive nitrogen species




Fig. 10 Mechanisms of different levels of nitration efficiency. When exposed to PMA and NO_2^- , NADPH oxidase (closed circle) is activated by PMA stimulation to form O_2^- . Then, formation of NO_2 was catalyzed by secreted EPO (gray circle) with external NO_2^- and H_2O_2 generated from O_2^- . Tyrosine nitration is facilitated by NO_2 . However, PMA stimulates NADPH oxidase continuously; O_2NOO^- can be formed from existing NO_2 and O_2^- , and it rapidly degrades to singlet oxygen molecule ($^1\text{O}_2$). This molecule is not reactive with tyrosine residue, and the formation of nitrotyrosine was suppressed as a result (A). However, when exposed to nitrite and H_2O_2 , the amount of NO_2 production catalyzed by EPO would increase, because H_2O_2 is not supplied via O_2^- and then the formation of O_2NOO^- is lacking (B).

responsible for free or protein-bound tyrosine nitration via a peroxidase-catalyzed system. Moreover, we found a difference in the nitration capacity of eosinophils between eosinophils in the gastric mucosa and those in the peritoneal cavity. However, the mechanisms of heterogeneity in nitration are unknown and should be clarified in future studies.

Acknowledgements. This study was supported, in part, by a Grant-in-Aid from the Ministry of Education, Science and Culture and, in part, by Special Coordination Funds from the Ministry of Education, Culture, Sports, Science and Technology, of the Japanese Government.

References

- Kazura JW, Fanning MM, Blumer JL and Mahmoud AA: Role of cell-generated hydrogen peroxide in granulocyte-mediated killing of *Schistosoma mansoni* in vitro. *J Clin Invest* (1981) 67: 93–102.
- Behm CA and Ovington KS: The role of eosinophils in parasitic helminth infections: insights from genetically modified mice. *Parasitol Today* (2000) 16: 202–209.
- Gleich GJ, Ottesen EA, Leiferman KM and Ackerman SJ: Eosinophils and human disease. *Int Arch Allergy Appl Immunol* (1989) 88: 59–62.
- Horwitz RJ and Busse WW: Inflammation and asthma. *Clin Chest Med* (1995) 16: 583–602.
- Holgate ST: The epidemic of allergy and asthma. *Nature* (1999) 402: B2–B4.
- Allen JN, Davis WB and Pacht ER: Diagnostic significance of increased bronchoalveolar lavage fluid eosinophils. *Am Rev Respir Dis* (1990) 142: 642–647.
- Sanz ML, Parra A, Prieto I, Dieguez I and Oehling AK: Serum eosinophil peroxidase (EPO) levels in asthmatic patients. *Allergy* (1997) 52: 417–422.
- Gleich GJ: Mechanisms of eosinophil-associated inflammation. *J Allergy Clin Immunol* (2000) 105: 651–663.
- Babior BM: Oxygen-dependent microbial killing by phagocytosis. *N Engl J Med* (1978) 298: 659–668.
- Saleh D, Ernst P, Lim S, Barnes PJ and Giaid A: Increased formation of the potent oxidant peroxynitrite in the airways of asthmatic patients is associated with induction of nitric oxide synthase: effect of inhaled glucocorticoid. *FASEB J* (1998) 12: 929–937.
- MacPherson JC, Comhair SA, Erzurum SC, Klein DF, Lipscomb MF, Kavuru MS, Samoszuk MK and Hazen SL: Eosinophils are a major source of nitric oxide-derived oxidants in severe asthma: characterization of pathways available to eosinophils for generating reactive nitrogen species. *J Immunol* (2001) 166: 5763–5772.
- Beckman JS, Beckman TW, Chen J, Marshall PA and Freeman BA: Apparent hydroxyl radical production from peroxynitrite: implications for endothelial injury by nitric oxide and superoxide. *Proc Natl Acad Sci U S A* (1990) 87: 1620–1624.
- del Pozo V, de Arruda-Chaves E, de Andres B, Cardaba B, Lopez-Farre A, Gallardo S, Cortegano I, Vidarte L, Jurado A, Sastre J, Palomino P and Lahoz C: Eosinophils transcribe and translate messenger RNA for inducible nitric oxide synthase. *J Immunol* (1997) 158: 859–864.
- Miles AM, Owens MW, Milligan S, Johnson GG, Fields JZ, Ing TS, Kottapalli V, Keshavarzian A and Grisham MB: Nitric oxide synthase in circulating vs. extravasated polymorphonuclear leukocytes. *J Leukoc Biol* (1995) 58: 616–622.
- Ischiropoulos H, Zhu L, Chen J, Tsai M, Martin JC, Smith CD, Beckman JS: Peroxynitrite-mediated tyrosine nitration catalyzed by superoxide dismutase. *Arch Biochem Biophys* (1992) 298: 431–437.
- Beckman JS, Ischiropoulos H, Zhu L, van der Woerd M, Smith C, Chen J, Harrison J, Martin JC and Tsai M: Kinetics of superoxide dismutase- and iron-catalyzed nitration of phenolics by peroxynitrite. *Arch Biochem Biophys* (1992) 298: 438–445.
- Abe K, Pan LH, Watanabe M, Konno H, Kato T and Itoyama Y: Upregulation of protein-tyrosine nitration in the anterior horn cells of amyotrophic lateral sclerosis. *Neuro Res* (1997) 19: 124–128.
- Beckmann JS, Ye YZ, Anderson PG, Chen J, Accavitti MA, Tarpey MM and White CR: Extensive nitration of protein tyrosines in human atherosclerosis detected by immunohistochemistry. *Biol Chem Hoppe Seyler* (1994) 375: 81–88.
- Sakaguchi AA, Miura S, Takeuchi T, Hokari R, Mizumori M, Yoshida H, Higuchi H, Mori M, Kimura H, Suzuki H, Ishii H: Increased expression of inducible nitric oxide synthase and peroxynitrite in *Helicobacter pylori* gastric ulcer. *Free Radic Biol Med* (1999) 27: 781–789.
- Mannick EE, Bravo LE, Zarama G, Realpe JL, Zhang XJ, Ruiz B, Fontham ET, Mera R, Miller MJ and Correa P: Inducible nitric oxide synthase, nitrotyrosine, and apoptosis in *Helicobacter pylori* gastritis: effect of antibiotics and antioxidants. *Cancer Res* (1996) 56: 3238–3243.
- van der Vliet A, Eiserich JP, Halliwell B and Cross CE: Formation of reactive nitrogen species during peroxidase-catalyzed oxidation of nitrite. A potential additional mechanism of nitric oxide-dependent toxicity. *J Biol Chem* (1997) 272: 7617–7625.
- van Dalen CJ, Winterbourn CC, Senthilmohan R and Kettle AJ: Nitrite as a substrate and inhibitor of myeloperoxidase. *J Biol Chem* (2000) 275: 11638–11644.
- Wu W, Chen Y and Hazen SL: Eosinophil peroxidase nitrates protein tyrosyl residues. Implications for oxidative damage by nitrating intermediates in eosinophilic inflammatory disorders. *J Biol Chem* (1999) 274: 25933–25944.
- Ueda T, Maekawa T, Sadamitsu D, Oshita S, Ogino K and Nakamura K: The determination of nitrite and nitrate in human plasma by capillary zone electrophoresis. *Electrophoresis* (1995) 16: 1002–1004.
- Torre D, Ferrario G, Speranza F, Orani A, Fiori GP and Zeroli C: Serum concentrations of nitrite in patients with HIV-1 infection. *J Clin Pathol* (1996) 49: 574–576.
- Hunt T, Byrns RE, Ignarro LJ and Gaston B: Condensed expirate nitrite as a home maker for acute asthma. *Lancet* (1995) 346: 1235–1236.
- Johnson GR and Metcalf D: Detection of new type of mouse eosinophil colony by Luxol-fast-blue staining. *Exp Haematol* (1980) 8: 549–561.
- Ogino K, Izumi Y, Segawa H, Takeyama Y, Ishiyama H, Houbara T, Uda T and Yamashita S: Zinc hydroxide induced respiratory burst in rat neutrophils. *Eur J Pharmacol* (1994) 270: 73–78.
- Babior BM, Kipnes RS and Curnutte JT: Biological defense mechanisms. The production by leukocytes of superoxide, a potential bactericidal agent. *J Clin Invest* (1973) 52: 741–744.
- Krueger GG, Ogden BE and Weston WL: In vitro quantitation of

- cell-mediated immunity in guinea-pigs by macrophage reduction of nitro-blue tetrazolium. *Clin Exp Immunol.* (1976) 23: 517-524.
31. Beckman JS, Chen J, Ischiropoulos H and Crow JP: Oxidative chemistry of peroxynitrite. *Methods Enzymol* (1994) 233: 229-240.
 32. Ogino K, Kodama N, Nakajima M, Yamada A, Nakamura H, Nagase H, Sadamitsu D and Maekawa T: Catalase catalyzes nitrotyrosine formation from sodium azide and hydrogen peroxide. *Free Radic Res* (2001) 35: 735-747.
 33. Sokolovsky M, Riordan JF and Vallee BL: Conversion of 3-nitrotyrosine to 3-aminotyrosine in peptides and proteins. *Biochem Biophys Res Commun* (1967) 27: 20-25.
 34. Smith PF, Krack G, McKee RL, Johnson DG, Gandolfi AJ, Hruby VJ, Krumdieck CL and Brendel K: Maintenance of adult rat liver slices in dynamic organ culture. *In Vitro Cell Dev Biol* (1986) 22: 706-712.
 35. Misko TP, Schilling RJ, Salvemini D, Moore WM and Currie MG: A fluorometric assay for the measurement of nitrite in biological samples. *Anal Biochem* (1993) 214: 11-16.
 36. Yamaguchi Y, Suda T, Ohta S, Tominaga K, Miura Y and Kasahara T: Analysis of the survival of mature human eosinophils: interleukin-5 prevents apoptosis in mature human eosinophils. *Blood* (1991) 78: 2542-2547.
 37. Mishra A, Hogan SP, Lee JJ, Foster PS and Rothenberg ME: Fundamental signals that regulate eosinophil homing to the gastrointestinal tract. *J Clin Invest* (1999) 103: 1719-1727.
 38. Dewson G, Cohen GM and Wardlaw AJ: Interleukin-5 inhibits translocation of Bax to the mitochondria, cytochrome c release, and activation of caspases in human eosinophils. *Blood* (2001) 98: 2239-2247.
 39. Sampson JB, Ye Y, Rosen H and Beckman JS: Myeloperoxidase and horseradish peroxidase catalyze tyrosine nitration in proteins from nitrite and hydrogen peroxide. *Arch Biochem Biophys* (1998) 356: 207-213.
 40. Goss SP, Hogg N and Kalyanaraman B: The effect of alpha-tocopherol on the nitration of gamma-tocopherol by peroxynitrite. *Arch Biochem Biophys.* (1999) 363: 333-340.
 41. Goss SP, Singh RJ and Kalyanaraman B: Bicarbonate enhances the peroxidase activity of Cu, Zn-superoxide dismutase. Role of carbonate anion radical. *J Biol Chem* (1999) 274: 28233-28239.
 42. Crow JP, Spruell C, Chen J, Gunn C, Ischiropoulos H, Tsai M, Smith CD, Radi R, Koppenol WH and Beckman JS: On the pH-dependent yield of hydroxyl radical products from peroxynitrite. *Free Radic Biol Med* (1994) 16: 331-338.
 43. Bos AJ, Wever R, Hamers MN and Roos D: Some enzymatic characteristics of eosinophil peroxidase from patients with eosinophilia and from healthy donors. *Infect Immun* (1981) 32: 427-431.
 44. Wever R, Hamers MN, de Graaf CJ, Weening RS and Roos D: Characterization of the peroxidase in human eosinophils. *Adv Exp Med Biol* (1982) 141: 501-509.
 45. Pryor WA, Houk KN, Foote CS, Fukuto JM, Ignarro LJ, Squadrito GL and Davies KJ: Free radical biology and medicine: it's a gas, man! *Am J Physiol Regul Integr Comp Physiol* (2006) 291: R491-R511.
 46. Logager T and Sehested K: Formation and decay of peroxynitric acid: a pulse radiolysis study. *J Phys Chem* (1993) 97: 10047-10052.
 47. Kirsch M, Lehnig M, Korth HG, Sustmann R and de Groot H: Inhibition of peroxynitrite-induced nitration of tyrosine by glutathione in the presence of carbon dioxide through both radical repair and peroxynitrate formation. *Chemistry* (2001) 7: 3313-3320.
 48. Goldstein S, Czapski G, Lind J and Merenyi G: Mechanism of Decomposition of Peroxynitric Ion (O_2NOO^-): Evidence for the Formation of $O_2^{\cdot -}$ and $^{\cdot}NO_2$ Radicals. *Inorg Chem* (1998) 37: 3943-3947.
 49. Pecci L, Montefoschi G, Antonucci A, Costa M and Cavallini D: Methylene blue photosensitized oxidation of hypotaurine in the presence of azide generates reactive nitrogen species: formation of nitrotyrosine. *Biochem Biophys Res Commun* (2003) 301: 411-416.