

Mathematical Journal of Okayama University

Volume 25, Issue 1

1983

Article 14

JUNE 1983

On the two decompositions of a measure space by an operator semigroup

Hisakichi Suzuki*

*Chiba Institute Of Technology

Copyright ©1983 by the authors. *Mathematical Journal of Okayama University* is produced by
The Berkeley Electronic Press (bepress). <http://escholarship.lib.okayama-u.ac.jp/mjou>

ON THE TWO DECOMPOSITIONS OF A MEASURE SPACE BY AN OPERATOR SEMIGROUP

HISAKICHI SUZUKI

Let $T = \{T_t; t > 0\}$ be a strongly continuous semigroup of positive linear operators on $L_1(X) = L_1(X, \mathfrak{F}, \mu)$, where $(X, \mathfrak{F}, \mu)$ is a σ -finite measure space. The semigroup T decomposes X into two parts C and D , called the initially conservative and initially dissipative parts, respectively. It holds that

$$(1) \quad T_t f = 0 \text{ on } D \text{ for any } f \in L_1(X) \text{ and } t > 0,$$

and that if $E \in \mathfrak{F}$, instead of D , satisfies the property (1), then $E \subset D$, where the inclusion holds up to a set of measure zero. (Throughout this paper, inclusions and equalities of sets or functions are considered in this sense.)

The space X has another decomposition into C^* and D^* by the same semigroup T , such that

$$(2) \quad \|T_t(f 1_{D^*})\|_1 = 0 \text{ for any } f \in L_1(X) \text{ and } t > 0,$$

and that if $E \in \mathfrak{F}$, instead of D^* , satisfies the property (2), then $E \subset D^*$, where 1_{D^*} denotes the characteristic function of D^* .

For the definitions and other characterizations of those decompositions we refer the reader to [1] in the case of contraction semigroups, and in the case of more general ones, to [2] and [3], in the latter of which C^* and D^* are denoted by P and N respectively.

In this paper we establish some theorems on the inclusion relations between D and D^* , motivated by the statement in [2] that $D \subset D^*$. If T_t are contractions, that is,

$$\|T_t\|_1 \leq 1 \text{ for any } t > 0,$$

then an inclusion relation between D and D^* holds (Theorem 1). But, in general, there is no relation between them as Theorem 3 shows.

Theorem 1. *If $T = \{T_t; t > 0\}$ is a strongly continuous semigroup of positive contraction operators, then $D^* \subset D$.*

Proof. If $T_t(t > 0)$ are contractions, then it holds that

$$(3) \quad T_t(L_1(C^*)) \subset L_1(C^*)$$

by [3, Remark 1], where $L_1(C^*)$ denotes the subspace consisting of functions vanishing outside C^* . Now, (3) implies together with (2) that

$$T_t f = 0 \text{ on } D^* \text{ for any } f \in L_1(X) \text{ and } t > 0,$$

and hence $D^* \subset D$.

Even in the case of contractions, it does not hold that $D = D^*$ in general.

Theorem 2. *There exists a strongly continuous semigroup $T = \{T_t; t > 0\}$ of positive contractions such that $D^* \subsetneq D$.*

Proof. Let $T = \{T_t\}$ be a strongly continuous semigroup of Markovian operators, that is,

$$(4) \quad \int_X T_t f d\mu = \int_X f d\mu \quad \text{for any } f \in L_1(X) \text{ and } t > 0.$$

By (2) and (4) we can easily conclude that $\mu(D^*) = 0$ for Markovian operators. If in addition $\mu(D) > 0$ holds for this semigroup T , then the theorem will be proved.

Now we assume that $X = C$ for the present Markovian semigroup T , and construct, making use of T , another Markovian semigroup $\tilde{T} = \{\tilde{T}_t\}$ such that the initially dissipative part $\tilde{D}$ determined by $\tilde{T}$ has positive measure.

Step 1. Definition of the underlying measure space $(\tilde{X}, \tilde{\mathfrak{F}}, \tilde{\mu})$. Let $A \in \mathfrak{F}$ have positive measure, and $(A, \mathfrak{F}_A, \mu_A)$ be the measure space defined as $\mathfrak{F}_A = \{B \cap A; B \in \mathfrak{F}\}$ and $\mu_A(B) = \mu(B \cap A)$. Now let $(X', \mathfrak{F}', \mu')$ be a measure space isomorphic to $(A, \mathfrak{F}_A, \mu_A)$ with $X \cap X' = \emptyset$. By an isomorphism of two measure spaces A and X' , we mean a one-to-one mapping τ from A onto X' such that

$$\tau^{-1}\mathfrak{F}' \subset \mathfrak{F}_A, \text{ and } \mu_A(\tau^{-1}B) = \mu'(B) \text{ for any } B \in \mathfrak{F}'.$$

We define a σ -finite measure space $(\tilde{X}, \tilde{\mathfrak{F}}, \tilde{\mu})$ as

$$\begin{aligned} \tilde{X} &= X \cup X', \\ \tilde{\mathfrak{F}} &= \{B \cup B'; B \in \mathfrak{F} \text{ and } B' \in \mathfrak{F}'\} \end{aligned}$$

and for $\tilde{B} = B \cup B' \in \tilde{\mathfrak{F}}$

$$\tilde{\mu}(\tilde{B}) = \mu(B) + \mu'(B').$$

Step 2. Construction of Markovian operators $\tilde{T}_t$ on $L_1(\tilde{X})$. Let $\tilde{f}$ be in $L_1(\tilde{X})$, and define a corresponding f in $L_1(X)$ as

$$(5) \quad f(x) = \begin{cases} \tilde{f}(x) + \tilde{f}(\tau x) & \text{if } x \in A \\ \tilde{f}(x) & \text{if } x \in X \setminus A, \end{cases}$$

where $\tilde{f}$ may be any one of versions of the equivalence class.

Now we define $\tilde{T} = \{\tilde{T}_t; t > 0\}$ as

$$(\tilde{T}_t \tilde{f})(x) = \begin{cases} 0 & \text{if } x \in X' \\ (T_t f)(x) & \text{if } x \in X = \tilde{X} \setminus X' \end{cases}$$

It is shown as follows that $\tilde{T}$ forms a semigroup. Let $s, t > 0$ and $\tilde{f} \in L_1(\tilde{X})$. Now let f and h correspond respectively to $\tilde{f}$ and $\tilde{h} = \tilde{T}_t \tilde{f}$ as in (5). Then since $h = \tilde{h} = T_t f$ on X , we have

$$\tilde{T}_s(\tilde{T}_t \tilde{f}) = T_s h = T_s(T_t f) = T_{s+t} f \text{ on } X.$$

Hence $\tilde{T}_s(\tilde{T}_t \tilde{f}) = \tilde{T}_{s+t} \tilde{f}$ holds.

It is easily shown that $\tilde{T}_t$ are Markovians, and that the initially dissipative part $\tilde{D}$ contains X' , which has positive measure.

Without the assumption that T is a semigroup of contractions, neither inclusion $D \subset D^*$ nor $D^* \subset D$ holds in general.

Theorem 3. *There exists a strongly continuous semigroup of positive operators which satisfies*

$$(a) \quad \mu(D) = 0 \text{ and } \mu(D^*) > 0;$$

and also there exists a strongly continuous semigroup of positive operators which satisfies

$$(b) \quad \mu(D) > 0, \mu(D^*) > 0 \text{ and } \mu(D \cap D^*) = 0.$$

Proof. Let X be the closed unit interval $[0, 1]$, and μ the Lebesgue measure on X . We construct two semigroups $T = \{T_t; t \geq 0\}$ on $L_1(X)$.

1) For $f \in L_1(X)$ and $t \geq 0$ we define T_t as

$$(T_t f)(x) = \begin{cases} e^{\alpha t} f(x) & \text{if } x \in [0, 1/2) \\ e^{\alpha t} f(x-1/2) & \text{if } x \in [1/2, 1], \end{cases}$$

where α is a nonzero constant. Then $T = \{T_t; t \geq 0\}$ is a semigroup of positive operators with the norms

$$\|T_t\|_1 \leq 2e^{\alpha t} \quad (t \geq 0).$$

We have $\mu(D) = 0$, while $\mu(D^*) \geq 1/2$, since it holds that for any $E \subset [1/2, 1]$ and $t \geq 0$

$$0 \leq \|T_t(1_E)\|_1 \leq \|T_t(1_{[1/2, 1]})\|_1 = 0.$$

This semigroup T satisfies the property (a).

2) Next we define T_t as

$$(T_t f)(x) = \begin{cases} 0 & \text{if } x \in [0, 1/3) \\ e^{at} f\{(x-1/3)+f(x+1/3)\} & \text{if } x \in [1/3, 2/3) \\ e^{at} f\{(x)+f(x-2/3)\} & \text{if } x \in [2/3, 1] \end{cases}$$

Clearly it holds that $\|T_t\|_1 \leq 2e^{at}$ ($t \geq 0$), $D = [0, 1/3)$ and $D^* = [1/3, 2/3)$. Hence a semigroup with the property (b) is constructed.

REFERENCES

- [1] M.A. AKCOGLU and R.V. CHACON: A local ratio theorem, *Canad. J. Math.* **22** (1970), 545—552.
- [2] M.A. AKCOGLU and U. KRENGEL: A differentiation theorem in L_p , *Math. Z.* **169** (1979), 31—40.
- [3] R. SATO: On local ergodic theorems for positive semigroups, *Studia Math.* **63** (1978), 45—55.

CHIBA INSTITUTE OF TECHNOLOGY

(Received June 7, 1982)