

Mathematical Journal of Okayama University

Volume 30, Issue 1

1988

Article 4

JANUARY 1988

Fixed rings of simple rings

Yoshimi Kitamura*

*Tokyo Gakugei University

Copyright ©1988 by the authors. *Mathematical Journal of Okayama University* is produced by
The Berkeley Electronic Press (bepress). <http://escholarship.lib.okayama-u.ac.jp/mjou>

FIXED RINGS OF SIMPLE RINGS

Dedicated to Professor Hisao Tominaga on his 60th birthday

YOSHIMI KITAMURA

Let A be a ring with identity and G a finite group of ring automorphisms of A . We denote by A^G the subring of A consisting of elements $a \in A$ such that $\sigma(a) = a$ for all $\sigma \in G$.

A need not be finitely generated over A^G even if A is a simple artinian ring as shown by Björk [1]. However, if the order of G is invertible in A , we can obtain the following result.

Theorem. *If A is a finite direct sum of simple rings and if the order of G is invertible in A , then A is a Frobenius extension of A^G .*

The purpose of this paper is to show the above result.

Throughout this paper, all rings, subrings, ring homomorphisms and modules are assumed to be unital.

According to Kasch [2], a ring extension A/B is called a *Frobenius extension* provided that A is finitely generated projective as a right B -module and $A \cong \text{Hom}(A_B, B_B)$ as B - A -bimodules. As shown in Onodera [7], A/B is a Frobenius extension if and only if there exist a B - B -homomorphism h of A to B and a finite number of elements r_i 's, l_i 's in A such that $x = \sum_i r_i h(l_i x) = \sum_i h(x r_i) l_i$ for all $x \in A$. When this is the case, we shall call $(h; r_i, l_i)_i$ a *Frobenius system* for A/B .

The following is obvious from the definition of Frobenius extension.

Lemma 1. *Let A_i/B_i ($i = 1, \dots, n$) be ring extensions. Then the finite product $A_1 \times \cdots \times A_n$ of rings is a Frobenius extension of $B_1 \times \cdots \times B_n$ if and only if each A_i/B_i is a Frobenius one.*

The following is well-known (see [7]).

Lemma 2. *Let A/B be a Frobenius extension. Then*

- (1) *For any Frobenius extension A'/A , A'/B is a Frobenius one.*
- (2) *Suppose B is contained in the center of A . For any algebra B' over B , $A \otimes_B B'/B' (\cong B \otimes_B B')$ is a Frobenius extension.*

(3) For any left A -module X , $\text{Hom}({}_A X, {}_A A) \cong \text{Hom}({}_B X, {}_B B)$.

Lemma 3. Let A_0/B_0 be a Frobenius extension. Let $f_j: A_0 \rightarrow A_j$ ($j = 1, \dots, m$) be ring isomorphisms. Then the finite product $A = A_0 \times A_1 \times \dots \times A_m$ of rings is a Frobenius extension of $B = \{(x, f_1(x), \dots, f_m(x)); x \in B_0\}$.

Proof. Let $(h_0; r_i, l_i)_{1 \leq i \leq n}$ be a Frobenius system for A_0/B_0 . Let define a mapping h of A to B by

$h(x_0, f_1(x_1), \dots, f_m(x_m)) = (y, f_1(y), \dots, f_m(y))$ for $x_0, x_1, \dots, x_m \in A_0$, where $y = h_0(x_0 + x_1 + \dots + x_m)$. Let $r_{j,i}, l_{j,i}$ ($0 \leq j \leq m, 1 \leq i \leq n$) be the elements of A defined as follows:

$$\begin{aligned} r_{0,i} &= (r_i, 0, \dots, 0), & l_{0,i} &= (l_i, 0, \dots, 0) \\ &\dots\dots\dots \\ r_{m,i} &= (0, \dots, 0, f_m(r_i)), & l_{m,i} &= (0, \dots, 0, f_m(l_i)) \quad (i = 1, \dots, n). \end{aligned}$$

Then one can see that $(h; r_{j,i}, l_{j,i})_{j,i}$ is a Frobenius system for A/B .

A ring A is simple if it has no proper two-sided ideals. Let G be a finite group of ring automorphisms of A . G is inner if every element σ of G is inner, that is, there exists a unit u in A such that $\sigma(a) = uau^{-1}$ for all $a \in A$, and G is outer if the identity element of G is the only inner automorphism in G .

The following result is due to Miyashita [4].

Lemma 4. If a ring A is simple and if G is outer, then A is a Frobenius extension of A^G .

Lemma 5. If a ring A is simple and if G is inner such that its order is invertible in A , then A is a Frobenius extension of A^G .

Proof. Let S be the algebra of G , that is $S = \sum_{\sigma \in G} J(\sigma)$, where $J(\sigma) = \{x \in A; xa = \sigma(a)x \text{ for all } a \in A\}$. Let C be the center of A . S is then a finite dimensional separable algebra over C (see, for example, [6], page 28). Further, the centralizer of S in A coincides with A^G . Let $S = S_1 \oplus \dots \oplus S_n$ be a decomposition of S into simple rings. Let $T = A \otimes {}_C S^o$, $T_i = A \otimes {}_C S_i^o$ ($i = 1, \dots, n$), where S^o and S_i^o denote the opposite rings of S and S_i respectively. We will show that when we consider A a left T -module by means of $(a \otimes s^o)x = axs$, A is a generator. Since S^o and S_i^o 's are Frobenius extensions of C , T and T_i 's are so over A by Lemma 2(2). Hence, by Lemma 2(3), $\text{Hom}({}_{T_i} A e_i, {}_{T_i} T_i) \cong \text{Hom}({}_A A e_i, {}_A A) \neq 0$, where e_i denotes the identity element of S_i . Since T_i is simple, $A e_i$ is a generator over T_i , and so A is a

generator over T as desired. Thus $T \otimes_A A (\cong T)$ is isomorphic to a direct summand of a finite direct sum of copies of A as a left T -module. Therefore, recalling T/A a Frobenius extension, we have by Theorem 2.10 of [5] that $\text{End}({}_A A)/\text{End}({}_T A)$, or equivalently, A/A^G is a Frobenius extension.

We are now in position to prove the theorem.

Proof of Theorem. We assume first A is simple, and show the theorem by induction on the order $|G|$ of G . Let N be the normal subgroup of G consisting of inner automorphisms in G . By Lemmas 4, 5, we may assume that $1 < |N| < |G|$. Let $T = A^N$, $\bar{G} = G/N$. $\bar{G}$ acts as automorphisms on T . By our induction hypothesis, A is a Frobenius extension of T . We shall show that $T/T^{\bar{G}}$ is a Frobenius extension. By [3], T is a finite direct sum of simple rings. Let $T = T_0 \oplus T_1 \oplus \cdots \oplus T_m$ be a decomposition of T into simple rings. Since every element of $\bar{G}$ induces a permutation of the finite set $\{T_0, T_1, \dots, T_m\}$, we can assume by Lemma 1 that $\bar{G}$ is transitive on the set. Let $\bar{\sigma}_i$ be elements of $\bar{G}$ such that $\bar{\sigma}_i(T_0) = T_i$ ($i = 1, \dots, m$), and let $\bar{G}_0$ be the set of $\bar{\sigma} \in \bar{G}$ such that $\bar{\sigma}(T_0) = T_0$. Then it is easy to see that $T^{\bar{G}} = \{x + \bar{\sigma}_1(x) + \cdots + \bar{\sigma}_m(x); x \in T_0^{\bar{G}_0}\}$. Since T_0 is a Frobenius extension of $T_0^{\bar{G}_0}$ by our induction hypothesis, T is so over $T^{\bar{G}}$ ($= A^G$) by Lemma 3. Hence A is a Frobenius extension of A^G by Lemma 2(1).

We shall next consider a general case. Let $A = A_0 \oplus A_1 \oplus \cdots \oplus A_m$ be a representation of A as a finite direct sum of simple rings. Let G_0 be the set of $\sigma \in G$ with $\sigma(A_0) = A_0$. Then A_0 is a Frobenius extension of $A_0^{G_0}$ from the first case, so that A is a Frobenius extension of A^G by the same argument as in proving T a Frobenius one over $T^{\bar{G}}$ above.

REFERENCES

- [1] J. E. BJÖRK : Conditions which imply that subrings of semiprimary rings are semiprimary, *J. Algebra* **19** (1971), 384–395.
- [2] F. KASCH : Projective Frobenius-Erweiterungen, *Sitzungsber. Heidelberger Akad. Wiss.* 1960/1961, 89–109.
- [3] V. K. KHARCHENKO : Galois subrings of simple rings, *Math. Notes* (transl.) **17** (1975), 533–536.
- [4] Y. MIYASHITA : Finite outer Galois theory of noncommutative rings, *J. Fac. Sci. Hokkaido Univ.* **19** (1966), 114–134.
- [5] Y. MIYASHITA : On Galois extensions and crossed products, *J. Fac. Sci. Hokkaido Univ.* **21**(1970), 97–121.
- [6] S. MONTGOMERY : Fixed rings of finite automorphism groups of associative rings, *Lecture Notes in Math.* **818**, Springer, Berlin-Heidelberg-New York, 1980.
- [7] T. ONODERA : Some studies on projective Frobenius extensions, *J. Fac. Sci. Hokkaido Univ.* **18** (1964), 89–107.

DEPARTMENT OF MATHEMATICS
TOKYO GAKUGEI UNIVERSITY
KOGANEI, TOKYO 184, JAPAN

(Received July 30, 1987)