

Mathematical Journal of Okayama University

Volume 39, Issue 1

1997

Article 10

JANUARY 1997

The Steenrod Algebra and Braid Groups

Mizuho Hikida*

*Hiroshima Prefectural University

Math. J. Okayama Univ. **39** (1997), 89–112

THE STEENROD ALGEBRA AND BRAID GROUPS

MIZUHO HIKIDA

1. Introduction. Let $q \neq 0$ be an element of $\mathbf{Z}/(p)$ for a prime p . In this paper, we define a map $\phi_q: B_t \rightarrow \mathbf{Z}/(p)$ from the braid group B_t to $\mathbf{Z}/(p)$ related to a cohomology of the Steenrod algebra with

$$(1.1) \quad \phi_q(\sigma\tau) = \phi_q(\tau) + (-q)^{|\tau|}\phi_q(\sigma),$$

where $|\tau| = \sum_i \epsilon_i$ for $\tau = \sigma_{i_1}^{\epsilon_1} \sigma_{i_2}^{\epsilon_2} \cdots \sigma_{i_n}^{\epsilon_n}$ (σ_i are generators of B_t). This map emerged from computation of the stable homotopy groups of spheres. In this paper, we argue the relation between this map and a cohomology of the Steenrod algebra. In the forthcoming paper, we shall argue the stable homotopy groups of spheres by using this map.

The braid group B_t of degree t is generated by σ_i ($1 \leq i \leq t-1$) and relations

$$\begin{aligned} (R1) \quad & \sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1} (1 \leq i \leq t-2) \\ (R2) \quad & \sigma_i \sigma_j = \sigma_j \sigma_i \quad (|i-j| \geq 2). \end{aligned}$$

Hence B_t has a presentation $F_t/\langle R1, R2 \rangle$ for a free group $F_t = \langle \sigma_1, \dots, \sigma_{t-1} \rangle$. We notice that the symmetric group S_t has a presentation $F_t/\langle R1, R2, \sigma_i^2 = e \rangle = B_t/\langle \sigma_i^2 = e \rangle$ (e is a unit). We denote $\bar{\sigma} \in S_t$ for $\sigma \in F_t$. For example, $\bar{\sigma}_i$ is a transposition $(i, i+1)$. Then we denote $\sigma N = (n_{\bar{\sigma}^{-1}(1)}, \dots, n_{\bar{\sigma}^{-1}(t)})$.

Definition 1.1. Inductively we define $\bar{\phi}_q: F_t \rightarrow \mathbf{Z}/(p)$ by taking

$$\bar{\phi}_q(e) = 0, \quad \bar{\phi}_q(\sigma_i) = 1, \quad \bar{\phi}_q(\sigma_i^{-1}) = q^{-1} \quad \text{and}$$

$$\bar{\phi}_q(\sigma_i \sigma) = \bar{\phi}_q(\sigma) + (-q)^{|\sigma|}, \quad \bar{\phi}_q(\sigma_i^{-1} \sigma) = \bar{\phi}_q(\sigma) - (-q)^{|\sigma|-1}.$$

Then $\bar{\phi}_q(\gamma\sigma) = \bar{\phi}_q(\sigma) = \bar{\phi}_q(\sigma\gamma)$ for $\gamma \in \langle R1, R2 \rangle$. $\phi_q: B_t \rightarrow \mathbf{Z}/(p)$ is induced from $\bar{\phi}_q$.

At Proposition 5.2, we show (1.1) and that ϕ_q is well defined.

Let A_* be the dual of Steenrod algebra for a prime p . Then

$$(1.2) \quad A_* = \begin{cases} \mathbf{Z}/(2)[\xi_1, \xi_2, \dots] & \text{for } p = 2, \\ E(\tau_0, \tau_1, \dots) \otimes_{\mathbf{Z}/(p)} \mathbf{Z}/(p)[\xi_1, \xi_2, \dots] & \text{for odd prime } p. \end{cases}$$

where $E(\tau_0, \tau_1, \dots)$ is the exterior algebra, $\deg \xi_i = 2^i - 1$ for $p = 2$, $= 2(p^i - 1)$ for an odd prime p , and $\deg \tau_i = 2p^i - 1$. We have a subalgebra

$$(1.3) \quad P_* = \begin{cases} \mathbf{Z}/(2)[\xi_1^2, \xi_2^2, \dots] & \text{for } p = 2, \\ \mathbf{Z}/(p)[\xi_1, \xi_2, \dots] & \text{for odd prime } p. \end{cases}$$

Let $c : A_* \rightarrow A_*$ be a conjugation. In this paper, we use elements

$$(1.4) \quad \eta_i = \begin{cases} c(\xi_{i+1}) & \text{for } p = 2 \\ c(\tau_i) & \text{for } p > 2 \end{cases} \quad \text{and} \quad t_i = \begin{cases} c(\xi_i^2) & \text{for } p = 2 \\ c(\xi_i) & \text{for } p > 2 \end{cases}.$$

For a coproduct $\Delta : A_* \rightarrow A_* \otimes_{\mathbf{Z}/(p)} A_*$,

$$(1.5) \quad \Delta \tau_n = \tau_n \otimes 1 + \sum_{i=0}^n \xi_{n-i}^{p^i} \otimes \tau_i \quad \text{and} \quad \Delta \xi_n = \sum_{i=0}^n \xi_{n-i}^{p^i} \otimes \xi_i \quad (\xi_0 = 1).$$

Hence,

$$(1.6) \quad \Delta \eta_n = 1 \otimes \eta_n + \sum_{i=0}^n \eta_i \otimes t_{n-i}^{p^i} \quad \text{and} \quad \Delta t_n = \sum_{i=0}^n t_i \otimes t_{n-i}^{p^i} \quad (t_0 = 1).$$

Let $C = \{C^t; \delta : C^t \rightarrow C^{t+1}\}$ be a cochain complex defined by

$$\begin{aligned} C^t &= A_*^t \otimes_{\mathbf{Z}/(p)} P_* = \underbrace{A_* \otimes_{\mathbf{Z}/(p)} \cdots \otimes_{\mathbf{Z}/(p)} A_*}_{t \text{ times}} \otimes_{\mathbf{Z}/(p)} P_* \quad \text{and} \\ (1.7) \quad x_t \otimes \cdots \otimes x_2 \otimes x_1 \otimes m &= x_t \otimes \cdots \otimes x_2 \otimes x_1 \otimes \Delta m \\ &\quad + \sum_{i=1}^t (-1)^i x_t \otimes \cdots \otimes \Delta x_i \otimes \cdots \otimes x_1 \otimes m \\ &\quad (-1)^{t+1} 1 \otimes x_t \otimes \cdots \otimes x_1 \otimes m, \end{aligned}$$

in which $x_i \in A_*$, $m \in P_*$ and $\Delta : N \rightarrow A_* \otimes N$ for $N = A_*$ or P_* . This cochain complex has the cohomology

$$(1.8) \quad \begin{aligned} H^t(C^*; \delta) &= \mathrm{Ext}_{A_*}^t(\mathbf{Z}/(p), P_*) \\ &= \mathbf{Z}/(p)[a_0, a_1, \dots] \quad (a_i \in \mathrm{Ext}^1) \quad (\text{see (2.1-2)}). \end{aligned}$$

At Definition 2.2, we define $X(N) \in C^t$ representing $a_N = a_{n_1} \cdots a_{n_t}$ for any sequence $N = (n_1, \dots, n_t)$ of integers $n_i \geq 0$ (see Proposition 2.3 iii)). Then, for any $\sigma \in F_t$, we define $Y_\sigma(N) \in C^{t-1}$ in Definition 3.1 such that

$$(1.9) \quad \delta Y_\sigma(N) = X(N) - X(\sigma N) \quad (\text{see Proposition 3.3 i)}) \quad \text{and}$$

$$(1.10) \quad Y_{\sigma\tau}(N) = Y_\tau(N) + Y_\sigma(\tau N) \quad (\text{see Proposition 3.4 i)}).$$

Moreover we define $R_\gamma(N) \in C^{t-2}$ for $\gamma \in \langle R1, R2 \rangle$ in Definition 4.1 such that

$$(1.11) \quad \delta R_\gamma(N) = Y_\gamma(N) \quad (\text{see Proposition 4.2 i)}).$$

Definition 1.2. (1) We define submodules of C^t , C^{t-1} and C^{t-2} by taking

$$C_X^t = \langle X(N) \rangle, \quad C_Y^{t-1}(N) = \langle Y_\sigma(\tau N) | \sigma, \tau \in F_t \rangle,$$

$$C_R^{t-2}(N) = \langle R_\gamma(N) | \gamma \in \langle R1, R2 \rangle \rangle,$$

$$C_Y^{t-1} = \cup_N C_Y^{t-1}(N) \quad \text{and} \quad C_R^{t-2} = \cup_N C_R^{t-2}(N)$$

(2) We define a map $\bar{\phi}_{q,N} : F_t \rightarrow C_Y^{t-1}(N)$ by taking

$$\bar{\phi}_{q,N}(e) = 0, \quad \bar{\phi}_{q,N}(\sigma_i) = Y_{\sigma_i}(N), \quad \bar{\phi}_{q,N}(\sigma_i^{-1}) = q^{-1}Y_{\sigma_i}(N) \quad \text{and}$$

$$\bar{\phi}_{q,N}(\sigma_i^{\pm 1}\sigma) = \bar{\phi}_{q,N}(\sigma) + q^{|\sigma|} \bar{\phi}_{q,\sigma N}(\sigma_i^{\pm 1}).$$

At Proposition 5.3, we see that $\bar{\phi}_{q,N}$ is well defined and

$$(1.12) \quad \bar{\phi}_{q,N}(\sigma\tau) = \bar{\phi}_{q,N}(\tau) + q^{|\tau|} \bar{\phi}_{q,\tau N}(\sigma).$$

The definitions of ϕ_q , $\bar{\phi}_{q,N}$ and the following theorem are the main purposes in this paper.

Theorem 1.3.

- i) $C_X^t / \delta C_Y^{t-1} = H^t(C^*; \delta) = \text{Ext}_{A_*}^t(\mathbf{Z}/(p), P_*) = \mathbf{Z}/(p)[a_0, a_1, \dots]$.
- ii) $\bar{\phi}_{1,N}(\sigma) = Y_\sigma(N)$, and so $\bar{\phi}_{1,N}(\langle R1, R2 \rangle) = \delta C_R^{t-2}(N)$.
- iii) $\bar{\phi}_{q,N}(\langle R1, R2 \rangle)$ is a submodule of $C_Y^{t-1}(N)$.

iv) We assume that $n_i \neq n_j$ for any $i \neq j$. If $t \geq 5$ and $q^2 \not\equiv 1 \pmod{p}$ then

$$C_Y^{t-1}(N)/\bar{\phi}_{q,N}(\langle R1, R2 \rangle) = \mathbf{Z}/(p) \quad \text{and}$$

$\bar{\phi}_{q,N}$ induces

$$\phi_q : B_t = F_t/\langle R1, R2 \rangle \rightarrow C_Y^{t-1}(N)/\bar{\phi}_{q,N}(\langle R1, R2 \rangle) = \mathbf{Z}/(p)$$

in Definition 1.1.

The cochain complex C and its cohomology are useful in the field of the stable homotopy theory. Let $H\mathbf{Z}/(p)$ be the Eilenberg-MacLane spectrum and MU_* a ring spectrum representing the complex cobordism theory. Then the Brown-Peterson spectrum BP is a minimal wedge summand of MU localized at a prime p . Now $H\mathbf{Z}/(p)_*(BP) = P_*$ and the Adams spectral sequence $E_2 = \mathrm{Ext}_{A_*}^{s,t}(\mathbf{Z}/(p), P_*) \Rightarrow \pi_*(BP)$ collapses and converges (see [9]). The Adams spectral sequence $\{E(H\mathbf{Z}/(p))_r^{s,t}, d_r^{H\mathbf{Z}/(p)}\}$ with

$$E(H\mathbf{Z}/(p))_2^{s,t} = \mathrm{Ext}_{A_*}^{s,t}(\mathbf{Z}/(p), \mathbf{Z}/(p))$$

and the Novikov-Adams spectral sequence $\{E(BP)_r^{s,t}, d_r^{BP}\}$ with

$$E(BP)_2^{s,t} = \mathrm{Ext}_{BP_* BP}^{s,t}(BP_*, BP_*)$$

are used to calculate the stable homotopy groups of the spheres. For these E_2 -terms, we have the Mahowald and May spectral sequences

$$\{E(\mathrm{Mah})_{u,r}^{s,t}, d_r^{\mathrm{Mah}}\} \quad \text{and} \quad \{E(\mathrm{May})_{u,r}^{s,t}, d_r^{\mathrm{May}}\} \quad \text{with}$$

$$E(\mathrm{Mah})_{u,2}^{s,t} = \mathrm{Ext}_{P_*}^{s,u}(\mathbf{Z}/(p), \mathrm{Ext}_{A_*}^{t,*}(\mathbf{Z}/(p), P_*)) \Rightarrow E(H\mathbf{Z}/(p))_2^{s+t,u} \quad \text{and}$$

$$E(\mathrm{May})_{u,1}^{s,t} = \mathrm{Ext}_{P_*}^{s,u}(\mathbf{Z}/(p), \mathrm{Ext}_{A_*}^{t,*}(\mathbf{Z}/(p), P_*)) \Rightarrow E(BP)_2^{s,u-t}.$$

By [4, 6], we can calculate $d_r^{H\mathbf{Z}/(p)}$ and d_r^{BP} from d_r^{May} and d_r^{Mah} , respectively. Moreover d_r^{Mah} is calculated by using δ of (1.7), $X(N)$, $Y_\sigma(N)$ and $R_\gamma(N)$ by [4]. We shall use the argument in this paper to calculate d_r^{BP} for $p = 2$ in the forthcoming paper.

This paper is organized as follows. We define $X(N)$, $Y_\sigma(N)$ and $R_\gamma(N)$ in § 2, § 3 and § 4, respectively. The maps ϕ_q and $\bar{\phi}_{q,N}$ are argued in § 5.

2. Definition of $X(N)$. In this section, we define an element $X(N) \in C^t$. Then we argue the properties of this element.

Let $C_A = \{C_A^t; \delta^A : C_A^t \rightarrow C_A^{t+1}\}$ be a cochain complex defined by

$$\begin{aligned} C_A^t &= A_*^t \otimes_{\mathbf{Z}/(p)} A_* \quad \text{and} \\ \delta^A(x_t \otimes \cdots \otimes x_2 \otimes x_1 \otimes x_0) &= \sum_{i=0}^t (-1)^i x_t \otimes \cdots \otimes \Delta x_i \otimes \cdots \otimes x_0 \\ &\quad + (-1)^{t+1} 1 \otimes x_t \otimes \cdots \otimes x_0 \quad (x_i \in A_*). \end{aligned}$$

Then we notice that

$$(2.1) \quad C_A \supset C, \quad \delta^A|C = \delta,$$

(2.2)

$$H^t(C_A^*; \delta^A) = \mathrm{Ext}_{A_*}^t(\mathbf{Z}/(p), A_*) = \begin{cases} 0 & \text{for } t > 0, \\ \mathbf{Z}/(p) & \text{for } t = 0 \end{cases} \quad \text{and}$$

(2.3)

$$\delta^A(x \otimes y) = -\delta^A x \otimes y + x \otimes \Delta y \quad \text{for } x \in A_*^t = C_A^{t-1}, y \in A_*.$$

Hence we have the following lemma.

Lemma 2.1. *For $x \in C^t$ ($t > 0$), $\delta x = 0$ if and only if there exists an element $y^A \in C_A^{t-1}$ such that $\delta^A y^A = x$*

Let $B_* = A_*//P_*$ be a quotient algebra, $pr : A_* \rightarrow B_*$ a projection and $\Delta_B : B_* \rightarrow B_* \otimes_{\mathbf{Z}/(p)} B_*$ an induced coproduct. We have a cochain complex

$$C_B = \{C_B^t = B_*^t; \delta^B : C_B^t \rightarrow C_B^{t+1}\}$$

defined by

$$\begin{aligned} \delta^B(x_t \otimes \cdots \otimes x_1) &= x_t \otimes \cdots \otimes x_1 \otimes 1 \\ &\quad + \sum_{i=1}^t (-1)^i x_t \otimes \cdots \otimes \Delta_B x_i \otimes \cdots \otimes x_1 \\ &\quad + (-1)^{t+1} 1 \otimes x_t \otimes \cdots \otimes x_1, \quad (x_i \in B_*). \end{aligned}$$

Now pr induces a cochain homomorphism $pr_* : C = A^t \otimes_{\mathbf{Z}/(p)} P_* \rightarrow C_B = B_*^t \otimes_{\mathbf{Z}/(p)} \mathbf{Z}/(p)$ and an isomorphism

(2.4)

$$pr_* : \mathrm{Ext}_{A_*}^t(\mathbf{Z}/(p), P_*) = H^t(C^*) \xrightarrow{\cong} \mathrm{Ext}_{B_*}^t(\mathbf{Z}/(p), \mathbf{Z}/(p)) = H^t(C_B^*)$$

by the change-of-ring theorem. Then

$$(2.5) \quad \mathrm{Ext}_{B_*}^*(\mathbf{Z}/(p), \mathbf{Z}/(p)) = \mathbf{Z}/(p)[a_0, a_1, \dots]$$

and $a_i \in \mathrm{Ext}_{B_*}^1(\mathbf{Z}/(p), \mathbf{Z}/(p))$ is represented by $pr(\eta_n) \in B_* = C_B^1$, where $\eta_n \in A_*$ is the one of (1.4). Consider elements $x(n) = \eta_n \in C_A^0 = A_*$ and $X(n) = \delta^A x(n) = \sum_{i=0}^n \eta_i \otimes t_{n-i}^{p^i} \in C^1$. $X(n)$ represents a_n since $pr_*(X(n)) = \eta_n \in B_* = C_B^1$. Let $N = (n_1, n_2, \dots, n_t)$ be a sequence of integers with $n_i \geq 0$. We shall define an element $x(N) \in C_A^{t-1} = A_*^t$ so that $\delta^A x(N)$ is included in C^t and represents a monomial $a_N = a_{n_1} a_{n_2} \cdots a_{n_t}$ as follows:

Definition 2.2. Inductively we define $x(N) \in C_A^{t-1}$ by taking $x(n_1) = \eta_{n_1}$ and

$$x(N, n_{t+1}) = x(n_1, \dots, n_t, n_{t+1}) = \sum_{I=0}^N x(I) \otimes t_{N-I}^{p^I} \eta_{n_{t+1}},$$

where

$$\begin{aligned} I &= (i_1, i_2, \dots, i_t), \\ \sum_{I=0}^N &= \sum_{i_1=0}^{n_1} \sum_{i_2=0}^{n_2} \cdots \sum_{i_t=0}^{n_t} \end{aligned}$$

and

$$t_{N-I}^{p^I} = t_{n_1-i_1}^{p^{i_1}} t_{n_2-i_2}^{p^{i_2}} \cdots t_{n_t-i_t}^{p^{i_t}}.$$

Moreover we define

$$X(N) = \sum_{I=0}^N x(I) \otimes t_{N-I}^{p^I} \in C^t,$$

and so

$$x(N, n) = X(N) \eta_n.$$

THE STEENROD ALGEBRA AND BRAID GROUPS

95

We shall prove that $\delta^A x(N) = X(N)$. For this purpose, we notice the following:

(2.6)

$$(a \otimes x)(b \otimes y) = (-1)^{\deg x \cdot \deg b} ab \otimes xy \quad (\text{see [11, Theorem 17.8]}),$$

$$(2.7) \quad \Delta t_{N-I}^{p^I} = \sum_{J=I}^N t_{J-I}^{p^I} \otimes t_{N-J}^{p^J} \quad \text{and}$$

$$(2.8) \quad \Delta t_{N-I}^{p^I} \eta_n = \sum_{J=I}^N t_{J-I}^{p^I} \otimes t_{N-J}^{p^J} \eta_n + \sum_{J=I}^N \sum_{i=0}^n t_{J-I}^{p^I} \eta_i \otimes t_{N-J}^{p^J} t_{n-i}^{p^i}$$

by (1.6). For any expression $f(I)$,

$$(2.9) \quad \sum_{I=0}^N \sum_{J=I}^N f(I) \otimes t_{J-I}^{p^I} \otimes t_{N-J}^{p^J} = \sum_{J=0}^N \sum_{I=0}^J f(I) \otimes t_{J-I}^{p^I} \otimes t_{N-J}^{p^J} \\ = \sum_{I=0}^N \sum_{J=0}^I f(J) \otimes t_{I-J}^{p^J} \otimes t_{N-I}^{p^I}.$$

Proposition 2.3. *For the above $X(N)$, we have the following.*

i) $\delta^A x(N) = X(N)$.

ii) $\Delta_* X(N) = \sum_{I=0}^N X(I) \otimes t_{N-I}^{p^I} \in C^t \otimes P_*$, in which $\Delta_*: C^t \rightarrow C^t \otimes_{\mathbf{Z}/(p)} P_*$
 P_* is the one induced by $\Delta: P_* \rightarrow P_* \otimes_{\mathbf{Z}/(p)} P_* \subset A_* \otimes_{\mathbf{Z}/(p)} P_*$.

iii) $X(N)$ represents a monomial a_N .

Proof. i) We shall use induction on t . For the case of $t = 1$, (1.6) implies this part. By (2.3), (2.8), (2.9) and induction,

$$\begin{aligned} \delta^A x(N, n) &= \delta^A \sum_{I=0}^N x(I) \otimes t_{N-I}^{p^I} \eta_n \\ &= - \sum_{I=0}^N \delta^A x(I) \otimes t_{N-I}^{p^I} \eta_n + \sum_{I=0}^N x(I) \otimes \Delta t_{N-I}^{p^I} \eta_n \\ &= - \sum_{I=0}^N X(I) \otimes t_{N-I}^{p^I} \eta_n + \sum_{I=0}^N \sum_{J=0}^I x(J) \otimes t_{I-J}^{p^J} \otimes t_{N-I}^{p^I} \eta_n \end{aligned}$$

$$+ \sum_{I=0}^N \sum_{J=0}^I \sum_{i=0}^n x(J) \otimes t_{I-J}^{p^J} \eta_i \otimes t_{N-I}^{p^I} t_{n-i}^{p^i}.$$

By definition,

$$X(I) = \sum_{J=0}^I x(J) \otimes t_{I-J}^{p^J}$$

and

$$x(I, i) = \sum_{J=0}^I x(J) \otimes t_{I-J}^{p^J} \eta_i.$$

Hence

$$\delta^A x(N, n) = \sum_{I=0}^N \sum_{i=0}^n x(I, i) \otimes t_{N-I}^{p^I} t_{n-i}^{p^i} = X(N, n).$$

ii) By the definition, (2.7) and (2.9),

$$\begin{aligned} \Delta_* X(N) &= \sum_{I=0}^N x(I) \otimes \sum_{J=I}^N t_{J-I}^{p^I} \otimes t_{N-J}^{p^J} = \sum_{I=0}^N \sum_{J=0}^I x(J) \otimes t_{I-J}^{p^J} \otimes t_{N-I}^{p^I} \\ &= \sum_{I=0}^N X(I) \otimes t_{N-I}^{p^I}. \end{aligned}$$

iii) Let $pr_* = pr \otimes \cdots \otimes pr : C_A^{t-1} = A_*^t \rightarrow C_B^t = B_*^t$ be a homomorphism. By induction, we see that $pr_* x(N, n) = pr_* x(N) \otimes \eta_n = \eta_{n_1} \otimes \cdots \otimes \eta_{n_t} \otimes \eta_n$. This implies this proposition.

3. Definition of $Y_\sigma(N)$. In this section, we define an element $Y_\sigma(N) \in C^{t-1}$ for $\sigma \in F_t$. Then we argue the properties of this element.

Definition 3.1. (1) For $t \geq 0$, we define $y_{t+1}(N, n, m) \in C_A^t$ for $N = (n_1, \dots, n_t)$ by taking

$$y_1(n, m) = -\eta_n \eta_m \quad \text{and}$$

$$y_{t+1}(N, n, m) = -X(N) \eta_n \eta_m = -\sum_{I=0}^N x(I) \otimes t_{N-I}^{p^I} \eta_n \eta_m.$$

Inductively we define $y_i(N, n) \in C_A^{t-1}$ for $1 \leq i \leq t-1$ by taking

$$y_i(N, n) = -\sum_{I=0}^N y_i(I) \otimes t_{N-I}^{p^I} \eta_n.$$

(2) Next we denote $y_e(N) = 0$, $y_{\sigma_i}(N) = y_i(N)$ and $y_{\sigma_i^{-1}}(N) = y_i(N)$ for generators $\sigma_i \in F_t$ and define y_σ inductively by $y_{\sigma_i^{\pm 1}\sigma}(N) = y_\sigma(N) + y_i(\sigma N)$ for $\sigma \in F_t$. Now we take $Y_a(N) = \sum_{I=0}^N y_a(I) \otimes t_{N-I}^{p^I} \in C^{t-1}$ for $a = i$ or σ .

Remark 3.2. In the above definition, we notice that $\eta_n \eta_m = -\eta_m \eta_n$, and so

$$(3.1) \quad y_i(\sigma_i N) = -y_i(N)$$

by induction on t for a fixed i . Then

$$y_{\sigma_i^{-1}\sigma_i\sigma}(N) = y_\sigma(N) + y_i(\sigma N) + y_i(\sigma_i\sigma N) = y_\sigma(N).$$

Hence $y_\sigma(N)$ for any $\sigma \in F_t$ is well-defined.

By (1.6), (2.6) and (2.7), we see that

$$(3.2) \quad \begin{aligned} \Delta \eta_n \eta_m &= (1 \otimes \eta_n + \sum_{i=0}^n \eta_i \otimes t_{n-i}^{p^i})(1 \otimes \eta_m + \sum_{j=0}^m \eta_j \otimes t_{m-j}^{p^j}) \\ &= 1 \otimes \eta_n \eta_m + \sum_{i=0}^n \eta_i \otimes t_{n-i}^{p^i} \eta_m \\ &\quad - \sum_{j=0}^m \eta_j \otimes \eta_n t_{m-j}^{p^j} + \sum_{i=0}^n \sum_{j=0}^m \eta_i \eta_j \otimes t_{n-i}^{p^i} t_{m-j}^{p^j}, \end{aligned}$$

and

(3.3)

$$\begin{aligned} \Delta t_{N-I}^{p^I} \eta_n \eta_m &= \sum_{J=I}^N t_{J-I}^{p^I} \otimes t_{N-J}^{p^J} \eta_n \eta_m + \sum_{J=I}^N \sum_{i=0}^n t_{J-I}^{p^I} \eta_i \otimes t_{N-J}^{p^J} t_{n-i}^{p^i} \eta_m \\ &\quad - \sum_{J=I}^N \sum_{j=0}^m t_{J-I}^{p^I} \eta_j \otimes t_{N-J}^{p^J} \eta_n t_{m-j}^{p^j} \\ &\quad + \sum_{J=I}^N \sum_{i=0}^n \sum_{j=0}^m t_{J-I}^{p^I} \eta_i \eta_j \otimes t_{N-J}^{p^J} t_{n-i}^{p^i} t_{m-j}^{p^j}. \end{aligned}$$

Proposition 3.3. *For the above elements, we have the following.*

i) [(1.9)] $\delta Y_\sigma(N) = X(N) - X(\sigma N)$.

ii) $\Delta_* Y_\sigma(N) = \sum_{I=0}^N Y_\sigma(I) \otimes t_{N-I}^{p^I}$.

Proof. i) In the first place, we prove that

$$(3.4) \quad \delta^A y_i(N) = -x(N) + x(\sigma_i N) + Y_i(N).$$

For example, by (2.3), (2.9) and (3.3),

$$\begin{aligned} \delta^A y_{t+1}(N, n, m) &= \sum_{I=0}^N \delta^A x(I) \otimes t_{N-I}^{p^I} \eta_n \eta_m - \sum_{I=0}^N x(I) \otimes \Delta t_{N-I}^{p^I} \eta_n \eta_m \\ &= \sum_{I=0}^N X(I) \otimes t_{N-I}^{p^I} \eta_n \eta_m \\ &\quad - \sum_{I=0}^N \sum_{J=0}^I x(J) \otimes t_{I-J}^{p^J} \otimes t_{N-I}^{p^I} \eta_n \eta_m \\ &\quad - \sum_{I=0}^N \sum_{J=0}^I \sum_{i=0}^n x(J) \otimes t_{I-J}^{p^J} \eta_i \otimes t_{N-I}^{p^I} t_{n-i}^{p^i} \eta_m \\ &\quad + \sum_{I=0}^N \sum_{J=0}^I \sum_{j=0}^m x(J) \otimes t_{I-J}^{p^J} \eta_j \otimes t_{N-I}^{p^I} \eta_n t_{m-j}^{p^j} \\ &\quad - \sum_{I=0}^N \sum_{J=0}^I \sum_{i=0}^n \sum_{j=0}^m x(J) \otimes t_{I-J}^{p^J} \eta_i \eta_j \otimes t_{N-I}^{p^I} t_{n-i}^{p^i} t_{m-j}^{p^j}. \end{aligned}$$

By the definitions of $X(I)$, $X(I, i)$, $X(I, j)$ and $y_{t+1}(I, i, j)$,

$$\begin{aligned}\delta^A y_{t+1}(N, n, m) &= - \sum_{I=0}^N \sum_{i=0}^n x(I, i) \otimes t_{N-I}^{p^I} t_{n-i}^{p^i} \eta_m \\ &\quad + \sum_{I=0}^N \sum_{j=0}^m x(I, j) \otimes t_{N-I}^{p^I} \eta_n t_{m-j}^{p^j} \\ &\quad + \sum_{I=0}^N \sum_{i=0}^n \sum_{j=0}^m y_{t+1}(I, i, j) \otimes t_{N-I}^{p^I} t_{n-i}^{p^i} t_{m-j}^{p^j} \\ &= -x(N, n, m) + x(N, m, n) + Y_{t+1}(N, n, m).\end{aligned}$$

Then (3.4) for any i is proved by induction on t . In fact, by (2.3), (2.8) and (2.9),

$$\begin{aligned}\delta^A y_i(N, n) &= \sum_{I=0}^N \delta^A y_i(I) \otimes t_{N-I}^{p^I} \eta_n - \sum_{I=0}^N y_i(I) \otimes \Delta t_{N-I}^{p^I} \eta_n \\ &= \sum_{I=0}^N \{-x(I) + x(\sigma_i I) + Y_i(I)\} \otimes t_{N-I}^{p^I} \eta_n \\ &\quad - \sum_{I=0}^N \sum_{J=0}^I y_i(J) \otimes t_{I-J}^{p^J} \otimes t_{N-I}^{p^I} \eta_n \\ &\quad - \sum_{I=0}^N \sum_{J=0}^I \sum_{a=0}^n y_i(J) \otimes t_{I-J}^{p^J} \eta_a \otimes t_{N-I}^{p^I} t_{n-a}^{p^a} \\ &= -x(N, n) + x(\sigma_i N, n) + Y_i(N, n).\end{aligned}$$

Next, by (2.3), (2.9) and (3.4),

$$\begin{aligned}\delta Y_i(N) &= - \sum_{I=0}^N \delta^A y_i(I) \otimes t_{N-I}^{p^I} + \sum_{I=0}^N y_i(I) \otimes \Delta t_{N-I}^{p^I} \\ &= - \sum_{I=0}^N \{-x(I) + x(\sigma_i I) + Y_i(I)\} \otimes t_{N-I}^{p^I} \\ &\quad + \sum_{I=0}^N \sum_{J=0}^I y_i(J) \otimes t_{I-J}^{p^J} \otimes t_{N-I}^{p^I}\end{aligned}$$

$$= X(N) - X(\sigma_i N).$$

In the last place, we assume that i) of this proposition holds for $\sigma \in F_t$, then

$$\begin{aligned} \delta Y_{\sigma_i^{\pm 1}\sigma}(N) &= \delta \left\{ Y_\sigma(N) + Y_{\sigma_i^{\pm 1}}(\sigma N) \right\} \\ &= X(N) - X(\sigma N) + X(\sigma N) - X(\sigma_i^{\pm 1}\sigma N) \\ &= X(N) - X(\sigma_i^{\pm 1}\sigma N). \end{aligned}$$

By induction, we see i) of this proposition.

ii) This is proved by the same manner as ii) of Proposition 2.3.

Next we prepare the following proposition.

Proposition 3.4.

- i) $y_{\sigma\tau}(N) = y_\tau(N) + y_\sigma(\tau N)$, $Y_{\sigma\tau}(N) = Y_\tau(N) + Y_\sigma(\tau N)$,
 $y_i(\sigma_i N) = -y_i(N)$ and $Y_i(\sigma_i N) = -Y_i(N)$.
- ii) $Y_\sigma(N) + Y_{\sigma^{-1}}(\sigma N) = 0$ and $Y_{\sigma^{-1}}(N) = -Y_\sigma(\sigma^{-1}N)$.
- iii) If $\tau\sigma N = \sigma N$ then $Y_{\sigma^{-1}\tau\sigma}(N) = Y_\tau(\sigma N)$.
- iv) $Y_{\sigma_i^n}(N) = (\frac{1-(-1)^n}{2})Y_i(N)$.

- Proof.* i) This is trivial by induction on a length of the word σ .
ii) By i), $Y_\sigma(N) + Y_{\sigma^{-1}}(\sigma N) = Y_{\sigma^{-1}\sigma}(N) = Y_e(N) = 0$.
iii) By i), ii) and the assumption,

$$\begin{aligned} Y_{\sigma^{-1}\tau\sigma}(N) &= Y_{\tau\sigma}(N) + Y_{\sigma^{-1}}(\tau\sigma N) \\ &= Y_\sigma(N) + Y_\tau(\sigma N) + Y_{\sigma^{-1}}(\sigma N) \\ &= Y_\tau(\sigma N). \end{aligned}$$

- iv) By i) and induction,

$$\begin{aligned} Y_{\sigma_i^{n+1}}(N) &= Y_{\sigma_i}(N) + Y_{\sigma_i^n}(\sigma_i N) \\ &= Y_i(N) + (\frac{1-(-1)^n}{2})Y_i(\sigma_i N) \\ &= \{1 - (\frac{1-(-1)^n}{2})\}Y_i(N) \\ &= (\frac{1-(-1)^{n+1}}{2})Y_i(N). \end{aligned}$$

Definition 3.5. We define elements $\gamma_i, \gamma_{i,j} \in F_t$ ($|i - j| \geq 2$) by

$$\gamma_i = \sigma_i^{-1} \sigma_{i+1}^{-1} \sigma_i^{-1} \sigma_{i+1} \sigma_i \sigma_{i+1} \quad \text{and} \quad \gamma_{i,j} = \sigma_i^{-1} \sigma_j^{-1} \sigma_i \sigma_j,$$

and normal subgroups

$$\langle R1 \rangle = \langle \gamma_i \rangle, \quad \langle R2 \rangle = \langle \gamma_{i,j} \rangle \quad \text{and} \quad \langle R1, R2 \rangle = \langle \gamma_i, \gamma_{i,j} \rangle$$

generated by $\gamma_i, \gamma_{i,j}$ for $i + 1 < j$.

Remark 3.6. We notice that $\gamma_{i,j}^{-1} = \gamma_{j,i}$.

Consider $Y_\gamma(N)$ for these elements $\gamma \in \langle R1, R2 \rangle$. The following lemma is shown by the definitions and Proposition 3.4.

Lemma 3.7. *For the above elements, we have the following.*

- i) $\gamma N = N$ for $\gamma \in \langle R1, R2 \rangle$, and so $Y_{\gamma\gamma'}(N) = Y_{\gamma'\gamma}(N) = Y_\gamma(N) + Y_{\gamma'}(N)$ for $\gamma\gamma' \in \langle R1, R2 \rangle$ and $Y_{\sigma^{-1}\gamma\sigma}(N) = Y_\gamma(\sigma N)$ for $\gamma \in \langle R1, R2 \rangle$ and $\sigma \in F_t$.
- ii) $y_{\gamma_i}(N) = y_{i+1}(N) - y_{i+1}(\sigma_i N) + y_{i+1}(\sigma_i \sigma_{i+1} N) - y_i(N) + y_i(\sigma_{i+1} N) - y_i(\sigma_{i+1} \sigma_i N)$ and $Y_{\gamma_i}(N) = Y_{i+1}(N) - Y_{i+1}(\sigma_i N) + Y_{i+1}(\sigma_i \sigma_{i+1} N) - Y_i(N) + Y_i(\sigma_{i+1} N) - Y_i(\sigma_{i+1} \sigma_i N)$.
- iii) $y_{\gamma_{i,j}}(N) = y_j(N) - y_i(N) + y_j(\sigma_i N) + y_i(\sigma_j N)$ and $Y_{\gamma_{i,j}}(N) = Y_j(N) - Y_i(N) + Y_j(\sigma_i N) + Y_i(\sigma_j N)$.

4. Definition of $R_\gamma(N)$. In this section, we define $R_\gamma(N) \in C^{t-2}$ for $\gamma \in \langle R1, R2 \rangle$ and prove (1.11).

Definition 4.1. For $n, m, l \geq 0$, $N = (n_1, \dots, n_t)$ and i ($1 \leq i \leq t - 1$), we define elements $r_{t+1}(N, n, m, l) \in C_A^t$ and $r_{i,t+1}(N, n, m) \in C_A^{t-1}$ as follows:

$$r_1(n, m, l) = \eta_n \eta_m \eta_l, \quad r_{t+1}(N, n, m, l) = X(N) \eta_n \eta_m \eta_l \text{ and}$$

$$r_{i,t+1}(N, n, m) = Y_i(N) \eta_n \eta_m.$$

Then we define $r_i(N) \in C_A^{t-3}$ for $1 \leq i \leq t-2$ and $r_{i,j}(N) \in C_A^{t-3}$ for $1 \leq i \leq j-2 \leq t-3$ by taking inductively

$$r_i(N, n) = \sum_{I=0}^N r_i(I) \otimes t_{N-I}^{p^I} \eta_n \quad \text{for } 1 \leq i \leq t-2 \quad \text{and}$$

$$r_{i,j}(N, n) = \sum_{I=0}^N r_{i,j}(I) \otimes t_{N-I}^{p^I} \eta_n \quad \text{for } 1 \leq i \leq j-2 \leq t-3.$$

Next we take

$$r_{\gamma_i}(N) = r_i(N), r_{\gamma_i^{-1}}(N) = -r_i(N), r_{\gamma_{i,j}}(N) = r_{i,j}(N) \quad \text{and}$$

$$r_{\gamma_{i,j}^{-1}}(N) = -r_{i,j}(N).$$

Then we define $r_{\sigma^{-1}\gamma\pm 1\sigma}(N) = r_{\gamma\pm 1}(\sigma N)$ for $\gamma = \gamma_i, \bar{\gamma}_i, \gamma_{i,j}$ and $\bar{\gamma}_{i,j}$. Now any element $\rho \in \langle R1, R2 \rangle$ is given by $\rho = \rho_1 \cdots \rho_k \cdots \rho_n \in \langle R1, R2 \rangle$ where $\rho_k = \sigma^{-1}\gamma\sigma$ for some σ and $\gamma = \gamma_i^{\pm 1}, \gamma_{i,j}^{\pm 1}$. We define $r_\rho(N) = \sum_{i=1}^n r_{\rho_i}(N)$.

Moreover we take

$$R_a(N) = \sum_{I=0}^N r_a(I) \otimes t_{N-I}^{p^I},$$

where $r_a = r_i, r_{i,j}$ and r_ρ for $\rho \in \langle R1, R2 \rangle$.

We notice that

$$(4.1) \quad \begin{aligned} \Delta \eta_n \eta_m \eta_l &= 1 \otimes \eta_n \eta_m \eta_l + \sum_{i=0}^n \eta_i \otimes t_{n-i}^{p^i} \eta_m \eta_l - \sum_{j=0}^m \eta_j \otimes \eta_n t_{m-j}^{p^j} \eta_l \\ &\quad + \sum_{k=0}^l \eta_k \otimes \eta_n \eta_m t_{l-k}^{p^k} + \sum_{i=0}^n \sum_{j=0}^m \eta_i \eta_j \otimes t_{n-i}^{p^i} t_{m-j}^{p^j} \eta_l \\ &\quad - \sum_{i=0}^n \sum_{k=0}^l \eta_i \eta_k \otimes t_{n-i}^{p^i} \eta_m t_{l-k}^{p^k} + \sum_{j=0}^m \sum_{k=0}^l \eta_j \eta_k \otimes \eta_n t_{m-j}^{p^j} t_{l-k}^{p^k} \\ &\quad + \sum_{i=0}^n \sum_{j=0}^m \sum_{k=0}^l \eta_i \eta_j \eta_k \otimes t_{n-i}^{p^i} t_{m-j}^{p^j} t_{l-k}^{p^k} \end{aligned}$$

by (1.6), (2.6) and (3.2). Hence, by (2.6) and (2.7),

$$\begin{aligned}
 (4.2) \quad \Delta t_{N-I}^{p^I} \eta_n \eta_m \eta_l = & \sum_{J=I}^N t_{J-I}^{p^I} \otimes t_{N-J}^{p^J} \eta_n \eta_m \eta_l \\
 & + \sum_{J=I}^N \sum_{i=0}^n t_{J-I}^{p^I} \eta_i \otimes t_{N-J}^{p^J} t_{n-i}^{p^i} \eta_m \eta_l \\
 & - \sum_{J=I}^N \sum_{j=0}^m t_{J-I}^{p^I} \eta_j \otimes t_{N-J}^{p^J} \eta_n t_{m-j}^{p^j} \eta_l \\
 & + \sum_{J=I}^N \sum_{k=0}^l t_{J-I}^{p^I} \eta_k \otimes t_{N-J}^{p^J} \eta_n \eta_m t_{l-k}^{p^k} \\
 & + \sum_{J=I}^N \sum_{i=0}^n \sum_{j=0}^m t_{J-I}^{p^I} \eta_i \eta_j \otimes t_{N-J}^{p^J} t_{n-i}^{p^i} t_{m-j}^{p^j} \eta_l \\
 & - \sum_{J=I}^N \sum_{i=0}^n \sum_{k=0}^l t_{J-I}^{p^I} \eta_i \eta_k \otimes t_{N-J}^{p^J} t_{n-i}^{p^i} \eta_m t_{l-k}^{p^k} \\
 & + \sum_{J=I}^N \sum_{j=0}^m \sum_{k=0}^l t_{J-I}^{p^I} \eta_j \eta_k \otimes t_{N-J}^{p^J} \eta_n t_{m-j}^{p^j} t_{l-k}^{p^k} \\
 & + \sum_{J=I}^N \sum_{i=0}^n \sum_{j=0}^m \sum_{k=0}^l t_{J-I}^{p^I} \eta_i \eta_j \eta_k \otimes t_{N-J}^{p^J} t_{n-i}^{p^i} t_{m-j}^{p^j} t_{l-k}^{p^k}.
 \end{aligned}$$

Now we have (1.11) as follows:

Proposition 4.2.

- i) [(1.11)] $\delta R_\gamma(N) = Y_\gamma(N)$ for $\gamma \in \langle R1, R2 \rangle$.
- ii) $\Delta_* R_\gamma(N) = \sum_{I=0}^N R_\gamma(I) \otimes t_{N-I}^{p^I}$ for $\gamma \in \langle R1, R2 \rangle$.
- iii) $R_{\sigma^{-1}\gamma\sigma}(N) = R_\gamma(\sigma N)$
 $R_{\gamma'\gamma}(N) = R_{\gamma\gamma'}(N) = R_\gamma(N) + R_{\gamma'}(N)$ for $\gamma, \gamma' \in \langle R1, R2 \rangle$ and
 $\sigma \in F_t$.

Proof. i) By using Proposition 2.3 i), Lemma 3.7 ii), (2.3), (2.9) and (4.2), we see that

$$\begin{aligned}
 \delta^A r_{t+1}(N, n, m, l) &= - \sum_{I=0}^N \delta^A x(I) \otimes t_{N-I}^{p^I} \eta_n \eta_m \eta_l + \sum_{I=0}^N x(I) \otimes \Delta t_{N-I}^{p^I} \eta_n \eta_m \eta_l \\
 &= \sum_{I=0}^N \sum_{i=0}^n x(I, i) \otimes t_{N-I}^{p^I} t_{n-i}^{p^i} \eta_m \eta_l \\
 &\quad - \sum_{I=0}^N \sum_{j=0}^m x(I, j) \otimes t_{N-I}^{p^I} \eta_n t_{m-j}^{p^j} \eta_l \\
 &\quad + \sum_{I=0}^N \sum_{k=0}^l x(I, k) \otimes t_{N-I}^{p^I} \eta_n \eta_m t_{l-k}^{p^k} \\
 &\quad + \sum_{I=0}^N \sum_{i=0}^n \sum_{j=0}^m y_{t+1}(I, i, j) \otimes t_{N-I}^{p^I} t_{n-i}^{p^i} t_{m-j}^{p^j} \eta_l \\
 &\quad - \sum_{I=0}^N \sum_{i=0}^n \sum_{k=0}^l y_{t+1}(I, i, k) \otimes t_{N-I}^{p^I} t_{n-i}^{p^i} \eta_m t_{l-k}^{p^k} \\
 &\quad + \sum_{I=0}^N \sum_{j=0}^m \sum_{k=0}^l y_{t+1}(I, j, k) \otimes t_{N-I}^{p^I} \eta_n t_{m-j}^{p^j} t_{l-k}^{p^k} \\
 &\quad + \sum_{I=0}^N \sum_{i=0}^n \sum_{j=0}^m \sum_{k=0}^l r_{t+1}(I, i, j, k) \otimes t_{N-I}^{p^I} t_{n-i}^{p^i} t_{m-j}^{p^j} t_{l-k}^{p^k} \\
 &= -y_{t+2}(N, n, m, l) + y_{t+2}(N, m, n, l) \\
 &\quad - y_{t+2}(N, l, n, m) + y_{t+1}(N, n, m, l) \\
 &\quad - y_{t+1}(N, n, l, m) + y_{t+1}(N, m, l, n) \\
 &\quad + R_{t+1}(N, n, m, l) \\
 &= -y_{\gamma_{t+1}}(N, n, m, l) + R_{t+1}(N, n, m, l).
 \end{aligned}$$

Then, by induction, we can prove that

$$(4.3) \quad \delta^A r_i(N) = -y_{\gamma_i}(N) + R_i(N).$$

In fact, by (2.3), (2.8) and (2.9),

$$\delta^A r_i(N, n) = - \sum_{I=0}^N \delta^A r_i(I) \otimes t_{N-I}^{p^I} \eta_n + \sum_{I=0}^N r_i(I) \otimes \Delta t_{N-I}^{p^I} \eta_n$$

$$\begin{aligned}
&= - \sum_{I=0}^N \{-y_{\gamma_i}(I) + R_i(I)\} \otimes t_{N-I}^{p^I} \eta_n \\
&\quad + \sum_{I=0}^N R_i(I) \otimes t_{N-I}^{p^I} \eta_n \\
&\quad + \sum_{I=0}^N \sum_{a=0}^n r_i(I, a) \otimes t_{N-I}^{p^I} t_{n-a}^{p^a} \\
&= -y_{\gamma_i}(N, n) + R_i(N, n).
\end{aligned}$$

Now,

$$\begin{aligned}
\delta R_i(N) &= - \sum_{I=0}^N \delta^A r_i(I) \otimes t_{N-I}^{p^I} + \sum_{I=0}^N r_i(I) \otimes \Delta t_{N-I}^{p^I} \\
&= - \sum_{I=0}^N \{-y_{\gamma_i}(I) + R_i(I)\} \otimes t_{N-I}^{p^I} \\
&\quad + \sum_{I=0}^N \sum_{J=0}^I r_i(J) \otimes t_{I-J}^{p^J} \otimes t_{N-I}^{p^I} \\
&= Y_{\gamma_i}(N).
\end{aligned}$$

Next, by Lemma 3.7, (2.3), (2.9), (3.3) and (3.4),

$$\begin{aligned}
\delta^A r_{i,t+1}(N, n, m) &= - \sum_{I=0}^N \delta^A y_i(I) \otimes t_{N-I}^{p^I} \eta_n \eta_m + \sum_{I=0}^N y_i(I) \otimes \Delta t_{N-I}^{p^I} \eta_n \eta_m \\
&= - \sum_{I=0}^N \{-x(I) + x(\sigma_i I) + Y_i(I)\} \otimes t_{N-I}^{p^I} \eta_n \eta_m \\
&\quad + \sum_{I=0}^N Y_i(I) \otimes t_{N-I}^{p^I} \eta_n \eta_m \\
&\quad + \sum_{I=0}^N \sum_{a=0}^n y_i(I, a) \otimes t_{N-I}^{p^I} t_{n-a}^{p^a} \eta_m \\
&\quad - \sum_{I=0}^N \sum_{b=0}^m y_i(I, b) \otimes t_{N-I}^{p^I} \eta_m t_{m-b}^{p^b}
\end{aligned}$$

$$\begin{aligned}
 & + \sum_{I=0}^N \sum_{a=0}^n \sum_{b=0}^m r_{i,t+1}(I, a, b) \otimes t_{N-I}^{p^I} t_{n-a}^{p^a} t_{m-b}^{p^b} \\
 & = -y_{t+1}(N, n, m) + y_{t+1}(\sigma_i N, n, m) \\
 & \quad + y_i(N, n, m) - y_i(N, m, n) + R_{i,t+1}(N, n, m) \\
 & = -y_{\gamma_{i,t+1}}(N, n, m) + R_{i,t+1}(N, n, m).
 \end{aligned}$$

Then, by induction, we can prove that

$$(4.4) \quad \delta^A r_{i,j}(N) = -y_{\gamma_{i,j}}(N) + R_{i,j}(N).$$

In fact, by (2.3), (2.8) and (2.9),

$$\begin{aligned}
 \delta^A r_{i,j}(N, n) & = - \sum_{I=0}^N \delta^A r_{i,j}(I) \otimes t_{N-I}^{p^I} \eta_n + \sum_{I=0}^N r_{i,j}(I) \otimes \Delta t_{N-I}^{p^I} \eta_n \\
 & = - \sum_{I=0}^N \{-y_{\gamma_{i,j}}(I) + R_{i,j}(I)\} \otimes t_{N-I}^{p^I} \eta_n \\
 & \quad + \sum_{I=0}^N R_{i,j}(I) \otimes t_{N-I}^{p^I} \eta_n \\
 & \quad + \sum_{I=0}^N \sum_{a=0}^n r_{i,j}(I, a) \otimes t_{N-I}^{p^I} t_{n-a}^{p^a} \\
 & = -y_{\gamma_{i,j}}(N, n) + R_{i,j}(N, n).
 \end{aligned}$$

Now we see that

$$\begin{aligned}
 \delta R_{i,j}(N) & = - \sum_{I=0}^N \delta^A r_{i,j}(I) \otimes t_{N-I}^{p^I} + \sum_{I=0}^N r_{i,j}(I) \otimes \Delta t_{N-I}^{p^I} \\
 & = - \sum_{I=0}^N \{-y_{\gamma_{i,j}}(I) + R_{i,j}(I)\} \otimes t_{N-I}^{p^I} \\
 & \quad + \sum_{I=0}^N \sum_{J=0}^I r_{i,j}(J) \otimes t_{I-J}^{p^J} \otimes t_{N-I}^{p^I} \\
 & = Y_{\gamma_{i,j}}(N).
 \end{aligned}$$

By the definition of $R_\gamma(N)$ and the above results, Lemma 3.7 implies this proposition i).

- ii) This is proved by the same manner as ii) of Proposition 2.3.
- iii) By the definition, this is trivial.

5. ϕ_q and $\bar{\phi}_{q,N}$. In this section, we argue the properties of ϕ_q and $\bar{\phi}_{q,N}$ and prove Theorem 1.3.

In the first place, we prepare the following lemma.

Lemma 5.1.

- i) $\bar{\phi}_q(\sigma_i^{-1}\sigma_i\sigma) = \bar{\phi}_q(\sigma) = \bar{\phi}_q(\sigma_i\sigma_i^{-1}\sigma)$.
- ii) $\bar{\phi}_q(\sigma\tau) = \bar{\phi}_q(\tau) + (-q)^{|\tau|}\bar{\phi}_q(\sigma)$.
- iii) $\bar{\phi}_q((\sigma\tau)\delta) = \bar{\phi}_q(\sigma(\tau\delta)) = \bar{\phi}_q(\delta) + (-q)^{|\delta|}\bar{\phi}_q(\tau) + (-q)^{|\tau\delta|}\bar{\phi}_q(\sigma)$.
- iv) $\bar{\phi}_q(\sigma^{-1}) = -(-q)^{|\sigma|}\bar{\phi}_q(\sigma)$.
- v) $\bar{\phi}_q(\sigma^{-1}\gamma\sigma) = (-q)^{|\sigma|}\bar{\phi}_q(\gamma) + \bar{\phi}_q(\sigma) - (-q)^{|\gamma|}\bar{\phi}_q(\sigma)$.
- vi) $\bar{\phi}_q(\gamma_i) = 0$ and $\bar{\phi}_q(\gamma_{i,j}) = 0$.
- vii) For $\gamma \in \langle R1, R2 \rangle$, $\bar{\phi}_q(\gamma) = 0$ and $\bar{\phi}_q(\gamma\sigma) = \bar{\phi}_q(\sigma) = \bar{\phi}_q(\sigma\gamma)$

Proof. i) By definition,

$$\begin{aligned} \bar{\phi}_q(\sigma_i^{-1}\sigma_i\sigma) &= \bar{\phi}_q(\sigma_i\sigma) + (-q)^{|\sigma|+1}q^{-1} \\ &= \bar{\phi}_q(\sigma) + (-q)^{|\sigma|} + (-q)^{|\sigma|+1}q^{-1} = \bar{\phi}_q(\sigma). \end{aligned}$$

As the same way, we see $\bar{\phi}_q(\sigma_i\sigma_i^{-1}\sigma) = \bar{\phi}_q(\sigma)$.

ii) We assume $\bar{\phi}_q(\sigma\tau) = \bar{\phi}_q(\tau) + (-q)^{|\tau|}\bar{\phi}_q(\sigma)$. Then, by definition,

$$\begin{aligned} \bar{\phi}_q(\sigma_i\sigma\tau) &= \bar{\phi}_q(\sigma\tau) + (-q)^{|\sigma\tau|} \\ &= \bar{\phi}_q(\tau) + (-q)^{|\tau|}\bar{\phi}_q(\sigma) + (-q)^{|\tau|+|\sigma|} = \bar{\phi}_q(\tau) + (-q)^{|\tau|}\bar{\phi}_q(\sigma_i\sigma). \end{aligned}$$

As the same way, $\bar{\phi}_q(\sigma_i^{-1}\sigma\tau) = \bar{\phi}_q(\tau) + (-q)^{|\tau|}\bar{\phi}_q(\sigma_i^{-1}\sigma)$. By induction on the word σ , we see ii).

iii) By ii), this is trivial.

iv) By ii), $\bar{\phi}_q(\sigma^{-1}) + (-q)^{-|\sigma|}\bar{\phi}_q(\sigma) = \bar{\phi}_q(\sigma\sigma^{-1}) = \bar{\phi}_q(e) = 0$.

v) This is easy by iii) and iv).

vi) By ii),

$$\begin{aligned}\bar{\phi}_q(\gamma_i) &= \bar{\phi}_q(\sigma_{i+1}) - q\bar{\phi}_q(\sigma_i) + q^2\bar{\phi}_q(\sigma_{i+1}) \\ &\quad - q^3\bar{\phi}_q(\sigma_i^{-1}) + q^2\bar{\phi}_q(\sigma_{i+1}^{-1}) - q\bar{\phi}_q(\sigma_i) \\ &= 1 - q + q^2 - q^2 + q - 1 = 0\end{aligned}$$

By the same way, we see the another equation.

vii) is seen by ii), v), vi) and $|\gamma| = 0$.

This lemma i) implies that $\bar{\phi}_q$ is well defined. By vii), $\bar{\phi}_q$ induces

$$\phi_q : B_t = F_t / \langle R1, R2 \rangle \rightarrow \mathbf{Z}/(p).$$

Hence we have the following.

Proposition 5.2.

i) ϕ_q is well defined.

ii) $\phi_q(\sigma\tau) = \phi_q(\tau) + (-q)^{|\tau|}\phi_q(\sigma)$.

iii) $\phi_q(\sigma^{-1}) = -(-q)^{-|\sigma|}\phi_q(\sigma)$.

iv) $\phi_q(\sigma^{-1}\gamma\sigma) = (-q)^{|\sigma|}\phi_q(\gamma) + \phi_q(\sigma) - (-q)^{|\gamma|}\phi_q(\sigma)$.

Next consider the map $\bar{\phi}_{q,N} : F_t \rightarrow C_Y^{t-1}(N)$ of Definition 1.2 (2).

Proposition 5.3.

i) $\bar{\phi}_{q,N}(\sigma_i^{-1}\sigma_i\sigma) = \bar{\phi}_{q,N}(\sigma) = \bar{\phi}_{q,N}(\sigma_i\sigma_i^{-1}\sigma)$. Hence $\bar{\phi}_{q,N}$ is well defined.

ii) $\bar{\phi}_{q,N}(\sigma\tau) = \bar{\phi}_{q,N}(\tau) + q^{|\tau|}\bar{\phi}_{q,\tau N}(\sigma)$.

iii) $\bar{\phi}_{q,N}((\sigma\tau)\delta) = \bar{\phi}_{q,N}(\sigma(\tau\delta)) = \bar{\phi}_{q,N}(\delta) + q^{|\delta|}\bar{\phi}_{q,\delta N}(\tau) + q^{|\tau\delta|}\bar{\phi}_{q,\tau\delta N}(\sigma)$.

iv) $\bar{\phi}_{q,N}(\sigma^{-1}) = -q^{-|\sigma|}\bar{\phi}_{q,\sigma^{-1}N}(\sigma)$.

v) $\bar{\phi}_{q,N}(\sigma^{-1}\gamma\sigma) = q^{|\sigma|}\bar{\phi}_{q,\sigma N}(\gamma) + \bar{\phi}_{q,N}(\sigma) - q^{|\gamma|}\bar{\phi}_{q,\sigma^{-1}\gamma\sigma N}(\sigma)$.

Proof. i) By definition and Proposition 3.4 i),

$$\begin{aligned}\bar{\phi}_{q,N}(\sigma_i^{-1}\sigma_i\sigma) &= \bar{\phi}_{q,N}(\sigma) + q^{|\sigma|}\bar{\phi}_{q,\sigma N}(\sigma_i) + q^{|\sigma|+1}\bar{\phi}_{q,\sigma_i\sigma N}(\sigma_i^{-1}) \\ &= \bar{\phi}_{q,N}(\sigma) + q^{|\sigma|}Y_i(\sigma N) + q^{|\sigma|+1}q^{-1}Y_i(\sigma_i\sigma N) \\ &= \bar{\phi}_{q,N}(\sigma)\end{aligned}$$

The another equation is seen by the same way.

ii) By induction on word σ and

$$\begin{aligned}\bar{\phi}_{q,N}(\sigma_i^{\pm 1}\sigma\tau) &= \{\bar{\phi}_{q,N}(\tau) + q^{|\tau|}\bar{\phi}_{q,\tau N}(\sigma)\} + q^{|\sigma|+|\tau|}\bar{\phi}_{q,\sigma\tau N}(\sigma_i^{\pm 1}) \\ &= \bar{\phi}_{q,N}(\tau) + q^{|\tau|}\bar{\phi}_{q,\tau N}(\sigma_i^{\pm 1}\sigma),\end{aligned}$$

we see ii).

iii) is given by ii).

iv) is given by ii) and $\bar{\phi}_{q,N}(\sigma_i\sigma_i^{-1}) = \bar{\phi}_{q,N}(e) = 0$.

v) is given by iii) and iv).

Now we argue $\bar{\phi}_{q,N}(\langle R1, R2 \rangle)$. The following lemma is proved by Proposition 5.3.

Lemma 5.4. *For $\gamma, \gamma' \in \langle R1, R2 \rangle$ and $\sigma \in F_t$, we have the following.*

i) $\bar{\phi}_{q,N}(\sigma^{-1}\gamma\sigma) = q^{|\sigma|}\bar{\phi}_{q,\sigma N}(\gamma)$.

ii) $\bar{\phi}_{q,N}(\gamma\gamma') = \bar{\phi}_{q,N}(\gamma) + \bar{\phi}_{q,N}(\gamma')$. Hence $\bar{\phi}_{q,N}(\langle R1, R2 \rangle)$ is a submodule of $C_Y^{t-1}(N)$.

iii)

$$\begin{aligned}\bar{\phi}_{q,N}(\sigma\gamma) &= \bar{\phi}_{q,N}(\sigma) + \bar{\phi}_{q,N}(\gamma) \quad \text{and} \\ \bar{\phi}_{q,N}(\gamma\sigma) &= \bar{\phi}_{q,N}(\sigma(\sigma^{-1}\gamma\sigma)) = \bar{\phi}_{q,N}(\sigma) + \bar{\phi}_{q,N}(\sigma^{-1}\gamma\sigma).\end{aligned}$$

Hence $\bar{\phi}_{q,N}$ induces a map

$$\phi_{q,N} : B_t = F_t/\langle R1, R2 \rangle \rightarrow C_Y^{t-1}(N)/\bar{\phi}_{q,N}(\langle R1, R2 \rangle).$$

iv)

$$\begin{aligned}\bar{\phi}_{q,N}(\gamma_i) &= Y_{i+1}(N) - Y_i(N) - q\{Y_{i+1}(\sigma_i N) - Y_i(\sigma_{i+1} N)\} \\ &\quad + q^2\{Y_{i+1}(\sigma_i \sigma_{i+1} N) - Y_i(\sigma_{i+1} \sigma_i N)\} \quad \text{and} \\ \bar{\phi}_{q,N}(\gamma_{i,j}) &= Y_j(N) - Y_i(N) - q\{Y_j(\sigma_i N) - Y_i(\sigma_j N)\}.\end{aligned}$$

Next we define elements $Z_{i,\sigma}(N) = Y_1(N) - (-1)^{|\sigma|}Y_i(\sigma N)$ and a module $C_Z^{t-1}(N) = \langle Z_{i,\sigma}(N) \rangle$.

Lemma 5.5.

- i) If $p = 2$ then $C_Y^{t-1}(N)/C_Z^{t-1}(N) = \mathbf{Z}/(2)$. If $p > 2$ and $n_i \neq n_j$ for any $i \neq j$ then $C_Y^{t-1}(N)/C_Z^{t-1}(N) = \mathbf{Z}/(p)$. If $p > 2$ and $n_i = n_j$ for some $i \neq j$ then $C_Y^{t-1}(N)/C_Z^{t-1}(N) = 0$.
- ii) $\bar{\phi}_{q,N}(\langle R1, R2 \rangle) \subset C_Z^{t-1}(N)$.
- iii) If $q^2 \not\equiv 1 \pmod{p}$ and $t \geq 5$ then $\bar{\phi}_{q,N}(\langle R1, R2 \rangle) = C_Z^{t-1}(N)$.

Proof. i) $\{Y_i(\sigma N) | \sigma \in F_t\}$ is a basis of $C_Y^{t-1}(N)$, and so is $\{Y_1(N)\} \cup \{Z_{i,\sigma}(N) | \sigma \in F_t\}$. Hence $C_Y^{t-1}(N)/C_Z^{t-1}(N)$ has a generator $Y_1(N)$. If $p = 2$ then $Y_i(\sigma N) \neq 0$. Therefore $C_Y^{t-1}(N)/C_Z^{t-1}(N) = \mathbf{Z}/(2)$.

We assume $p > 3$. If $n_i \neq n_j$ for any $i \neq j$ then $Y_i(\sigma N) \neq 0$ for any i and $\sigma \in F_t$, and so $C_Y^{t-1}(N)/C_Z^{t-1}(N) = \mathbf{Z}/(p)$. If $n_i = n_j$ for some $i \neq j$ then there exists an element $\sigma \in F_t$ so that $n_{\bar{\sigma}^{-1}(1)} = n_{\bar{\sigma}^{-1}(2)}$, i.e., $\sigma_i \sigma N = \sigma N$, and so $Y_1(\sigma N) = Y_1(\sigma_1 \sigma N) = -Y_1(\sigma N)$. Hence $Y_1(\sigma N) = 0$ by $p > 2$ and $Y_1(N) = Z_{1,\sigma}(N) \in C_Z^{t-1}(N)$. Thus $C_Y^{t-1}(N)/C_Z^{t-1}(N) = 0$.

ii) By definition, if $|\sigma| = |\tau|$ then $Y_i(\sigma N) - Y_j(\tau N) = (-1)^{|\sigma|}\{Z_{j,\tau}(N) - Z_{i,\sigma}(N)\}$. We see ii) by Lemma 5.4.

iii) By Lemma 5.4,

$$\begin{aligned} \bar{\phi}_{q,N}(\sigma^{-1}\gamma_{i,j}\sigma) &= q^{|\sigma|}\{Y_j(\sigma N) - Y_i(\sigma N)\} \\ &\quad - q^{|\sigma|+1}\{Y_j(\sigma_i \sigma N) - Y_i(\sigma_j \sigma N)\} \end{aligned}$$

and

$$\begin{aligned} \bar{\phi}_{q,N}((\sigma_i \sigma_j \sigma)^{-1}\gamma_{i,j}\sigma_i \sigma_j \sigma) &= q^{|\sigma|+2}\{Y_j(\sigma_i \sigma_j \sigma N) - Y_i(\sigma_i \sigma_j \sigma N)\} \\ &\quad - q^{|\sigma|+3}\{Y_j(\sigma_i \sigma_i \sigma_j \sigma N) - Y_i(\sigma_j \sigma_i \sigma_j \sigma N)\} \\ &= q^{|\sigma|+3}\{Y_j(\sigma N) - Y_i(\sigma N)\} \\ &\quad - q^{|\sigma|+2}\{Y_j(\sigma_i \sigma N) - Y_i(\sigma_j \sigma N)\}. \end{aligned}$$

Therefore

$$\begin{aligned} \bar{\phi}_{q,N}(\sigma^{-1}\gamma_{i,j}\sigma) - q^{-1}\bar{\phi}_{q,N}((\sigma_i \sigma_j \sigma)^{-1}\gamma_{i,j}\sigma_i \sigma_j \sigma) \\ = (1 - q^2)q^{|\sigma|}\{Y_j(\sigma N) - Y_i(\sigma N)\}, \end{aligned}$$

and so $Y_j(\sigma N) \equiv Y_i(\sigma N) \pmod{\bar{\phi}_{q,N}(\langle R1, R2 \rangle)}$ for $|i - j| \geq 2$ if $q^2 \not\equiv 1 \pmod{p}$. Hence $Y_i(\sigma N) \equiv Y_1(\sigma N)$ for $i \geq 3$. By $t \geq 5$, $Y_2(\sigma N) \equiv Y_4(\sigma N) \equiv Y_1(\sigma N)$. Thus

$$(5.1) \quad Y_i(\sigma N) \equiv Y_1(\sigma N) \pmod{\bar{\phi}_{q,N}(\langle R1, R2 \rangle)} \text{ for any } i.$$

By this equation,

$$Y_i(\sigma_j^{\pm 1}\sigma N) \equiv Y_1(\sigma_j^{\pm 1}\sigma N) \equiv Y_j(\sigma_j^{\pm 1}\sigma N) = -Y_j(\sigma N) \equiv -Y_1(\sigma N).$$

By induction, $Y_i(\sigma N) \equiv (-1)^{|\sigma|}Y_1(N) \pmod{\bar{\phi}_{q,N}(\langle R1, R2 \rangle)}$. Thus

$$Z_{i,\sigma}(N) \in \bar{\phi}_{q,N}(\langle R1, R2 \rangle).$$

Now we can prove Theorem 1.3.

Proof of Theorem 1.3. i-ii) are trivial by the definitions, (1.8) and Proposition 3.3.

iii) is Lemma 5.4 ii).

iv) We define a map $\psi : C_Y^{t-1}(N) \rightarrow \mathbf{Z}/(p)$ by taking $\psi(Y_i(\sigma N)) = (-1)^{|\sigma|}$. If $n_i \neq n_j$ for any $i \neq j$ then ψ is well defined. By Definition 1.1 and 1.2, $\bar{\phi}_q = \psi \bar{\phi}_{q,N}$. Now $\text{Ker } \psi = C_Z^{t-1}(N) = \bar{\phi}_{q,N}(\langle R1, R2 \rangle)$ by Lemma 5.5, and so $\bar{\phi}_{q,N}$ induces ϕ_q .

REFERENCES

- [1] J. F. ADAMS: Stable Homotopy and Generalised Homology, Univ. of Chicago Press, Chicago Illinois London, 1974.
- [2] G. BURDE and H. ZIESCHANG: Knots, Walter de Gruyter, Berlin New York, 1985.
- [3] H. CARTAN and S. EILENBERG: Homological Algebra, Princeton Univ. Press, Princeton New Jersey, 1956.
- [4] M. HIKIDA: Relations between several Adams spectral sequences, Hiroshima Math. J. **19** (1989), 37–76.
- [5] M. HIKIDA and K. SHIMOMURA: An exact sequence related to Adams-Novikov E_2 -terms of a cofibering, J. Math. Soc. Japan **46** (1994), 645–661.
- [6] H. R. MILLER: On relations between Adams spectral sequences, with an application to the stable homotopy of a Moore space, J. Pure appl. Algebra **20** (1981), 287–312.
- [7] J. W. MILNOR: The Steenrod algebra and its dual, Ann. of Math. **67** (1958), 150–171.
- [8] J. W. MILNOR and J. C. MOORE: On the structure of Hopf algebras, Ann. of Math. **81** (1965), 211–264.
- [9] D. C. RAVENEL: Complex Cobordism and Stable Homotopy Groups of Spheres, Academic Press, New York, 1986.

- [10] N. E. STEENROD and D. B. A. EPSTEIN: Cohomology Operations, Annals of Math. Studies No. 50, Princeton Univ. Press, Princeton, 1962.
- [11] R. M. SWITZER: Algebraic Topology, Homotopy and Homology, Springer-Verlag, Berlin New York, 1975.

HIROSHIMA PREFECTURAL UNIVERSITY
SHOBARA-SHI, HIROSHIMA 727 JAPAN
E-mail: hikida@bus.hiroshima-pu.ac.jp

(*Received July 8, 1997*)