

# *Mathematical Journal of Okayama University*

---

*Volume 46, Issue 1*

2004

*Article 34*

JANUARY 2004

---

## Results on prime near-ring with $(\alpha, \gamma)$ -derivation

Oznur Golbasi\*

Neset Aydin<sup>†</sup>

\*Cumhuriyet University

<sup>†</sup>Canakkale 18 Mart University

Copyright ©2004 by the authors. *Mathematical Journal of Okayama University* is produced by The Berkeley Electronic Press (bepress). <http://escholarship.lib.okayama-u.ac.jp/mjou>

# Results on prime near-ring with $(\alpha, \gamma)$ -derivation

Oznur Golbasi and Neset Aydin

## Abstract

Let  $N$  be a prime left near-ring with multiplicative center  $Z$ ; and  $D$  be a  $(\alpha, \gamma)$ -derivation such that  $\delta D = D\delta$  and  $\Gamma D = D\Gamma$  (i) If  $D(N) \subset Z$ ; or  $[D(N); D(N)] = 0$  or  $[D(N); D(N)]\sigma, \gamma = 0$ ; then  $(N; +)$  is abelian. (ii) If  $N$  is 2-torsion free,  $d_1$  is a  $(\alpha, \gamma)$ -derivation and  $d_2$  is a derivation on  $N$  such that  $d_1 d_2(N) = 0$ , then  $d_1 = 0$  or  $d_2 = 0$ .

**KEYWORDS:** Prime Near-Ring, Derivation,  $(\alpha, \gamma)$ -Derivation.

Math. J. Okayama Univ. **46** (2004), 1–7

## RESULTS ON PRIME NEAR-RING WITH $(\sigma, \tau)$ -DERIVATION

ÖZNUR GÖLBAŞI AND NEŞET AYDIN

**ABSTRACT.** Let  $N$  be a prime left near-ring with multiplicative center  $Z$ , and  $D$  be a  $(\sigma, \tau)$ -derivation such that  $\sigma D = D\sigma$  and  $\tau D = D\tau$ . (i) If  $D(N) \subset Z$ , or  $[D(N), D(N)] = 0$  or  $[D(N), D(N)]_{\sigma, \tau} = 0$ , then  $(N, +)$  is abelian. (ii) If  $N$  is 2-torsion free,  $d_1$  is a  $(\sigma, \tau)$ -derivation and  $d_2$  is a derivation on  $N$  such that  $d_1 d_2(N) = 0$ , then  $d_1 = 0$  or  $d_2 = 0$ .

### 1. INTRODUCTION

Recently, some results concerning commutativity in prime near-rings with derivation have been generalized in several ways. The primary purpose of this paper is to generalize some results obtained by H. E. Bell and G. Mason [1], and A. A. M. Kamal [2].

Throughout this paper,  $N$  will denote a zero-symmetric left near-ring with multiplicative center  $Z$ .  $N$  is called a prime near-ring if  $aNb = \{0\}$  implies that  $a = 0$  or  $b = 0$ . Let  $\sigma$  and  $\tau$  be two near-ring automorphisms of  $N$ . An additive mapping  $D : N \rightarrow N$  is called a  $(\sigma, \tau)$ -derivation if  $D(xy) = \tau(x)D(y) + D(x)\sigma(y)$  holds for all  $x, y \in N$ . For  $x, y \in N$ , the symbol  $[x, y]$  will denote  $xy - yx$ , while the symbol  $(x, y)$  will denote the additive-group commutator  $x + y - x - y$ . Given  $x, y \in N$ , we write  $[x, y]_{\sigma, \tau} = x\sigma(y) - \tau(y)x$ ; in particular  $[x, y]_{1,1} = [x, y]$ , in the usual sense. As for terminologies used here without mention, we refer to G. Pilz [3].

### 2. RESULTS

We begin with two quite general and useful lemmas.

**Lemma 1.** *Let  $D$  be a  $(\sigma, \tau)$ -derivation of near ring  $N$ . Then  $D(xy) = D(x)\sigma(y) + \tau(x)D(y)$  for all  $x, y \in N$ .*

*Proof.* Note that

$$\begin{aligned} D(x(y + y)) &= \tau(x)D(y + y) + D(x)\sigma(y + y) \\ &= \tau(x)D(y) + \tau(x)D(y) + D(x)\sigma(y) + D(x)\sigma(y), \end{aligned}$$

and

$$D(xy + xy) = \tau(x)D(y) + D(x)\sigma(y) + \tau(x)D(y) + D(x)\sigma(y).$$

---

*Mathematics Subject Classification.* 16Y30, 16N60, 16W25.

*Key words and phrases.* Prime Near-Ring, Derivation,  $(\sigma, \tau)$ -Derivation.

Comparing these two expressions, one can obtain

$$\tau(x)D(y) + D(x)\sigma(y) = D(x)\sigma(y) + \tau(x)D(y)$$

and so,

$$D(xy) = D(x)\sigma(y) + \tau(x)D(y), \text{ for all } x, y \in N.$$

□

**Lemma 2.** *Let  $D$  be a  $(\sigma, \tau)$ -derivation on a near-ring  $N$  and  $a \in N$ . Then for all  $x, y \in N$ ,*

$$(\tau(x)D(y) + D(x)\sigma(y))\sigma(a) = \tau(x)D(y)\sigma(a) + D(x)\sigma(y)\sigma(a).$$

*Proof.* For all  $x, y \in N$ , we get

$$\begin{aligned} D((xy)a) &= \tau(xy)D(a) + D(xy)\sigma(a) \\ &= \tau(x)\tau(y)D(a) + (\tau(x)D(y) + D(x)\sigma(y))\sigma(a). \end{aligned}$$

On the other hand,

$$\begin{aligned} D(x(ya)) &= \tau(x)D(ya) + D(x)\sigma(ya) \\ &= \tau(x)\tau(y)D(a) + \tau(x)D(y)\sigma(a) + D(x)\sigma(y)\sigma(a). \end{aligned}$$

For these two expressions of  $D(xya)$ , we obtain that, for all  $x, y \in N$ ,

$$(\tau(x)D(y) + D(x)\sigma(y))\sigma(a) = \tau(x)D(y)\sigma(a) + D(x)\sigma(y)\sigma(a).$$

□

**Lemma 3.** *Let  $N$  be a prime near-ring,  $D$  a nonzero  $(\sigma, \tau)$ -derivation of  $N$  and  $a \in N$ .*

- i) *If  $D(N)\sigma(a) = 0$  then  $a = 0$ .*
- ii) *If  $aD(N) = 0$  then  $a = 0$ .*

*Proof.* i) For all  $x, y \in N$ , we get

$$0 = D(xy)\sigma(a) = \tau(x)D(y)\sigma(a) + D(x)\sigma(y)\sigma(a).$$

Using hypothesis and  $\sigma$  is an automorphism of  $N$ , we have

$$D(x)N\sigma(a) = 0.$$

Since  $N$  is prime near-ring and  $D$  is a nonzero  $(\sigma, \tau)$ -derivation of  $N$ , we obtain  $a = 0$ .

- ii) A similar argument works if  $aD(N) = 0$ .

□

**Lemma 4.** *Let  $D$  be a  $(\sigma, \tau)$ -derivation which commute  $\sigma$  and  $\tau$ . If  $N$  is a 2-torsion free near-ring and  $D^2 = 0$  then  $D = 0$ .*

*Proof.* For arbitrary  $x, y \in N$ , we have

$$\begin{aligned} 0 &= D^2(xy) = D(D(xy)) = D(\tau(x)D(y) + D(x)\sigma(y)) \\ &= \tau^2(x)D^2(y) + D(\tau(x))\sigma(D(y)) + \tau(D(x))D(\sigma(y)) + D^2(x)\sigma^2(y). \end{aligned}$$

By hypothesis,

$$2D(\tau(x))D(\sigma(y)) = 0 \quad \text{for all } x, y \in N.$$

Since  $N$  is 2-torsion free near-ring and  $\sigma$  is an automorphism on  $N$ , we get

$$D(\tau(x))D(N) = 0.$$

It gives  $D = 0$  by Lemma 3 (ii).  $\square$

**Theorem 1.** *Let  $N$  be a near-ring and  $D$  a nonzero  $(\sigma, \tau)$ -derivation of  $N$ . If  $u \in N$  is not a left zero divisor and  $[D(u), u]_{\sigma, \tau} = 0$  then  $(x, u)$  is constant (that is,  $D(x, u) = 0$ ) for every  $x \in N$ .*

*Proof.* Since  $u(u + x) = u^2 + ux$ , we have  $D(u(u + x)) = D(u^2 + ux)$ . Expanding this equation, we have

$$\tau(u)D(u + x) + D(u)\sigma(u + x) = D(u^2) + D(ux)$$

and so

$$\begin{aligned} \tau(u)D(u) + \tau(u)D(x) + D(u)\sigma(u) + D(u)\sigma(x) \\ = \tau(u)D(u) + D(u)\sigma(u) + \tau(u)D(x) + D(u)\sigma(x) \end{aligned}$$

which reduces to

$$\tau(u)D(x) + D(u)\sigma(u) - \tau(u)D(x) - D(u)\sigma(u) = 0.$$

Therefore

$$\tau(u)D(x, u) = 0$$

by using the assumption  $[D(u), u]_{\sigma, \tau} = 0$ . Since  $u$  is not a left zero divisor, we get  $D(x, u) = 0$ . Thus  $(x, u)$  is a constant for every  $x \in N$ .  $\square$

**Theorem 2.** *Let  $N$  be a prime near-ring with a nonzero  $(\sigma, \tau)$ -derivation  $D$  such that  $\sigma D = D\sigma$  and  $\tau D = D\tau$ . If  $D(N) \subset Z$  then  $(N, +)$  is abelian. Moreover, if  $N$  is 2-torsion free, then  $N$  is a commutative ring.*

*Proof.* Suppose that  $a \in N$  such that  $D(a) \neq 0$ . So,  $D(a) \in Z \setminus \{0\}$  and  $D(a) + D(a) \in Z \setminus \{0\}$ . For all  $x, y \in N$ , we have

$$(x + y)(D(a) + D(a)) = (D(a) + D(a))(x + y),$$

that is,

$$xD(a) + xD(a) + yD(a) + yD(a) = D(a)x + D(a)y + D(a)x + D(a)y.$$

Since  $D(a) \in Z$ , we get

$$D(a)x + D(a)y = D(a)y + D(a)x,$$

and so,

$$D(a)(x, y) = 0 \text{ for all } x, y \in N.$$

Since  $D(a) \in Z \setminus \{0\}$  and  $N$  is a prime near-ring, it follows that  $(x, y) = 0$ , for all  $x, y \in N$ . Thus  $(N, +)$  is abelian.

Using hypothesis, for any  $b, c \in N$ ,

$$\sigma(c)D(ab) = D(ab)\sigma(c).$$

By Lemma 2, we have

$$\sigma(c)\tau(a)D(b) + \sigma(c)D(a)\sigma(b) = \tau(a)D(b)\sigma(c) + D(a)\sigma(b)\sigma(c).$$

Comparing these two expressions, using  $D(N) \subset Z$  and  $(N, +)$  is abelian, we obtain that

$$\sigma(c)\tau(a)D(b) + D(a)\sigma(c)\sigma(b) = \tau(a)D(b)\sigma(c) + D(a)\sigma(b)\sigma(c)$$

so we have

$$D(b)[\tau(a), \sigma(c)] = D(a)\sigma([c, b]) \text{ for all } b, c \in N.$$

Suppose now that  $N$  is not commutative. Choosing  $b, c \in N$  such that  $[b, c] \neq 0$  and  $a = D(x) \in Z$ , we get

$$D^2(x)\sigma([c, b]) = 0 \text{ for all } x \in N.$$

Since the central element  $D^2(x)$  can not be a nonzero divisor of zero, we conclude  $D^2(x) = 0$  for all  $x \in N$ . By Lemma 4, this cannot happen for nontrivial  $D$ .  $\square$

**Theorem 3.** *Let  $N$  be a prime near-ring admitting a nonzero  $(\sigma, \tau)$ -derivation  $D$  such that  $\sigma D = D\sigma$  and  $\tau D = D\tau$ . If  $[D(N), D(N)] = 0$ , then  $(N, +)$  is abelian. Moreover, if  $N$  is 2-torsion free, then  $N$  is a commutative ring.*

*Proof.* The argument used in the proof of Theorem 2 shows that if both  $z$  and  $z + z$  commute elementwise with  $D(N)$ , then we have

$$(2.1) \quad zD(x, y) = 0 \text{ for all } x, y \in N.$$

Substituting  $D(t), t \in N$  for  $z$  in (2.1), we get  $D(t)D(x, y) = 0$ . Since  $\sigma$  is an automorphism of  $N$ , we have  $\sigma(D(t))\sigma(D(x, y)) = 0$ . Using  $\sigma D = D\sigma$ , we get

$$D(\sigma(t))\sigma(D(x, y)) = 0 \text{ for all } x, y, t \in N.$$

By Lemma 3 (i), we obtain that  $D(x, y) = 0$  for all  $x, y \in N$ . For  $w \in N$ , we have  $0 = D(wx, wy) = D(w(x, y))$  and so we obtain

$$D(w)\sigma((x, y)) = 0.$$

Again, applying Lemma 3 (i), we get  $(x, y) = 0$  for all  $x, y \in N$ .

Now, assume that  $N$  is 2-torsion free. By the assumption  $[D(N), D(N)] = 0$ ,

$$D(\sigma(z))D(D(x)y) = D(D(x)y)D(\sigma(z)) \text{ for all } x, y, z \in N.$$

Hence, we get

$$\begin{aligned} D(\sigma(z))\tau(D(x))D(y) + D(\sigma(z))D^2(x)\sigma(y) \\ = \tau(D(x))D(y)D(\sigma(z)) + D^2(x)\sigma(y)D(\sigma(z)) \end{aligned}$$

by Lemma 2. Using  $D(\tau(x))D(\sigma(z)) = D(\sigma(z))D(\tau(x))$ ,  $\sigma D = D\sigma$  and  $\tau D = D\tau$ , we have

$$\begin{aligned} D(\tau(x))D(\sigma(z))D(y) + D(\sigma(z))D^2(x)\sigma(y) \\ = D(\tau(x))D(y)D(\sigma(z)) + D^2(x)\sigma(y)D(\sigma(z)) \end{aligned}$$

Since  $(N, +)$  is abelian, we conclude that

$$D(\tau(x))[D(\sigma(z)), D(y)] = D^2(x)\sigma([D(z), y]) \text{ for all } x, y, z \in N.$$

The left term of this equation is zero by the hypothesis, so we get

$$(2.2) \quad D^2(x)\sigma(D(z))\sigma(y) = D^2(x)\sigma(y)\sigma(D(z)) \text{ for all } x, y, z \in N.$$

Replacing  $y$  by  $yt$ , ( $t \in N$ ) in (2.2) and using (2.2), we have

$$\begin{aligned} D^2(x)\sigma(y)\sigma(t)\sigma(D(z)) &= D^2(x)\sigma(D(z))\sigma(y)\sigma(t) \\ &= D^2(x)\sigma(y)\sigma(D(z))\sigma(t) \end{aligned}$$

and so,

$$(2.3) \quad D^2(x)N\sigma([t, D(z)]) = 0 \text{ for all } x, t, z \in N.$$

Since  $N$  is a prime near-ring, we have

$$D^2(N) = 0 \text{ or } D(N) \subset Z$$

by Brauers's Trick. If  $D^2(N) = 0$ , then it contradicts that  $D$  is a nonzero  $(\sigma, \tau)$ -derivation of  $N$  by Lemma 4. So,  $D(N) \subset Z$ . Thus,  $N$  is a commutative ring by Theorem 2.  $\square$

**Theorem 4.** *Let  $N$  be a 2-torsion free prime near-ring,  $d_1$  a  $(\sigma, \tau)$ -derivation of  $N$  and  $d_2$  a derivation of  $N$ . If  $d_1d_2(N) = 0$ , then  $d_1 = 0$  or  $d_2 = 0$ .*

*Proof.* For  $x, y \in N$ , we have

$$\begin{aligned} 0 &= d_1d_2(xy) = d_1(xd_2(y) + d_2(x)y) \\ &= \tau(x)d_1d_2(y) + d_1(x)\sigma(d_2(y)) + \tau(d_2(x))d_1(y) + d_1d_2(x)\sigma(y). \end{aligned}$$

That is,

$$(2.4) \quad d_1(x)\sigma(d_2(y)) + \tau(d_2(x))d_1(y) = 0 \quad \text{for all } x, y \in N.$$

If we take  $d_2(x)$  instead of  $x$  in (2.4), then

$$\tau(d_2^2(x))d_1(y) = 0 \quad \text{for all } x, y \in N.$$

Using Lemma 3 (ii) one can obtain  $d_1 = 0$  or  $d_2^2 = 0$ . If  $d_2^2 = 0$ , we have  $d_2 = 0$  by Lemma 4. This completes the proof of theorem.  $\square$

**Theorem 5.** *Let  $N$  be a 2-torsion free prime near-ring,  $d_1$  a derivation and  $d_2$  be a  $(\sigma, \tau)$ -derivation of  $N$  such that  $\tau d_2 = d_2 \tau$  and  $\tau d_1 = d_1 \tau$ . If  $d_1 d_2(N) = 0$ , then  $d_1 = 0$  or  $d_2 = 0$ .*

*Proof.* The same argument in the proof of Theorem 4, we can write

$$(2.5) \quad d_1(\tau(x))d_2(y) + d_2(x)d_1(\sigma(y)) = 0 \quad \text{for all } x, y \in N.$$

Replacing  $x$  by  $d_2(x)$  in (2.5) and using  $\tau d_2 = d_2 \tau$  and  $\tau d_1 = d_1 \tau$ , we have

$$d_2^2(x)d_1(\sigma(y)) = 0 \quad \text{for all } x, y \in N.$$

Applying [1, Lemma 3 (ii)], we obtain  $d_1 = 0$  or  $d_2^2 = 0$ . If  $d_2^2 = 0$ , then  $d_2 = 0$  by Lemma 4.  $\square$

**Theorem 6.** *Let  $D$  be a nonzero  $(\sigma, \tau)$ -derivation of a prime near-ring  $N$  and  $a \in N$ . If  $[D(N), a]_{\sigma, \tau} = 0$  then  $D(a) = 0$  or  $a \in Z$ .*

*Proof.* By hypothesis,

$$D(ax)\sigma(a) = \tau(a)D(ax) \quad \text{for all } x \in N$$

and so,

$$(\tau(a)D(x) + D(a)\sigma(x))\sigma(a) = \tau(a)(\tau(a)D(x) + D(a)\sigma(x)).$$

Since  $N$  satisfies the partial distributive law by Lemma 2, we get

$$\tau(a)D(x)\sigma(a) + D(a)\sigma(x)\sigma(a) = \tau(a)\tau(a)D(x) + \tau(a)D(a)\sigma(x).$$

Using the hypothesis, we have

$$\tau(a)\tau(a)D(x) + D(a)\sigma(x)\sigma(a) = \tau(a)\tau(a)D(x) + D(a)\sigma(a)\sigma(x),$$

that is,

$$(2.6) \quad D(a)\sigma([x, a]) = 0 \quad \text{for all } x \in N.$$

Substituting  $xy$ , ( $y \in N$ ) for  $x$  and using (2.6), we have

$$D(a)\sigma(x)\sigma([y, a]) = 0 \quad \text{for all } x, y \in N.$$

Since  $\sigma$  is automorphism of prime near-ring of  $N$ , we get  $D(a) = 0$  or  $a \in Z$ . This completes the proof.  $\square$



**Theorem 7.** *Let  $D$  be a nonzero  $(\sigma, \tau)$ -derivation of a prime near-ring  $N$  such that  $\sigma D = D\sigma$  and  $\tau D = D\tau$ . If  $[D(N), D(N)]_{\sigma, \tau} = 0$ , then  $(N, +)$  is abelian. Moreover, if  $N$  is 2-torsion free then  $N$  is a commutative ring.*

*Proof.* By Theorem 6, we have

$$N = \{x \in N \mid D^2(x) = 0\} \cup \{x \in N \mid D(x) \in Z\}.$$

By Brauer's Trick, we get  $D^2(N) = 0$  or  $D(N) \subset Z$ . Since  $D$  is a nonzero  $(\sigma, \tau)$ -derivation of  $N$ , we get  $D(N) \subset Z$ . By Theorem 2, we prove the theorem.  $\square$

#### REFERENCES

- [1] BELL, H. E. AND MASON G., *On Derivations in near-rings*, Near-rings and Near-fields, North-Holland Mathematical Studies **137**, (1987).
- [2] KAMAL, AHMED A. M.,  *$\sigma$ -derivations on prime near-rings*, Tamkang J. Math. **32**(2001), no.2, 89–93.
- [3] PILZ, G., *Near-rings 2nd Ed.*, North Holland, Amsterdam, (1983).

ÖZNUR GÖLBAŞI

CUMHURİYET UNIVERSITY

FACULTY OF ARTS AND SCIENCE

DEPARTMENT OF MATHEMATICS

SIVAS - TURKEY

*e-mail address:* ogolbasi@cumhuriyet.edu.tr

*URL:* <http://www.cumhuriyet.edu.tr>

NEŞET AYDIN

ÇANAKKALE 18 MART UNIVERSITY

FACULTY OF ARTS AND SCIENCE

DEPARTMENT OF MATHEMATICS

ÇANAKKALE - TURKEY

*e-mail address:* neseta@comu.edu.tr

*URL:* <http://www.comu.edu.tr>

(Received June 4, 2003)