

Source: Okayama University (Japan), Center for Public Affairs and Communications

For immediate release: 28 Sept 2015

Okayama University research: Metamaterial absorbers for infrared inspection technologies

(Okayama, 28 September) A metamaterial absorber capable of highly sensitive infrared detection enhances the spectroscopy of tiny molecular details

Plasmonic metamaterials are man-made substances whose structure can be manipulated to influence the way they interact with light. As such, metamaterials offer an attractive platform for sensing applications, including infrared (IR) absorption spectroscopy – a technique used to uncover details of the chemical make-up and structure of substances.

Now, Atsushi Ishikawa at Okayama University and his colleague (or collaborator), Takuo Tanaka at RIKEN's Metamaterials Laboratory, Japan, (Note: I just feel, not necessary to emphasize "Tanaka at RIKEN".) have fabricated a novel plasmonic metamaterial absorber comprised of gold and magnesium fluorine capable of high sensitivity IR detection¹. The metamaterial could prove invaluable in the development of next-generation IR inspection technologies.

The researchers carefully designed their absorber to maximise the IR signal and minimise background noise. The metamaterial consists of 50 nm gold ribbons on a thick gold film, separated by a layer of magnesium fluorine (see image).

The wavelength of IR is longer and has less energy than visible light, meaning that it is not strong enough to excite electrons, unlike other types of spectroscopy. IR absorption spectroscopy therefore exploits the ability of IR to induce vibrations in bonded atoms. Organic compounds will absorb IR radiation corresponding to the different types of molecular vibrations present; the resulting absorption spectra tell scientists about the unique chemical structure of the compounds.

To test the capabilities of the new metamaterial, the team decided to identify the stretching vibrational modes of carbon-hydrogen bonds in 16-Mercaptohexadecanoic (16-MHDA) acid. They dipped the absorber in 16-MHDA ethanol solution to encourage a self-assembling monolayer of the acid molecules to develop. Under IR radiation at different incident angles, the metamaterial-monolayer spectral output displayed distinct peaks corresponding to carbon-hydrogen stretching, with the most pronounced peaks under IR at an angle of 40°.

The new metamaterial approach gave highly-detailed measurements pertaining to tiny molecular details (at the attamole level) in the 16-MHDA monolayer. The researchers hope their new technique will open doors to the development of ultrasensitive IR inspection technologies for material science and security applications.

Background

Metamaterials

The ability to manipulate the light absorption of materials could revolutionize many technologies, such as photovoltaic cells and thermal devices. Research into the design and development of plasmonic metamaterials is still relatively new. These materials are synthetic, and scientists can design their surface structures to exploit the behavior of surface plasmons – quasiparticles that exist on metal surfaces and interact with light – to achieve tuneable optical properties.

Infrared absorption spectroscopy could be dramatically enhanced by the introduction of tuneable metamaterial-based absorbers designed to enable high-resolution detection of tiny molecular details.

Methodology

The metamaterial absorber built by the team comprised gold nano-ribbons (measuring 50 nm thick) on a gold film base, with a thin layer of magnesium fluoride separating the two gold layers. As molecular monolayers self-assemble on noble metal surfaces, they hypothesized that the gold-based metamaterial would prove a strong candidate for enabling the high-resolution measurement of IR-induced vibrational modes in self-assembling monolayers.

Their approach involved covering their metamaterial absorber with an ultrathin self-assembling monolayer of 16-MDHA acid molecules. They also covered a bare gold film sample with the same monolayer for comparison.

The researchers subjected the two monolayers to IR radiation at different incident angles. The monolayer on bare gold exhibited a low signal-noise ratio, and it was very difficult to see the absorption dips on the spectral output corresponding to the IR-induced carbon-hydrogen stretching in the monolayer.

In contrast, the absorption dips were very well-pronounced in the spectral read-out for the metamaterial-monolayer, because the vibrational modes of the 16-MHDA molecules resonated with the plasmonic modes of the metamaterial. This so-called 'resonant coupling' produced distinct peaks corresponding to IR-induced carbon-hydrogen stretching in the 16-MHDA molecular structure. The resonant coupling was dependent on the angle of the incident light, with the clearest, strongest signal at an angle of 40°.

Future work

The researchers believe their absorber may open doors to new ultra-sensitive IR detection technologies. Further, their technique could be exploited in other ways – by optimizing the surface structure of other metamaterials, they could enhance resonant coupling still further and enable sensitivities down to the zeptomole level.




Figure caption

Researchers at Okayama University have created a new IR spectroscopic technique utilizing the properties of a metamaterial-based absorber to enhance spectral output. Trials on a self-assembling monolayer of 16-MHDA acid showed distinct peaks corresponding to carbon-hydrogen stretching in the monolayer.

Reference

Atsushi Ishikawa & Takuo Tanaka. Metamaterial absorbers for infrared detection of molecular self-assembled monolayers. *Nature Scientific Reports* **5** (12570) Published online, 31 July 2015.

doi: 10.1038/srep12570.

<http://www.nature.com/articles/srep12570>

Correspondence to

Assistant Professor Atsushi Ishikawa, Ph.D.
Department of Electrical and Electronic Engineering,
Okayama University, 3-1-1 Tsushimanaka, Kitaku,
Okayama, 700-8530, Japan.
E-mail: a-ishikawa@okayama-u.ac.jp



Further information

Okayama University

1-1-1 Tsushima-naka, Kita-ku, Okayama 700-8530, Japan

Public Relations and Information Strategy, Okayama University

E-mail: www-adm@adm.okayama-u.ac.jp

Website: http://www.okayama-u.ac.jp/index_e.html

Okayama Univ. e-Bulletin: <http://www.okayama-u.ac.jp/user/kouhou/ebulletin/>

Okayama Univ. e-Bulletin (PDF Issues): [http://www.okayama-](http://www.okayama-u.ac.jp/en/tp/cooperation/ebulletin.html)

[u.ac.jp/en/tp/cooperation/ebulletin.html](http://www.okayama-u.ac.jp/en/tp/cooperation/ebulletin.html)

About Okayama University (You Tube):

<https://www.youtube.com/watch?v=iDL1coqPRYI>

Okayama University Image Movie (You Tube):

<https://www.youtube.com/watch?v=WnbJVk2eIA>

<https://www.youtube.com/watch?v=KU3hOIXS5kk>

Okayama University Medical Research Updates (OU-MRU)

Vol.1 : [Innovative non-invasive 'liquid biopsy' method to capture circulating tumor cells from blood samples for genetic testing](#)

Vol.2 : [Ensuring a cool recovery from cardiac arrest](#)

Vol.3 : [Organ regeneration research leaps forward](#)

Vol.4 : [Cardiac mechanosensitive integrator](#)

Vol.5 : [Cell injections get to the heart of congenital defects](#)

Vol.6 : [Fourth key molecule identified in bone development](#)

Vol.7 : [Anticancer virus solution provides an alternative to surgery](#)

Vol.8 : [Light-responsive dye stimulates sight in genetically blind patients](#)

Vol.9 : [Diabetes drug helps towards immunity against cancer](#)

Vol.10 : [Enzyme-inhibitors treat drug-resistant epilepsy](#)

Vol.11 : [Compound-protein combination shows promise for arthritis treatment](#)

Vol.12 : [Molecular features of the circadian clock system in fruit flies](#)

Vol.13 : [Peptide directs artificial tissue growth](#)

Vol.14 : [Simplified boron compound may treat brain tumours](#)

About Okayama University

Okayama University is one of the largest comprehensive universities in Japan with roots going back to the Medical Training Place sponsored by the Lord of Okayama and established in 1870. Now with 1,300 faculty and 14,000 students, the University offers courses in specialties ranging from medicine and pharmacy to humanities and physical sciences. Okayama University is located in the heart of Japan approximately 3 hours west of Tokyo by Shinkansen.

