

Erratum

In the article by Senoo M *et al.* entitled “**Anxiety and Feelings toward the Baby among Pregnant Women with Uterine Leiomyomas**”, which appeared in the December 2015 issue, Vol. 69, No. 6, pp 339-348, following corrections should be listed.

1. page 341. Line 2 from the top, in the right column.

Error:

Among the 25-29-year-old women ($n = 38$), the proportion of pregnant women with anxiety about pregnancy tended to be higher in the group without uterine leiomyomas compared to the uterine leiomyoma group ($p = 0.095$).

Correction:

Among the 40-44-year-old women ($n = 5$), the proportion of pregnant women with anxiety about pregnancy in the group without uterine leiomyomas had a significantly higher rate of anxiety than the group with uterine leiomyomas.

From the perspective of primipara only, among the 25-29-year-old women ($n = 38$), the proportion of pregnant women with anxiety about pregnancy tended to be higher in the group without uterine leiomyomas compared to the uterine leiomyoma group ($p = 0.063$).

2. page 341. Just above Table 1, in the right column.

Error:

significant difference.

Correction:

significant difference ($p = 0.079$).

3. page 342. Table 2

Error:

Table 2 Perceived anxiety in pregnant women with and without uterine leiomyomas

	Overall Uterine leiomyomas		P-value	Primipara Uterine leiomyomas		P-value	Multipara Uterine leiomyomas		P-value
	Without (n = 109)	With (n = 23)		Without (n = 58)	With (n = 11)		Without (n = 51)	With (n = 12)	
Overall	35 (32.1%)	7 (30.4%)	n.s.	25 (43.1%)	3 (27.3%)	n.s.	10 (19.6%)	4 (33.3%)	n.s.
Age									
>24 years (n = 6)	33.3% (2/6)	0% (0/0)	n.s.	33.3% (2/6)	0% (0/0)	n.s.	0% (0/0)	0% (0/0)	n.s.
25-29 years (n = 38)	35.3% (12/34)	0% (0/4)	n.s. (0.095)	50.0% (11/22)	0% (0/4)	n.s.	8.3% (1/12)	0% (0/0)	n.s.
30-34 years (n = 45)	30.6% (11/36)	55.6% (5/9)	n.s.	40.0% (6/15)	50.0% (2/4)	n.s.	23.8% (5/21)	60% (3/5)	n.s.
35-39 years (n = 38)	28.1% (9/32)	33.3% (2/6)	n.s.	35.7% (5/14)	50% (1/2)	n.s.	22.2% (4/18)	25.0% (1/4)	n.s.
40-44 years (n = 5)	100% (1/1)	0% (0/4)	<0.05	100% (1/1)	0% (0/1)	n.s.	0 (0%)	0% (0/3)	n.s.
History of miscarriage and/or stillbirth									
Without (n = 106)	28.1% (25/89)	23.5% (4/17)	n.s.	42.2% (19/45)	30.0% (3/10)	n.s.	13.6% (6/44)	14.3% (1/7)	n.s.
With (n = 26)	50.0% (10/20)	50.0% (3/6)	n.s.	46.2% (6/13)	0% (0/1)	n.s.	57.1% (4/7)	60.0% (3/5)	n.s.
History of fertility treatment									
Without (n = 107)	29.7% (27/91)	25.0% (4/16)	n.s.	39.6% (19/48)	0% (0/5)	n.s.	18.6% (8/43)	36.4% (4/11)	n.s.
With (n = 25)	44.4% (8/18)	42.9% (3/7)	n.s.	60.0% (6/10)	50.0% (3/6)	n.s.	25.0% (2/8)	0% (0/1)	n.s.
Pregnancy term									
1st trimester (n = 10)	28.6% (2/7)	66.7% (2/3)	n.s.	50.0% (2/4)	0% (0/0)	n.s.	0% (0/3)	66.7% (2/3)	n.s. (0.083)
2nd trimester (n = 44)	38.9% (14/36)	25.0% (2/8)	n.s.	52.6% (10/19)	66.7% (2/3)	n.s.	23.5% (4/17)	0% (0/5)	n.s.
3rd trimester (n = 74)	27.4% (17/62)	25.0% (3/12)	n.s.	37.5% (12/32)	12.5% (1/8)	n.s.	16.7% (5/30)	50.0% (2/4)	n.s.
Threatened abortion and/or preterm labor									
Without (n = 121)	28.3% (28/99)	27.3% (6/22)	n.s.	38.0% (19/50)	20.0% (2/10)	n.s.	18.4% (9/49)	33.3% (4/12)	n.s.
With (n = 11)	70.0% (7/10)	100% (1/1)	n.s.	75.0% (6/8)	100% (1/1)	n.s.	50.0% (1/2)	0% (0/0)	n.s.

Data are mean ± SD, median [range], n.s.: not significant.

Correction:

Table 2 Perceived anxiety in pregnant women with and without uterine leiomyomas

	Overall Uterine leiomyomas		P-value	Primipara Uterine leiomyomas		P-value	Multipara Uterine leiomyomas		P-value
	Without (n = 109)	With (n = 23)		Without (n = 58)	With (n = 11)		Without (n = 51)	With (n = 12)	
Overall	35 (32.1%)	7 (30.4%)	n.s.	25 (43.1%)	3 (27.3%)	n.s.	10 (19.6%)	4 (33.3%)	n.s.
Age									
>24 years (n = 6)	33.3% (2/6)	0% (0/0)	—	33.3% (2/6)	0% (0/0)	—	0% (0/0)	0% (0/0)	—
25-29 years (n = 38)	35.3% (12/34)	0% (0/4)	n.s.	50.0% (11/22)	0% (0/4)	n.s. (0.063)	8.3% (1/12)	0% (0/0)	—
30-34 years (n = 45)	30.6% (11/36)	55.6% (5/9)	n.s.	40.0% (6/15)	50.0% (2/4)	n.s.	23.8% (5/21)	60% (3/5)	n.s.
35-39 years (n = 38)	28.1% (9/32)	33.3% (2/6)	n.s.	35.7% (5/14)	50% (1/2)	n.s.	22.2% (4/18)	25.0% (1/4)	n.s.
40-44 years (n = 5)	100% (1/1)	0% (0/4)	<0.05	100% (1/1)	0% (0/1)	n.s.	0% (0/0)	0% (0/3)	—
History of miscarriage and/or stillbirth									
Without (n = 106)	28.1% (25/89)	23.5% (4/17)	n.s.	42.2% (19/45)	30.0% (3/10)	n.s.	13.6% (6/44)	14.3% (1/7)	n.s.
With (n = 26)	50.0% (10/20)	50.0% (3/6)	n.s.	46.2% (6/13)	0% (0/1)	n.s.	57.1% (4/7)	60.0% (3/5)	n.s.
History of fertility treatment									
Without (n = 107)	29.7% (27/91)	25.0% (4/16)	n.s.	39.6% (19/48)	0% (0/5)	n.s. (0.079)	18.6% (8/43)	36.4% (4/11)	n.s.
With (n = 25)	44.4% (8/18)	42.9% (3/7)	n.s.	60.0% (6/10)	50.0% (3/6)	n.s.	25.0% (2/8)	0% (0/1)	n.s.
Pregnancy term									
1st trimester (n = 10)	28.6% (2/7)	66.7% (2/3)	n.s.	50.0% (2/4)	0% (0/0)	—	0% (0/3)	66.7% (2/3)	n.s. (0.083)
2nd trimester (n = 44)	38.9% (14/36)	25.0% (2/8)	n.s.	52.6% (10/19)	66.7% (2/3)	n.s.	23.5% (4/17)	0% (0/5)	n.s.
3rd trimester (n = 74)	27.4% (17/62)	25.0% (3/12)	n.s.	37.5% (12/32)	12.5% (1/8)	n.s.	16.7% (5/30)	50.0% (2/4)	n.s.
Threatened abortion and/or preterm labor									
Without (n = 121)	28.3% (28/99)	27.3% (6/22)	n.s.	38.0% (19/50)	20.0% (2/10)	n.s.	18.4% (9/49)	33.3% (4/12)	n.s.
With (n = 11)	70.0% (7/10)	100% (1/1)	n.s.	75.0% (6/8)	100% (1/1)	n.s.	50.0% (1/2)	0% (0/0)	—

Data are mean ± SD, median [range], n.s.: not significant.