



Original Full Length Article

## CCN4/WISP-1 positively regulates chondrogenesis by controlling TGF- $\beta$ 3 function



Yuya Yoshioka<sup>a</sup>, Mitsuaki Ono<sup>a,b,\*</sup>, Azusa Maeda<sup>a,c</sup>, Tina M. Kilts<sup>c</sup>, Emilio Satoshi Hara<sup>a,d</sup>, Hany Khattab<sup>a,e</sup>, Junji Ueda<sup>a</sup>, Eriko Aoyama<sup>e</sup>, Toshitaka Oohashi<sup>e</sup>, Masaharu Takigawa<sup>e</sup>, Marian F. Young<sup>c</sup>, Takuo Kuboki<sup>a</sup>

<sup>a</sup> Department of Oral Rehabilitation and Regenerative Medicine, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Okayama, Japan

<sup>b</sup> Department of Molecular Biology and Biochemistry, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Okayama, Japan

<sup>c</sup> Craniofacial and Skeletal Diseases Branch, National Institute of Dental and Craniofacial Research, National Institutes of Health, Bethesda, MD, USA

<sup>d</sup> Department of Biomaterials, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Okayama, Japan

<sup>e</sup> Advanced Research Center for Oral & Craniofacial Sciences, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Okayama, Japan

### ARTICLE INFO

#### Article history:

Received 4 August 2015

Revised 6 October 2015

Accepted 6 November 2015

Available online 10 November 2015

#### Keywords:

Bone marrow stromal cell (BMSC)

Chondrogenesis

CCN4/WISP-1

TGF- $\beta$ 3

Articular cartilage

### ABSTRACT

The CCN family of proteins plays important roles in development and homeostasis of bone and cartilage. To understand the role of CCN4 in chondrogenesis, human bone marrow stromal cells (hBMSCs) were transduced with CCN4 adenovirus (adCCN4) or siRNA to CCN4 (siCCN4) in the presence or absence of transforming growth factor- $\beta$ 3 (TGF- $\beta$ 3). Overexpression of CCN4 enhanced TGF- $\beta$ 3-induced SMAD2/3 phosphorylation and chondrogenesis of hBMSCs in an *in vitro* assay using a micromass culture model. On the other hand, knockdown of CCN4 inhibited the TGF- $\beta$ 3-induced SMAD2/3 phosphorylation and synthesis of cartilage matrix in micromass cultures of hBMSCs. Immunoprecipitation–western blot analysis revealed that CCN4 bound to TGF- $\beta$ 3 and regulated the ability of TGF- $\beta$ 3 to bind to hBMSCs. *In vivo* analysis confirmed there was a significant decrease in the gene expression levels of chondrocyte markers in cartilage samples from *Ccn4*-knock out (KO) mice, compared to those from wild type (WT) control. In order to investigate the regenerative properties of the articular cartilage in *Ccn4*-KO mice, articular cartilage defects were surgically performed in the knee joints of young mice, and the results showed that the cartilage was partially repaired in WT mice, but not in *Ccn4*-KO mice. In conclusion, these results show, for the first time, that CCN4 has a positive influence on chondrogenic differentiation by modulating the effects of TGF- $\beta$ 3.

© 2015 Elsevier Inc. All rights reserved.

### 1. Introduction

Osteoarthritis (OA), which is characterized by the degeneration and loss of articular cartilage and subchondral bone, is one of the most common disorders of the musculoskeletal system, and is caused mainly by the effects of aging, trauma or injury and overloading, such as in the case of obesity. The main symptoms of OA of the hip or knee are pain and stiffness, causing loss of joint mobility. Articular cartilage consists basically of a single cell type, the chondrocyte, which can synthesize all components of the extracellular matrix (ECM), and the mechanical function and properties of articular cartilage is greatly dependent on the quality and quantity of ECM. However, adult cartilage has a poor capacity for repair and regeneration. Therefore, autologous chondrocyte transplantation has been widely used. Stem cell therapy, which employs the use of stem/progenitor cells from different tissue or induced-

pluripotent stem cells (iPSCs), has gained particular attention in recent years. However, a deeper understanding of the detailed mechanisms of chondrogenic differentiation of stem/progenitor cells, including the interaction with growth factors, is necessary to eventually improve the current clinical outcomes for cartilage regeneration.

Transforming growth factor-beta (TGF- $\beta$ ) is a member of a family of dimeric polypeptide growth factors and is classified into four major sub-families, TGF- $\beta$ s, bone morphogenetic proteins (BMPs), activins/inhibins, and glia cell line-derived neurotrophic factors (GDNFs). Among the TGF- $\beta$  superfamily, TGF- $\beta$ 1,  $\beta$ 2 and  $\beta$ 3 are known to play important roles not only in chondrocyte proliferation and differentiation but also in embryonic development of cartilage through activation of the SMAD2/3 signaling pathway; and alterations in TGF- $\beta$  signaling are known to be involved in the pathogenesis of OA and osteoporosis [1,2]. In early stages of chondrogenesis from mesenchymal stem cell condensation to limb bud formation, TGF- $\beta$  activates SMAD3, which binds to co-SMAD4, translocates to the nucleus and promotes the expression of *Sox9*, the master regulator of chondrogenesis [3].

The CCN family consists of six members, CCN1/Cyr61/Cef10, CCN2/CTGF/Fisp12, CCN3/Nov, CCN4/WISP-1/rCOP-1/Elm-1, CCN5/WISP-2, and CCN6/WISP-3. CCN family proteins are characterized by four

\* Corresponding author at: Department of Oral Rehabilitation and Regenerative Medicine, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, 2-5-1 Shikata-cho, Kita-ku, Okayama-shi, Okayama-ken 700-8525, Japan.

E-mail address: [mitsuaki@md.okayama-u.ac.jp](mailto:mitsuaki@md.okayama-u.ac.jp) (M. Ono).

distinct functional domains: an insulin-like growth factor binding protein like module (IGFBP), a von Willebrand factor type C repeat (VWC), a thrombospondin type 1 repeat (TSP1), and a cysteine-rich C-terminal module (CT). Because CCN family proteins have this unique modular structure, they can interact with various other proteins, cell membrane receptors and ECM components, and can regulate several cell functions, including adhesion, migration, proliferation, and differentiation both *in vitro* and *in vivo*. Interestingly, CCN family members can interact with and regulate the activity of members of the TGF- $\beta$  superfamily and orchestrate tissue development during osteogenesis and chondrogenesis [4,5]. For instance, CCN2 was shown to promote the proliferation and differentiation of chondrocytes by interacting with BMP-2, FGF-2, and IGF-1 [4,6]. Additionally, our research group showed that CCN4 regulates osteoblast function by controlling the activity of TGF- $\beta$ 1 and BMP-2 [7,8].

Clinical and experimental evidence also suggest that CCN4 plays a role in OA [9,10]. Previous studies reported a high expression of CCN4 in the synovium and cartilage in an experimental OA model in mice as well as in human OA patients [10,11]. These findings suggest that CCN4 may be involved in the chondrocyte differentiation and/or cartilage regeneration. However, the specific function of CCN4 in chondrogenesis and cartilage regeneration remains unclear.

To understand the role of CCN4 in chondrogenesis, we investigated the effect of CCN4 on the chondrogenic differentiation of human bone marrow stromal cells (hBMSCs) using gain and loss of function models, and found that CCN4 is a positive regulator of TGF- $\beta$ 3-induced chondrogenic differentiation of hBMSCs *in vitro*. Moreover, in an experimental animal model of full-thickness articular cartilage defect using *Ccn4*-knock out (KO) mice, we showed that CCN4 is important for cartilage repair *in vivo*.

## 2. Materials and methods

### 2.1. Cells and culture medium

Human bone marrow stromal cells (hBMSCs) were purchased from Lonza (Walkersville, MD, USA). Cells were cultured in  $\alpha$ -MEM containing 10% fetal bovine serum (FBS), 1% L-glutamine (Life Technologies, Rockville, MD, USA), and 1% penicillin and streptomycin (Sigma, St. Louis, MO, USA) [Basal medium].

For chondrogenic differentiation, hBMSCs at passages 4 to 5 were used for analysis of chondrogenic differentiation. hBMSCs were suspended ( $2 \times 10^5$  cells/10  $\mu$ L) and cultured in micromasses, as previously reported [12,13]. Chondrogenic medium consisted of DMEM low glucose, supplemented with 6  $\mu$ g/mL each of insulin, transferrin and selenous acid (ITS solution, BD, Bedford, MA, USA), 50  $\mu$ g/mL of L-Ascorbic Acid Phosphate Magnesium Salt *n*-Hydrate (Wako Pure Chemical Industries, Osaka, Japan), 100 nM dexamethasone (Sigma), 10 ng/mL of rhTGF- $\beta$ 3 (R&D Systems, Inc., Minneapolis, MN, USA) and antibiotics. Medium was changed every 3–4 days and samples were cultured for 28 days prior to analysis.

For the overexpression of the *CCN4* gene, hBMSCs were transfected with adenovirus that express human CCN4 (adCCN4) as reported previously [8]. Adenovirus CMV empty vector (adCMV) was used as the negative control. For the knockdown of *Ccn4* gene, hBMSCs were cultured in penicillin and streptomycin-free basal medium, following the manufacturer's instruction. Twenty four hours later, the medium was changed to Opti-MEM Reduced-Serum Medium (Life Technologies) supplemented with Lipofectamine RNAiMAX (Life Technologies) and 10 nM siRNA to decrease the expression of human *CCN4* (siCCN4; Life Technologies) according to the manufacturer's instructions. Stelth<sup>TM</sup> RNAi Negative Control High GC Duplex (Life Technologies) was used as the negative control.

For the isolation of chondrocytes, the growth plates of femur, which were collected from newborn WT and *Ccn4*-KO, were incubated with Trypsin-EDTA (Life Technologies) for 30 min and 86.5 U/mL of Type 1

collagenase (Worthington Biochemical Corporation, Lakewood, NJ, USA) overnight at 37 °C. Isolated chondrocytes were cultured with DMEM/F12 (Life Technologies) containing 10% fetal bovine serum (FBS), 1% L-glutamine, and 1% penicillin and streptomycin.

### 2.2. Reverse transcription and real time RT-PCR analysis

Total cellular RNA was extracted by RNA Minikit (Life Technologies), according to the manufacturer's instructions. Possible residual DNA was removed by DNase (Purelink<sup>TM</sup> DNase; Life Technologies). RNA samples were reverse-transcribed by using iScript cDNA synthesis kit (Bio-Rad; Hercules, CA, USA). Real time RT-PCR was performed to quantify the expression of the target gene by using KAPA SYBR FAST qPCR Master Mix (KAPA BIOSYSTEMS, Wilmington, MA, USA) and Chromo4 real-time detector. The levels of mRNAs of interest were normalized to that of the reference gene *S29*. Primer sequences are shown in Table 1.

### 2.3. Western blot analysis

Cells were lysed by M-PER (Mammalian Protein Extraction Reagent; Thermo, Waltham, Massachusetts, USA) supplemented with a Protease Inhibitor Cocktail (1 mg/mL; Roche, Indianapolis, IN, USA) and Phosphatase Inhibitor Signal-Use Cocktail (1:100; Thermo). Cell debris were removed from lysates by centrifugation at 12,000 rpm for 10 min at 4 °C. Protein concentration in the cell lysate was determined by using Pierce<sup>®</sup> BCA Protein assay kit (Thermo). Ten micrograms of the total protein lysate was separated in precast polyacrylamide gels (NuPage, Life Technologies) by electrophoresis and then transferred onto polyvinylidene fluoride membranes (PVDF; GE Healthcare Life sciences, Buckinghamshire, UK) at 30 V for 2 h. Blots were blocked and incubated in primary antibodies to SOX-9 (1:1000; Millipore, Billerica, MA, USA), phosphorylated SMAD2/3 (pSMAD2/3), total SMAD2/3 (1:1000; Cell Signaling Technologies, Danvers, MA, USA) and  $\beta$ -ACTIN as a loading control (1:5000; Sigma). Membranes were then washed, and incubated in HRP-labeled anti-rabbit IgG (1:2000; Santa Cruz Biotechnology, Santa Cruz, CA, USA) for 1 h at room temperature (RT). The blots were developed using Forte western HRP Substrate (Millipore), and visualized with Image Quant LAS 4000 mini (Fujifilm, Tokyo, Japan).

### 2.4. Immunoprecipitation

Cells were harvested using NP-40 buffer (1% NP-40, 0.15 M NaCl, 0.01 M sodium phosphate, and 1 mM ethylenediaminetetraacetic acid (EDTA)) containing a Complete Protease Inhibitor Mixture (Roche Diagnostics, Mannheim, Germany). Cleared cell lysates were incubated with magnetic beads conjugated with 4C5 anti-DDK mouse monoclonal antibody (OriGene, Rockville, MD, USA), or 1  $\mu$ g mouse IgG1 kappa monoclonal [MOPC-21] isotype control (Abcam, Cambridge, UK) that was previously bound to Dynabeads protein G (Life Technologies). The beads were washed extensively, and the bound proteins were boiled in loading buffer for 10 min at 70 °C and then separated by SDS-PAGE and analyzed by immunoblot with 4C5 anti-DDK monoclonal antibody (OriGene) or anti-CCN4 antibody (LF-187). Details on the construction and specificity of LF-187 have been described previously [14,15].

### 2.5. Cell protein binding assay

To assess the effect of CCN4 on the cell binding affinity of rhTGF- $\beta$ 3, hBMSCs were seeded at a concentration of  $1 \times 10^4$  cells/well in 96-well plates. After 24 h, cells were transfected with adCCN4 or siCCN4 for 48 h. Next, cells were fixed for 15 min at RT with 4% paraformaldehyde (PFA), washed and pre-blocked in 5% goat serum (Invitrogen) and 0.05% bovine serum albumin (BSA) for 1 h at 37 °C. Cells were incubated with 30 ng/mL of TGF- $\beta$ 3 for 4 h at RT. Bound TGF- $\beta$ 3 ligands were detected by incubating the cells with anti-TGF- $\beta$ 3 antibody (1:100; ProteinTech Group, Inc., Chicago, IL, USA) overnight at 4 °C. Cells were washed and

**Table 1**  
Primers used for Real time RT-PCR analysis.

Gene	Species	GenBank accession no.	Primer sequence	PCR product length (bp)
S29	Human	BC032813	5'-T CTCGCT CTT GTCGT GT CT GTTC-3'(S) 5'-ACACTGGCGGCACATATTGAGG-3'(AS)	75
CCN1	Human	AF003114	5'-GCCAATGAAGCAGCGTTTCCC-3'(S) 5'-CAAT GAGT CCCAT CACCCACACC-3'(AS)	171
CCN2	Human	BT019794	5'-T GCGAGGAGT GGGTGTGTGAC-3'(S) 5'-TGGACCAGGCAGTTGGCTCTAATC-3'(AS)	125
CCN3	Human	BC015028	5'-GAGCAGTGCCAATCTACAGCGAAG-3'(S) 5'-AGATGGAGAAGCAGGAAGGTCAGG-3'(AS)	138
CCN4	Human	AB034725	5'-ACACTCATTAAAGCGAGGAAGAAG-3'(S) 5'-TCAGGACACTGGAAGGACACG-3'(AS)	185
CCN5	Human	BC058074	5'-CACACCGAAGACCCACCTCTG-3'(S) 5'-GCAGCAGCCACAGCCATCC-3'(AS)	151
CCN6	Human	AY358350	5'-GGGTACAGGGCACTGGACCATTAG-3'(S) 5'-ACAGCATCCACGCCATCTCTCAC-3'(AS)	167
COL2A1	Human	BT007205	5'-TGGAGCAGCAAGAGCAAGGAGAAG-3'(S) 5'-CCGTGGACAGCAGCGTAGG-3'(AS)	136
S29	Mouse	NM_009093	5'-GGAGTCACCCACGGAAGTTCGG-3'(S) 5'-GGAAGCACTGGCGGCACATG-3'(AS)	108
Sox-9	Mouse	NM_011448	5'-TCAGATGCACTGAGGAGCAC-3'(S) 5'-CCAGCCACAGCAGTGAGTAA-3'(AS)	208
Acan	Mouse	L07049	5'-CCCAGTACCCTACAGAGACA-3'(S) 5'-ACAGTGACCCTGGAAGTTCGG-3'(AS)	215
Col2a1	Mouse	BC082331	5'-GGGCAACAGCAGGTTACATACAC-3'(S) 5'-TCAATAATGGGAAGCGGGAGGTC-3'(AS)	120
Col10a1	Mouse	NM_009925	5'-TTCTGCTGTAATGTTCTTGA-3'(S) 5'-AATGCCTTGTCTCTCTTAC-3'(AS)	164
Bgn	Mouse	L20276	5'-ACCTGTCCTTCCATCTCT-3'(S) 5'-CCGTGTGTGTGTGTGTGT-3'(AS)	201
Dcn	Mouse	X53929	5'-CAAACATACTCGCATCCCT-3'(S) 5'-TGTCCAAGTGGAGTTCCTC-3'(AS)	214

S, sense; AS, antisense.

incubated with the HRP-labeled anti-rabbit IgG (1:1000; Santa Cruz Biotechnology) for 1 h at RT. The amount of bound TGF- $\beta$ 3 was determined by tetramethylbenzidine (TMB) colorimetric substrate (Cell Signaling) after incubation for 10 min at RT. The reaction was terminated with an equal volume of 2 N H<sub>2</sub>SO<sub>4</sub>, and the optical absorbance was measured at 450 nm.

#### 2.6. In vivo full-thickness articular cartilage defect model in *Ccn4* knockout (KO) mice

Mice deficient in *Ccn4* (*Ccn4*<sup>-/-</sup>) were generated using standard homologous recombination and gene targeting as described before [16]. Wild type (WT) and *Ccn4*-KO mice (C57BL/6) at 6 weeks of age were utilized in the experiments, according to the guidelines for Animal Care and Use at the NIDCR (ASP #13-700), NIH. Full-thickness articular cartilage defects were created according to a previously published report [17]. Briefly, three female WT and *Ccn4*-KO mice were anesthetized with isoflurane (2–5% in O<sub>2</sub>) and the joint capsule of the knee was opened, and the patella was dislocated laterally to expose the distal articular surface of the femur. Then, full-thickness articular cartilage defects of 400  $\mu$ m diameter were created using 27 G needles, which has a diameter of 400  $\mu$ m, in order to make the same size defect. Penetration to subchondral bone was confirmed by bleeding from the cartilage defect site. Muscles and patella were replaced on their original position and muscles and skin were sutured in separate layers. Four weeks after the surgery, animals were euthanized for histological analysis.

#### 2.7. Isolation of articular cartilage from WT and *Ccn4*-KO mice

To compare the gene expression level of chondrocyte markers in the articular cartilage of WT and *Ccn4*-KO mice at 6 weeks of age, articular cartilage samples were carefully collected from the caput femur of WT

and *Ccn4*-KO mice under a surgical microscope. Total RNA was isolated from the tissue samples using Trizol (Life Technologies), and gene expression levels of *Sox9*, aggrecan (*Acan*) and collagen type II, alpha 1 (*Col2a1*) were analyzed by real time RT-PCR.

#### 2.8. Histological analysis and qualitative scoring

hBMSC pellets and mice samples were fixed in 4% PFA, dehydrated and embedded in paraffin. Serial sections of 5  $\mu$ m were made, hydrated, and subsequently stained with hematoxylin and eosin (H&E), toluidine blue or safranin O for staining of glycosaminoglycans, as described previously [12]. For immunohistochemical staining of collagen type II, paraffin sections were deparaffinized, rehydrated, antigen-retrieved, blocked with 10% normal goat serum (Life Technologies) for 10 min, and incubated overnight at 4 °C with anti-collagen type II antibody (1:200; Millipore). Sections were then washed, incubated with secondary antibody and detected with 3-Amino-9-ethylcarbazole (AEC) color developing reagent (SuperPicture™ HRP Polymer Conjugate Broad Spectrum, Life Technologies), according to the manufacturer's instructions. In a blinded fashion, an examiner used a modified ICRS score to evaluate the histological condition of the articular cartilage surface post-surgery, as reported previously [12]. In pellet culture experiments, quantitation of the positive area of collagen type II was performed using a BZ-X700 microscope equipped with a BZ analyzer (Keyence, Osaka, Japan).

#### 2.9. Statistical analysis

The results obtained from quantitative experiments were reported as the mean values  $\pm$  SD. Statistical differences were examined using a one-way factorial ANOVA followed by Tukey's multiple comparison tests or Student's unpaired *t*-tests.

### 3. Results

#### 3.1. Expression of CCN family members in hBMSC cultures during chondrogenic differentiation in vitro

To obtain a better insight into the molecular mechanisms underlying the role of CCN family members in the process of chondrogenic differentiation of hBMSCs, we first investigated the expression levels of mRNA encoding genes of CCN family members in 21-day hBMSC micromass cultures. Although the transcript levels of *CCN5* and *CCN6* fluctuated during the culture period, there was a relatively high increase in the gene expression levels of *CCN1*, *CCN2*, *CCN3* and *CCN4* after the day 7, which followed a similar expression pattern of *COL2A1*, suggesting that these proteins could be playing important roles in modulating chondrogenesis (Fig. 1).


#### 3.2. CCN4 regulates TGF- $\beta$ -induced chondrogenesis of hBMSCs

Since *CCN1*, *CCN3* and mainly *CCN2* had already been reported to be important factors regulating chondrocyte differentiation [3,18,19], we evaluated the influence of *CCN4* on the chondrogenic process of hBMSCs by inducing *CCN4* over-expression using adenovirus encoding *CCN4* (adCCN4) as well as by silencing *CCN4* gene using siCCN4. AdCCN4 was transduced into hBMSCs [8], and the increase in *CCN4* gene expression level was confirmed by real time RT-PCR, in comparison with adCMV-transduced cells (Fig. 2A). Conversely, *CCN4* gene expression level was decreased with a targeted gene knock down using siCCN4 (Fig. 2E). First, we confirmed that overexpression and down regulation of *CCN4* did not affect hBMSC proliferation by MTS assay (data not shown). Next, since *CCN4* is considered to be a downstream target of


TGF- $\beta$  [7], we tested the effect of *CCN4* over expression and down regulation on TGF- $\beta$ -induced chondrogenic differentiation of hBMSCs in a micromass culture system. After 24 h of culture, we observed that adCCN4-transduced cells presented a notable increase in SOX9 protein levels (Fig. 2B). In accordance, after 28 days of culture, we could observe an intense staining with safranin O and Alcian blue for detection of glycosaminoglycans, as well as an intense immunostaining for collagen type II in adCCN4-transduced hBMSC cultures (Figs. 2C, S1). A quantitative analysis of the major chondrocyte marker genes, *ACAN* and *COL2A1*, confirmed a marked increase in the levels of the two genes in adCCN4-transduced hBMSC cultures, compared to adCMV (Fig. 2D). By contrast, knockdown of *CCN4* suppressed chondrogenic differentiation of hBMSCs, as demonstrated by a decrease in SOX9 levels in 24 h hBMSC pellets (Fig. 2F). Additionally, chondrocyte-like cells were nearly absent, and *ACAN* and *COL2A1* transcript levels were also significantly down regulated in siCCN4-transduced hBMSC micromasses, after 28 days of culture (Figs. 2G, H, S1). Taken together, these results demonstrate that *CCN4* plays an important role in enhancing chondrogenic differentiation of hBMSCs in vitro.

#### 3.3. CCN4 enhances TGF- $\beta$ -induced phosphorylation of SMADs

Next, in order to investigate the mechanism of how *CCN4* interacts with TGF- $\beta$  during chondrogenic differentiation of hBMSCs, we cultured the cells in monolayer, and transduced the cells with adCCN4. Forty eight hours after adCCN4 transduction, stimulation with rhTGF- $\beta$ 3 (10 ng/mL) for 10 min significantly increased the phosphorylation of SMAD2 and SMAD3 as compared to adCMV or TGF- $\beta$ 3-treated control cells (Fig. 3A). In contrast, as a response to the *CCN4* knockdown, TGF- $\beta$ 3-induced activation of SMAD2 and SMAD3 was repressed in hBMSCs



**Fig. 1.** Expression pattern of CCN family genes during chondrogenesis. hBMSCs were cultured in a micromass culture system in chondrogenic induction medium for up to 21 days. Total RNA was collected at 1, 3, 7, 14, and 21 days, and the mRNA expression levels of *CCN1* to *CCN6* (A–F) and *COL2A1* (G) were evaluated by real time RT-PCR. The expression of each gene was normalized to that of *S29* ribosomal RNA. Data are reported as mean  $\pm$  SD ( $n = 3$ ). \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$  versus day 1 (one-way ANOVA/Tukey). Results are representative data of at least three independent experiments.



**Fig. 2.** The effect of CCN4 on the chondrogenic differentiation of hBMSCs *in vitro*. hBMSCs were transduced with adCCN4 (A–D) or siCCN4 (E–H) for two days, and cultured in micromasses. (A, E) Two days after the transduction, mRNA expression levels of *CCN4* gene were measured by real-time RT-PCR. (B, F) The cellular proteins were collected 24 h after chondrogenic induction, and protein levels of SOX9 were detected by western blot.  $\beta$ -ACTIN was used as protein loading control. (C, G) Twenty eight days after chondrogenic induction, histological analysis of hBMSC micromass cultures was performed for detection of glycosaminoglycans with safranin O staining, or collagen type II by immunohistochemistry. Graphs show the quantitation of the positive area of collagen type II. (D, H) mRNA expression level of *ACAN* *COL2* gene was measured by real time RT-PCR. The expression of each gene was normalized to that of S29 ribosomal RNA. Bars represent the mean values and standard deviation ( $\pm$ SD) ( $n = 3$ ). \*\* $p < 0.01$ , \*\*\* $p < 0.001$  (Student's *t*-tests). Results are representative data of at least three independent experiments.

transduced with siCCN4 (Fig. 3B). In order to confirm the effect of CCN4 on the TGF- $\beta$  signaling pathway, we isolated chondrocytes from growth plate of newborn WT and *Ccn4*-KO mice, and observed that TGF- $\beta$ -induced phosphorylation of SMAD2/3 was also reduced in chondrocytes derived from *Ccn4*-KO mice (Fig. 3C). These results explain in part the effect of CCN4 in enhancing TGF- $\beta$ -induced chondrogenic differentiation of hBMSCs by promoting a stronger activation of SMAD2 and SMAD3.

### 3.4. Interaction between CCN4 and TGF- $\beta$ 3

It has been reported that CCN family members bind to the members of the TGF- $\beta$  superfamily and regulate their function, thus we speculated that CCN4 could bind to TGF- $\beta$ 3. To test this hypothesis, hBMSCs were transduced with adCCN4 and/or TGF- $\beta$ 3 overexpression vector for 2 days and the protein levels of CCN4 or TGF- $\beta$ 3 were detected by western blot (Fig. 4A and B). Immunoprecipitation (IP)-western blot analysis showed a binding between DDK-tagged TGF- $\beta$ 3 and CCN4 (Fig. 4C). The direct binding between the two proteins was also evaluated using recombinant proteins (rhCCN4 and rhTGF- $\beta$ 3); however, the IP-western results did not confirm that there was a direct binding between the two recombinant proteins (data not shown). Thus in cell extracts our data shows that CCN4 binds to TGF- $\beta$ 3, however we cannot conclude definitively whether this binding is direct or, rather, requires additional intermediating factors.


Since both CCN4 and TGF- $\beta$ 3 are secreted proteins, we then hypothesized that they could interact to each other at the cell membrane or

pericellularly at the extracellular level. Therefore, to determine whether CCN4 could regulate TGF- $\beta$ 3-to-cell binding affinity, we performed an on-cell protein binding assay with CCN4 overexpression and knock-down conditions. First, we performed a quantitative analysis of the amount of CCN4 bound to the surface of cells using anti-CCN4 antibody. As shown in Fig. 4D and E, the amount of CCN4 on the surface of cells was increased in the adCCN4-transduced hBMSCs; and on the other hand, it was decreased in siCCN4-transduced hBMSCs.

Next, to perform the TGF- $\beta$ 3 affinity assay, adCCN4 or siCCN4-transduced hBMSCs were treated with rhTGF- $\beta$ 3 (1  $\mu$ g/mL) for 4 h and the amount of TGF- $\beta$ 3 bound to the cell membrane was detected by using anti-TGF- $\beta$ 3 antibody. As predicted, increased amounts of TGF- $\beta$ 3 were detected in hBMSCs transduced with adCCN4 (Fig. 4F). Conversely, the binding affinity of TGF- $\beta$ 3 to hBMSCs was decreased under CCN4 knockdown condition (Fig. 4G). Taken together, these data support the concept that CCN4 can enhance TGF- $\beta$ 3 function by facilitating its binding to the surface of hBMSCs and subsequently enhancing the activation of TGF- $\beta$  downstream signaling.

### 3.5. *Ccn4* deficiency decreases the chondrogenic differentiation related markers in articular cartilage and delays articular cartilage regeneration *in vivo*

Next, we isolated articular cartilage samples from the caput femur of WT and *Ccn4*-KO mice (Fig. 5A–C), and found that the gene expression levels of chondrocyte-related markers, *Sox9*, *Acan* and *Col2a1*, cartilage



**Fig. 3.** The effect of CCN4 on TGF- $\beta$ 3-SMAD signaling pathway *in vitro*. hBMSCs were transduced with adCCN4 (A) or siCCN4 (B) for 2 days. (A, B) Transduced-hBMSCs in monolayer culture were treated with 10 ng/mL of TGF- $\beta$ 3 for 10 min, and cellular proteins were collected for detection of SMAD2/3 activity.  $\beta$ -ACTIN was used as protein loading control. (C) Chondrocytes were collected from mandibular condylar cartilage of newborn WT and *Ccn4*-KO mice, and stimulated with TGF- $\beta$ 3 for 10 min. The cellular proteins were collected for analysis of TGF- $\beta$ 3-induced phosphorylation of SMAD2/3. SMAD2/3 activity was decreased in chondrocytes derived from *Ccn4*-KO mice, compared to WT mice.


ECM including Biglycan (*Bgn*) and Decorin (*Dcn*), were significantly decreased in *Ccn4*-KO mice (Figs. 5D, S2), indicating that CCN4 is necessary for the maintenance of articular cartilage homeostasis. Of note, however, gene expression levels of collagen type X, alpha 1 (*Col10a1*), which is hypertrophic chondrocyte marker and highly expressed in articular cartilage of OA, were not detected in articular cartilage of either WT or *Ccn4*-KO mice (data not shown). Moreover, histological analysis of the joints showed that there was no notable difference in morphology between WT and *Ccn4*-KO mice in the articular cartilage of the knees (Fig. S3A–C).

Finally, we performed full-thickness articular cartilage defects in the knee joints of WT and *Ccn4*-KO mice, and investigated whether CCN4 could work as a positive regulator of articular cartilage repair. One month after the surgical defect, histological analysis of the joints revealed the formation of a newly-formed cartilage-like tissue positively stained with toluidine blue in WT mice (Fig. 6A). Moreover, immunostaining analysis showed that the newly-formed articular cartilage-like tissue was positive for collagen type II (Fig. 6B). Conversely, no cartilage-like tissue could be observed in the defect region of *Ccn4*-KO (Fig. 6A), and a histopathology-based scoring of the repair cartilage showed a significant difference between the WT mice group and *Ccn4*-KO mice group, indicating a substantial decrease in the ability of the articular cartilage to repair in *Ccn4*-KO mice (Fig. 6C).

#### 4. Discussion

CCN family members are expressed in development of growth plate cartilage and, because of their unique structural domain, they can interact with various growth factors and ECM components [20]. CCN2 for example, is expressed dominantly in the pre-hypertrophic zone of growth plate and interacts with BMP-2 and ACAN, and regulates chondrocyte proliferation and differentiation [4,21]. Furthermore, CCN2 was shown to promote the regeneration of articular cartilage [22]; and aging-related cartilage degradation could be prevented by cartilage-specific overexpression of CCN2 [23]. Using the targeted mutation mice model of CCN3, Roddy K.A. et al. have also reported that CCN3 is required for maintenance of articular cartilage and for joint homeostasis [24].


CCN4 is also expressed in the hypertrophic zone [25], and our previous results show that CCN4 can regulate osteogenesis by controlling the affinity of BMP-2 to hBMSCs [8]. Despite the implications of our previous reports that elucidated the effect of CCN4 on osteoblast differentiation and function [8,26], no study has yet determined a decisive role of CCN4 in chondrogenesis and cartilage regeneration. In this study, we demonstrated that CCN4 overexpression significantly increased the TGF- $\beta$ 3-induced chondrogenic differentiation of hBMSCs, and that CCN4 plays important roles in joint homeostasis and in the process of articular cartilage repair. Nevertheless, the articular cartilage defect




**Fig. 4.** Interaction between CCN4 and TGF- $\beta$ 3. hBMSCs were transduced with adCCN4 and/or TGF- $\beta$ 3-DDK-over expression vector, and incubated for 2 days, before collection of cell lysates. (A, B) Western blotting was performed using anti-CCN4 antibody or anti-DDK antibody, respectively. (C) Immunoprecipitation assay was performed using an anti-DDK antibody, and CCN4 or TGF- $\beta$ 3/DDK was detected by immunoblotting using an anti-CCN4 or anti-DDK antibodies. (D, E) hBMSCs were transduced with adCCN4 or siCCN4, respectively. After two days, cells were fixed and, the amount of CCN4 bound to cells was measured by its immunoreactivity to anti-CCN4 antibody. (F, G) To analyze the interaction of CCN4 and TGF- $\beta$ 3, fixed cells were incubated with TGF- $\beta$ 3 for 2 h. The amount of TGF- $\beta$ 3 bound to cells was measured by its immunoreactivity using anti-DDK antibody. Bars represent the mean values and standard deviation ( $\pm$ SD) ( $n = 3$ ). \*\* $p < 0.01$ , \*\*\* $p < 0.001$  (Student's  $t$ -tests). Results are representative data of at least three independent experiments.

model used in this study did not enable a complete repair of the articular surface in WT mice (Fig. 6). A possible reasoning for these results could be the age of the mice, since Matsuoka et al. demonstrated that

age is one of the crucial factors for the articular cartilage repair, with superior results observed in young (3 week-old) or juvenile (4 week-old) mice [17]; and thus, the 6 week-old mice used in this study could be



**Fig. 5.** Comparison of gene expression level of chondrogenic markers in articular cartilage between WT and *Ccn4*-KO mice. Articular cartilage samples of *caput femoris* for analysis of gene expression level of chondrogenic markers in healthy articular cartilage. Photograph and histological image of safranin O staining before the isolation of the articular cartilage are shown in (A) and (B), respectively; and the histological image after isolation is shown in (C). (D) Total RNA was collected from the articular cartilage separated from the *caput femoris*, and the mRNA expression levels of chondrogenic markers were measured by real time RT-PCR. The expression of each gene was normalized to that of S29 ribosomal RNA. Bars represent the mean values and standard deviation ( $\pm$ SD) ( $n = 3$ ). \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$  (Student's  $t$ -tests). Results are representative data of at least three independent experiments.



**Fig. 6.** Comparison of the articular cartilage healing between WT and KO mice. The full-thickness articular cartilage defects of 400  $\mu$ m in diameter were surgically created in the knee joints of WT and KO mice to evaluate the healing ability of articular cartilage. Four weeks after the surgery, samples were collected and specimens were stained with toluidine blue (A), and immunostained for collagen type II (B). (C) Evaluation of the histological condition of the articular cartilage surface was performed using a modified ICRS score. Bars represent the mean values and standard deviation ( $\pm$ SD) ( $n = 3$ ). \* $p < 0.05$  (Student's  $t$ -tests).

relatively aged, with a limited capacity for natural healing of the articular cartilage. Despite the incomplete repair of the articular cartilage in WT mice, our results show that CCN4 may be playing important roles in cartilage metabolism, as we could detect a significant decrease in the gene expression levels of *Sox9*, *Acan*, and *Col2a1* in the articular cartilage of *Ccn4*-KO mice, which eventually, could also contribute to the observed impaired healing capacity of chondrocytes.

Additionally, our recent article showed that the body length of *Ccn4*-KO mice is slightly but significantly reduced compared to wild type mice [16]. We also analyzed the growth plates of growing mice at embryonic day E18.5 and showed that the hypertrophic zone of growth plate was expanded in the *Ccn4*-KO compared to wild type mice. From these data, we also suspect that CCN4 could regulate the long bone growth by controlling the apoptosis of hypertrophic chondrocytes. Further experiments will be required to elucidate this issue.

Of note, *in vitro* experiments showed that CCN4 alone does not have any function for osteo-chondrogenesis. Therefore, CCN4 action may be mediated by interacting with other growth factors associated with chondrogenesis, and regulating their function. Here we show that CCN4 can interact with and control TGF- $\beta$ 3 function; however, additional experiments are necessary to clarify the exact mechanisms of action, since the direct binding between the two factors could not be determined. Other cellular mechanisms, independent of TGF- $\beta$ 3, could also be involved in the CCN4's role in chondrogenesis. For instance, it has been reported that tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ), a pro-inflammatory cytokine, induces focal loss of cartilage in OA knee joints [27]. Additionally, in the cardiac fibroblast, TNF- $\alpha$  up-regulates the expression level of CCN4 via its cyclic AMP response element-binding protein (CREB) site, and that CCN4 protects from TNF- $\alpha$  induced cell death [28]. Indeed, elevated levels of both TNF- $\alpha$  and CCN4 have been found in the cartilage of OA patients [9,10,29,30] and, CCN4 could be not only a key component in the process of cartilage differentiation but also could be playing important roles in organizing the inflammatory condition in OA, and promoting cartilage repair.

In summary, for the first time, we showed that CCN4 exerts a positive effect on TGF- $\beta$ 3-induced chondrogenesis by modulating TGF- $\beta$ 3 binding to the surface of hBMSCs and enhancing its downstream signaling. Further investigations are necessary for a better understanding of the underlying molecular and cellular mechanisms involved in effect of CCN family proteins in the process of chondrogenesis and/or cartilage repair, as well as for advancing new therapeutic modalities for OA to increase current clinical outcomes.

### Conflict of interest

The authors declare no competing interests.

### Author contributions

Study design: MO, MFY and TK. Study conduct: MO and MFY. Data collection: YY, AM, TMK, ESH and EA. Data analysis: YY, AM, HK, and JU. Data interpretation: MO, TO, MT, and TK. Drafting manuscript: YY, MO and ESH. Revising manuscript content: MFY, and TK. Approving final version of manuscript: TO, MT and TK.

### Acknowledgments

This study was supported in part by JSPS KAKENHI Grant Numbers #23890122 and #26861639 and the Intramural Program of the NIH, NIDCR.

### Appendix A. Supplementary data

Supplementary data to this article can be found online at <http://dx.doi.org/10.1016/j.bone.2015.11.007>.

### References

- [1] Y. Yamada, H. Okuizumi, A. Miyauchi, Y. Takagi, K. Ikeda, A. Harada, Association of transforming growth factor beta1 genotype with spinal osteophytosis in Japanese women, *Arthritis Rheum.* 43 (2000) 452–460.
- [2] H.H. Lau, A.Y. Ho, K.D. Luk, A.W. Kung, Transforming growth factor-beta1 gene polymorphisms and bone turnover, bone mineral density and fracture risk in southern Chinese women, *Calcif. Tissue Int.* 74 (2004) 516–521.
- [3] C.I. Lorda-Diez, J.A. Montero, M.J. Diaz-Mendoza, J.A. Garcia-Porrero, J.M. Hurler, Defining the earliest transcriptional steps of chondrogenic progenitor specification during the formation of the digits in the embryonic limb, *PLoS One* 6 (2011), e24546.
- [4] A. Maeda, T. Nishida, E. Aoyama, S. Kubota, K.M. Lyons, T. Kuboki, M. Takigawa, CCN family 2/connective tissue growth factor modulates BMP signalling as a signal conductor, which action regulates the proliferation and differentiation of chondrocytes, *J. Biochem.* 145 (2009) 207–216.
- [5] J.G. Abreu, N.I. Ketpura, B. Reversade, E.M. De Robertis, Connective-tissue growth factor (CTGF) modulates cell signalling by BMP and TGF-beta, *Nat. Cell Biol.* 4 (2002) 599–604.
- [6] E. Aoyama, S. Kubota, M. Takigawa, CCN2/CTGF binds to fibroblast growth factor receptor 2 and modulates its signaling, *FEBS Lett.* 586 (2012) 4270–4275.
- [7] C.A. Inkson, M. Ono, S.A. Kuznetsov, L.W. Fisher, P.G. Robey, M.F. Young, TGF-beta1 and WISP-1/CCN-4 can regulate each other's activity to cooperatively control osteoblast function, *J. Cell. Biochem.* 104 (2008) 1865–1878.

- [8] M. Ono, C.A. Inkson, T.M. Kilts, M.F. Young, WISP-1/CCN4 regulates osteogenesis by enhancing BMP-2 activity, *J. Bone Miner. Res.* 26 (2011) 193–208.
- [9] T. Urano, K. Narusawa, M. Shiraki, T. Usui, N. Sasaki, T. Hosoi, Y. Ouchi, T. Nakamura, S. Inoue, Association of a single nucleotide polymorphism in the WISP1 gene with spinal osteoarthritis in postmenopausal Japanese women, *J. Bone Miner. Metab.* 25 (2007) 253–258.
- [10] A.B. Blom, S.M. Brockbank, P.L. van Lent, H.M. van Beuningen, J. Geurts, N. Takahashi, P.M. van der Kraan, F.A. van de Loo, B.W. Schreurs, K. Clements, P. Newham, W.B. van den Berg, Involvement of the Wnt signaling pathway in experimental and human osteoarthritis: prominent role of Wnt-induced signaling protein 1, *Arthritis Rheum.* 60 (2009) 501–512.
- [11] J.F. Liu, S.M. Hou, C.H. Tsai, C.Y. Huang, C.J. Hsu, C.H. Tang, CCN4 induces vascular cell adhesion molecule-1 expression in human synovial fibroblasts and promotes monocyte adhesion, *Biochim. Biophys. Acta* 2013 (1833) 966–975.
- [12] E.S. Hara, M. Ono, P.T. Hai, W. Sonoyama, S. Kubota, M. Takigawa, T. Matsumoto, M.F. Young, B.R. Olsen, T. Kuboki, Fluocinolone acetonide is a potent synergistic factor of TGF-beta3-associated chondrogenesis of bone marrow-derived mesenchymal stem cells for articular surface regeneration, *J. Bone Miner. Res.* (2015).
- [13] A. Woods, G. Wang, H. Dupuis, Z. Shao, F. Beier, Rac1 signaling stimulates N-cadherin expression, mesenchymal condensation, and chondrogenesis, *J Biol Chem* 282 (2007) 23500–23508.
- [14] L. Xu, R.B. Corcoran, J.W. Welsh, D. Pennica, A.J. Levine, WISP-1 is a Wnt-1- and beta-catenin-responsive oncogene, *Genes Dev.* 14 (2000) 585–595.
- [15] M. Ono, C.A. Inkson, R. Sonn, T.M. Kilts, L.F. de Castro, A. Maeda, L.W. Fisher, P.G. Robey, A.D. Berendsen, L. Li, N. McCartney-Francis, A.C. Brown, N.P. Crawford, A. Molinolo, A. Jain, N.S. Fedarko, M.F. Young, WISP1/CCN4: a potential target for inhibiting prostate cancer growth and spread to bone, *PLoS One* 8 (2013), e71709.
- [16] A. Maeda, M. Ono, K. Holmbeck, L. Li, T.M. Kilts, V. Kram, M.L. Noonan, Y. Yoshioka, E.M. McNerny, M.A. Tantiillo, D.H. Kohn, K.M. Lyons, P.G. Robey, M.F. Young, WNT1-induced Secreted Protein-1 (WISP1), a novel regulator of bone turnover and Wnt signaling, *J Biol Chem* 290 (2015) 14004–14018.
- [17] M. Matsuoka, T. Onodera, F. Sasazawa, D. Momma, R. Baba, K. Hontani, N. Iwasaki, An articular cartilage repair model in common C57Bl/6 mice, *Tissue Eng. Part C Methods* (2015).
- [18] J. Lafont, C. Jacques, G. Le Dreau, F. Calhabeu, H. Thibout, C. Dubois, F. Berenbaum, M. Laurent, C. Martinier, New target genes for NOV/CCN3 in chondrocytes: TGF-beta2 and type X collagen, *J. Bone Miner. Res.* 20 (2005) 2213–2223.
- [19] S. Ivkovic, B.S. Yoon, S.N. Popoff, F.F. Safadi, D.E. Libuda, R.C. Stephenson, A. Daluiski, K.M. Lyons, Connective tissue growth factor coordinates chondrogenesis and angiogenesis during skeletal development, *Development* 130 (2003) 2779–2791.
- [20] H. Kawaki, S. Kubota, A. Suzuki, N. Lazar, T. Yamada, T. Matsumura, T. Ohgawara, T. Maeda, B. Perbal, K.M. Lyons, M. Takigawa, Cooperative regulation of chondrocyte differentiation by CCN2 and CCN3 shown by a comprehensive analysis of the CCN family proteins in cartilage, *J. Bone Miner. Res.* 23 (2008) 1751–1764.
- [21] E. Aoyama, T. Hattori, M. Hoshijima, D. Araki, T. Nishida, S. Kubota, M. Takigawa, N-terminal domains of CCN family 2/connective tissue growth factor bind to aggrecan, *Biochem. J.* 420 (2009) 413–420.
- [22] T. Nishida, S. Kubota, S. Kojima, T. Kuboki, K. Nakao, T. Kushibiki, Y. Tabata, M. Takigawa, Regeneration of defects in articular cartilage in rat knee joints by CCN2 (connective tissue growth factor), *J. Bone Miner. Res.* 19 (2004) 1308–1319.
- [23] S. Itoh, T. Hattori, N. Tomita, E. Aoyama, Y. Yutani, T. Yamashiro, M. Takigawa, CCN family member 2/connective tissue growth factor (CCN2/CTGF) has anti-aging effects that protect articular cartilage from age-related degenerative changes, *PLoS One* 8 (2013), e71156.
- [24] K.A. Roddy, C.A. Boulter, Targeted mutation of NOV/CCN3 in mice disrupts joint homeostasis and causes osteoarthritis-like disease, *Osteoarthr. Cartil.* 23 (2015) 607–615.
- [25] T. Yanagita, S. Kubota, H. Kawaki, K. Kawata, S. Kondo, T. Takano-Yamamoto, S. Tanaka, M. Takigawa, Expression and physiological role of CCN4/Wnt-induced secreted protein 1 mRNA splicing variants in chondrocytes, *FEBS J* 274 (2007) 1655–1665.
- [26] A. Maeda, M. Ono, K. Holmbeck, L. Li, T.M. Kilts, V. Kram, M.L. Noonan, Y. Yoshioka, E. McNerny, M. Tantiillo, D. Kohn, K.M. Lyons, P.G. Robey, M.F. Young, WNT1 induced secreted protein-1 (WISP1): a novel regulator of bone turnover and Wnt signaling, *J Biol Chem* (2015).
- [27] C.I. Westcott, A.F. Barakat, L. Wood, M.J. Perry, P. Neison, I. Bisbinas, L. Armstrong, A.B. Millar, C.J. Elson, Tumor necrosis factor alpha can contribute to focal loss of cartilage in osteoarthritis, *Osteoarthr. Cartil.* 8 (2000) 213–221.
- [28] K. Venkatchalam, B. Venkatesan, A.J. Valente, P.C. Melby, S. Nandish, J.E. Reusch, R.A. Clark, B. Chandrasekar, WISP1, a pro-mitogenic, pro-survival factor, mediates tumor necrosis factor-alpha (TNF-alpha)-stimulated cardiac fibroblast proliferation but inhibits TNF-alpha-induced cardiomyocyte death, *J Biol Chem* 284 (2009) 14414–14427.
- [29] O. Stannus, G. Jones, F. Cicuttini, V. Parameswaran, S. Quinn, J. Burgess, C. Ding, Circulating levels of IL-6 and TNF-alpha are associated with knee radiographic osteoarthritis and knee cartilage loss in older adults, *Osteoarthr. Cartil.* 18 (2010) 1441–1447.
- [30] J. Xue, J. Wang, Q. Liu, A. Luo, Tumor necrosis factor-alpha induces ADAMTS-4 expression in human osteoarthritis chondrocytes, *Mol. Med. Rep.* 8 (2013) 1755–1760.