

指 導 教 授 氏 名	指 導 役 割
高柴 正悟 印	全般的な指導
久保田 聡 印	実験、研究方針、論文作成指導
印	

## 学 位 論 文 要 旨

岡山大学大学院医歯薬学総合研究科

専攻分野 歯周病態学分野	身分 大学院生	氏名 堀 綾花
<p>論 文 題 名 Regulatory mechanism of CCN2 production by serotonin (5-HT) via 5-HT<sub>2A</sub> and 5-HT<sub>2B</sub> receptors in chondrocytes.  (軟骨細胞における 5-HT<sub>2A</sub> および 5-HT<sub>2B</sub> 受容体を介したセロトニン (5-HT) による CCN2 産生の調節機構)</p>		
<p>論文内容の要旨 (2000字程度)  Serotonin (5-hydroxytryptamine: 5-HT) is recognized as a neurotransmitter in the central nerve system and as a regulator of systemic blood pressure in the peripheral tissues. Recently, it was reported that 5-HT<sub>2</sub> receptors (5-HT<sub>2Rs</sub>) were expressed in cartilage tissues lacking both vessels and neurons, suggesting possible novel functions of 5-HT during cartilage development and regeneration. Our previous data indicated that CCN family protein 2/connective tissue growth factor (CCN2/CTGF) plays a central role in cartilage development and regeneration. Therefore, the aim of this study was to investigate the effect of 5-HT on the production of CCN2 in chondrocytes. Firstly, we showed that the mRNAs of 5-HT<sub>2R</sub> subtypes 5-HT<sub>2AR</sub> and 5-HT<sub>2BR</sub>, were expressed in a human chondrocytic cell line, HCS-2/8; however, 5-HT<sub>2CR</sub> mRNA was not detected. In addition, exogenously added 5-HT did not affect the 5-HT<sub>2AR</sub> and 5-HT<sub>2BR</sub> expressions. Next, we demonstrated that CCN2 production was increased by treatment with a 5-HT<sub>2AR</sub> agonist and the combination of 5-HT and 5-HT<sub>2BR</sub> antagonist. In contrast, treatment with a 5-HT<sub>2BR</sub> agonist and the combination of 5-HT and 5-HT<sub>2AR</sub> antagonist decreased CCN2 production. Furthermore, we showed that phosphorylation of Akt and p38 MAPK were increased by treatment with 5-HT<sub>2AR</sub> agonist, and that phosphorylation of PKCε, PKCζ, ERK1/2 and JNK were increased by treatment with 5-HT<sub>2BR</sub> agonist. Finally, we found that 5-HT<sub>2AR</sub> was localized in the growth plate, whereas 5-HT<sub>2BR</sub> was localized in the articular cartilage.</p>		

論文内容の要旨（2000 字程度）

These findings suggest that 5-HT promotes CCN2 production through the 5-HT<sub>2A</sub>R in growth plates, and that it represses CCN2 production through the 5-HT<sub>2B</sub>R in articular cartilage for harmonized development of long bones.