



Journal Name

COMMUNICATION

Catalytic Enantioselective Hosomi–Sakurai Reaction of α -Ketoesters Promoted by Chiral Copper(II) Complexes

Received 00th January 20xx,
Accepted 00th January 20xx

Yutaro Niwa, Mayu Miyake, Ichiro Hayakawa, Akira Sakakura*

DOI: 10.1039/x0xx00000x

www.rsc.org/

A catalytic enantioselective Hosomi–Sakurai reaction of α -ketoesters has been developed. A copper(II) complex with a chiral bis(oxazoline) ligand bearing methanesulfonamide groups shows excellent catalytic activity to give α,α -disubstituted α -hydroxyesters in high yields with high enantioselectivities. This is the first successful method for the catalytic enantioselective 1,2-addition of α -ketoesters with allylic silanes.

Catalytic asymmetric allylations of ketones are useful methods for the synthesis of chiral tertiary alcohols.¹ Several effective methods that use chiral catalysts have been developed. Among these methods, the Hosomi–Sakurai reaction^{2,3} is promising, since it employs inexpensive and nontoxic allylic silanes as carbon nucleophiles. However, little is known about the catalytic enantioselective Hosomi–Sakurai reaction of ketones,^{4,5} because ketones generally show lower reactivity and selectivity than aldehydes. In addition, the use of a chiral Lewis acid catalyst sometimes initiates undesired background reactions through a non-enantioselective silylium ion catalysis to decrease the enantioselectivity.⁶

α -Ketoesters are useful substrates for asymmetric 1,2-addition with a carbon nucleophile, since the reaction affords optically active α,α -disubstituted α -hydroxyesters, which are important structural motifs in biologically active compounds. Accordingly, methods for the catalytic asymmetric allylation of α -ketoesters have been reported.⁷ For example, Feng and colleagues reported asymmetric addition with tetraallylstannane catalyzed by *N,N'*-dioxide–In(III) complexes.^{7a} Yoda and colleagues developed an asymmetric addition with β -amido-functionalized allylstannanes catalyzed by chiral In(III) complexes.^{7b} Aminophenol-catalyzed asymmetric addition with allylboronates was also reported by Hoveyda and colleagues.^{7c} However, to the best of our knowledge, an efficient method for the asymmetric addition of α -ketoesters

with allylic silanes has not yet been reported.

Previously, Ishihara and Sakakura's group developed a new chiral bis(oxazoline) ligand **1a** bearing methanesulfonamide groups (Table 1).⁸ A copper(II) complex of **1a**, which is called *n*-cation catalyst, catalytically promotes the hetero-Diels–Alder reaction of β,γ -unsaturated α -ketoesters with allylsilanes to give the corresponding adducts in high yields with high diastereo- and enantioselectivities.^{8c} Since **1a**-CuX₂ successfully activates β,γ -unsaturated α -ketoesters, this catalyst may also be suitable for promoting the asymmetric 1,2-addition of α -ketoesters. We report herein the first catalytic enantioselective Hosomi–Sakurai reaction of α -ketoesters.



Our study commenced with an examination of the catalytic activities of **1a**-Cu(NTf₂)₂ in the Hosomi–Sakurai reaction of α -ketoester **2a** (Table 1). The reaction of **2a** with an allyltrimethylsilane was conducted in the presence of **1a**-Cu(NTf₂)₂ (5 mol%) at ambient temperature. Since the crude product included a small amount of TMS ether of **3a**,[†] the crude product was treated with TBAF to remove the TMS group. As a result, the reaction proceeded smoothly in EtNO₂ at ambient temperature to give **3a** in 78% yield with 74% ee (entry 1). In sharp contrast, the reaction in other solvents such as dichloromethane, acetonitrile, THF and isopropyl alcohol did not proceed at all even under heating conditions.[‡] The use of allyltriisopropylsilane gave **3a** with a slightly better enantioselectivity (80% ee), but the yield of **3a** was significantly decreased (50%) (entry 2). We next evaluated other chiral bis(oxazoline) ligands.⁹ Ligand **1b** bearing trifluoromethanesulfonamide groups is also effective for the Diels–Alder reaction of *N*-acryloyloxazolidinones.^{8a} However, the use of **1b** in the present reaction significantly decreased the enantioselectivity (53% ee), although the reactivity was improved (90% yield) (entry 3). When the reaction was conducted in the presence of ligand **1c**, which has sterically bulky *tert*-butyl groups at the 4,4'-positions, only a trace amount of **3a** was obtained (entry 4). In contrast, the use of **1d** bearing benzyl groups at the 4,4'-positions gave **3a** in 83% yield, while the enantioselectivity was poor (–7% ee)[§] (entry 5).

Graduate School of Natural Science and Technology, Okayama University, 3-1-1, Tsushima-naka, Kita-ku, Okayama, Japan.

Electronic Supplementary Information (ESI) available: Experimental procedures and characterization data for all new compounds (PDF). See DOI: 10.1039/x0xx00000x

The use of ligand **1e** or **1f**, which have four phenyl groups at the 4,4',5,5'-positions, gave **3a** in moderate yields (68 and 67%) in low enantioselectivity (−16 and −35% ee)⁸ (entries 6 and 7).

Table 1 Catalytic activities of chiral copper(II) complexes in the addition reaction of **2a** with allylsilanes^a



Entry	1	R	Yield (%) ^b	Ee (%) ^c
1	1a	Me	78	74
2	1a	<i>i</i> -Pr	50	80
3	1b	Me	90	53
4	1c	Me	3	ND
5	1d	Me	83	−7
6	1e	Me	68	−16
7	1f	Me	67	−35


^a The reaction of **2a** (0.2 mmol) was conducted with allylsilane (3 equiv) in the presence of **1**-Cu(NTf₂)₂ (5 mol%) in EtNO₂ at ambient temperature for 20 h. The crude product was treated with TBAF (1 equiv) in THF at ambient temperature for 0.5 h. ^b Isolated yield. ^c Evaluated by chiral HPLC analysis.

The optimized reaction conditions could be applied to the allylation of various α -ketoesters **2** (Table 2). The reaction of benzyl pyruvate **2b** gave the corresponding adduct **3b** in 71% yield with 65% ee (entry 1). The absolute configuration of **3b** was assigned to be *R* from the sign of the measured optical rotation, compared to that of the reported *S*-isomer.^{7a} This enantioselectivity could be explained by the proposed transition state assembly (Scheme 1). Intramolecular interaction of the sulfonamide groups of **1a** with copper(II) cation would preferentially shield the *Si* face of the coordinated **2b**.⁸ Allyltrimethylsilane would approach the *Re* face of **2b** to give (*R*)-**3b** as the major enantiomer.

Table 2 **1a**-Cu(NTf₂)₂-catalyzed Hosomi–Sakurai reaction of **2** with allyltrimethylsilane^a

Entry	2	R ¹	R ²	Yield (%) ^b	Ee (%) ^c
1	2b	Me	Bn	3b , 71	65 (<i>R</i>)
2	2c	CH ₃ (CH ₂) ₅	Et	3c , 74	74
3	2d	CH ₂ =CH(CH ₂) ₂	Et	3d , 41	74
4	2e	BnO(CH ₂) ₃	Et	3e , 75	73
5	2f	<i>c</i> -C ₅ H ₉	Et	3f , 67	69
6	2g	<i>c</i> -C ₆ H ₁₁	Et	3g , 0	—
7	2h	Ph	Me	3h , 0	—
8	2i	4-CF ₃ C ₆ H ₄	Et	3j , 72	79

^a The reaction of **2** (0.2 mmol) was conducted with allyltrimethylsilane (3 equiv) in the presence of **1**-Cu(NTf₂)₂ (5 mol%) in EtNO₂ at ambient temperature for 1–24 h. The crude product was treated with TBAF (1 equiv) in THF at ambient temperature. ^b Isolated yield. ^c Evaluated by chiral HPLC analysis.




Scheme 1 Proposed transition state assembly for the **1a**-Cu(NTf₂)₂-catalyzed allylation of **2b**

The reaction of primary alkyl-substituted α -ketoesters such as **2c** [R¹ = CH₃(CH₂)₅], **2d** [R¹ = CH₂=CH(CH₂)₂] and **2e** [R¹ = BnO(CH₂)₂] also gave the corresponding adducts **3c–e** in moderate to good yields (41–75%) with good enantioselectivities (73–74% ee) (entries 2–4).

The reactivity of the present allylation highly depended on the size of the R¹ group. For example, the reaction of cyclopentyl-substituted α -ketoester **2f** gave the corresponding adduct **3f** in 67% yield with 69% ee (entry 5), while cyclohexyl-substituted α -ketoester **2g** (R¹ = *c*-C₆H₁₁) and methyl benzoylformate (**2h**, R¹ = Ph) were inert under the present reaction conditions, and the starting materials were completely recovered (entries 6 and 7). The introduction of an electron-withdrawing CF₃ group on the phenyl group of **2h** successfully improved the reactivity to give **3i** in 72% yield with 79% ee (entry 8).

We next examined the catalytic enantioselective Hosomi–Sakurai reaction with methallyltrimethylsilane, which is more nucleophilic than allyltrimethylsilane¹⁰ (Scheme 2). As a result, the reaction proceeded rapidly even at −30 °C to give the corresponding adduct **4a** with high enantioselectivity (97% ee).

However, the yield of **4a** was low (48%) and a significant amount (32%) of **5a** was also formed as a byproduct. This byproduct should be generated via deprotonation of the tertiary cation intermediate, while desilylation of the same cationic intermediate gives the desired product **4a**. According to previous reports,¹¹ **5a** could be converted into **4a** under acidic conditions via formation of the tertiary cation intermediate. Indeed, treatment of **5a** with 5 mol% of **1a**-Cu(NTf₂)₂ at ambient temperature gave **4a** in 75% yield without any loss of enantiomeric excess (97% ee) (Scheme 2).



Scheme 2 **1a**-Cu(NTf₂)₂-catalyzed addition reaction of **2a** with methallyltrimethylsilane


The above results implied that when the reaction mixture, which was obtained from the reaction of **2a** with methallyltrimethylsilane at $-30\text{ }^{\circ}\text{C}$, was allowed to warm to ambient temperature, then the corresponding adducts **4a** would be formed as a single product. As we expected, the reaction of **2a** with methallyltrimethylsilane under these conditions ($-30\text{ }^{\circ}\text{C}$, 1 h; then rt, 2 h) gave **4a** in 90% yield with 96% ee (Table 3, entry 1).

Since adduct **4a** was obtained in high yield with high enantioselectivity, we next examined the substrate scope and limitations. As in the reaction with allyltrimethylsilane, the crude products included a small amount of TMS ether of **4**, so the crude products were treated with TBAF to remove the TMS group. The reaction of **2b–f** gave the corresponding adducts **4b–f** in good yields (53–79%) with high enantioselectivity (88–98% ee) (entries 2–6). However, the reaction of **2g**, which has a rather bulky cyclohexyl group, showed a slightly low enantioselectivity (78% ee), although the yield of **4g** was good (81%) (entry 7). The aryl-substituted α -ketoesters **2h–l** were also good substrates for the reaction with methallyltrimethylsilane, and gave the corresponding adducts **4h–l** in high yields (78–99%) with high enantioselectivity (93–98% ee) (entries 8–12). The introduction of a methyl group at the ortho-position of the phenyl group significantly decreased the reactivity, probably due to steric hindrance (14% yield, entry 13). In contrast to the 2-methylphenyl substituted **2m**, 2-fluoro-derivative **2n** showed good reactivity (75% yield) and high enantioselectivity (97% ee) (entry 14).

Table 3 Enantioselective addition reaction of **2** with methallyltrimethylsilane^a

Entry	2	R ¹	R ²	Yield (%) ^b	Ee (%) ^c
1	2a	Ph(CH ₂) ₂	Et	4a , 90	96
2	2b	Me	Bn	4b , 73	94
3	2c	CH ₃ (CH ₂) ₅	Et	4c , 79	94
4 ^d	2d	CH ₂ =CH(CH ₂) ₂	Et	4d , 75	96
5	2e	BnO(CH ₂) ₃	Et	4e , 53	98
6	2f	<i>c</i> -C ₅ H ₉	Et	4f , 53	88
7	2g	<i>c</i> -C ₆ H ₁₁	Et	4g , 81	78
8	2h	Ph	Me	4h , 99	98
9	2i	4-CF ₃ C ₆ H ₄	Et	4i , 81	97
10	2j	4-MeOC ₆ H ₄	Et	4j , 95	93
11	2k	4-MeC ₆ H ₄	Et	4k , 80	96
12	2l	3-MeC ₆ H ₄	Et	4l , 78	95
13 ^e	2m	2-MeC ₆ H ₄	Et	4m , 14	ND
14	2n	2-FC ₆ H ₄	Et	4n , 75	97

^a The reaction of **2** (0.2 mmol) with methallyltrimethylsilane (3 equiv) was conducted in EtNO₂ (1 mL) in the presence of **1a**-Cu(NTf₂)₂ at $-30\text{ }^{\circ}\text{C}$ for 1 h, and then the reaction mixture was warmed to ambient temperature. The crude product was treated with TBAF (0.5–3 equiv) in THF at $0\text{ }^{\circ}\text{C}$. ^b Isolated yield. ^c Evaluated by chiral HPLC analysis. ^d The reaction in the first step was conducted at $-50\text{ }^{\circ}\text{C}$. ^e The reaction in the first step was conducted at $60\text{ }^{\circ}\text{C}$.



Scheme 3 Ring-closing metathesis of adducts **3d** and **4d**.

α,α -Disubstituted α -hydroxyesters **3** and **4**, the adducts of the present Hosomi–Sakurai reaction, are synthetically useful chiral building blocks. For example, ring-closing metathesis of **3d** and **4d** using Grubbs' 2nd-generation catalyst (2 mol%) gave the corresponding cyclohexenes **6** and **7** in respective yields of 82 and 83% without loss of enantiomeric excess (Scheme 2). Compound **6** is a key chiral intermediate for the synthesis of quinic acid.¹²

In conclusion, we developed a catalytic enantioselective Hosomi–Sakurai reaction of α -ketoesters **2**. Chiral copper(II) complex with bis(oxazoline) ligand **1a** showed high catalytic activity, and chiral α,α -disubstituted α -hydroxyesters **3** and **4** were obtained in high yields with high enantioselectivities. Financial support for this project was partially provided by a Grant-in-Aid from JSPS KAKENHI (18K05123), the Sumitomo

Foundation and the Okayama Foundation for Science and Technology. The authors gratefully thank the Division of Instrumental Analysis, Department of Instrumental Analysis & Cryogenics, Advanced Science Research Center, Okayama University for the NMR and HRMS measurements.

Conflicts of interest

There are no conflicts to declare.

Notes and references

‡ See the Electronic Supplementary Information for details.

§ The minus sign indicates that the opposite enantiomer was obtained as a major product.

- For selected reviews, see: (a) S. E. Denmark, J. Fu, *Chem. Rev.*, 2003, **103**, 2763; (b) H. Lachance, D. G. Hall, *Org. React.* Vol. 73, Wiley, 2008; (c) M. Yus, J. C. González-Gómez, F. Foubelo, *Chem. Rev.*, 2011, **111**, 7774.
- (a) A. Hosomi, M. Endo, H. Sakurai, *Chem. Lett.*, 1976, **5**, 941; (b) A. Hosomi, H. Sakurai, *Tetrahedron Lett.*, 1976, **17**, 1295; (c) A. Hosomi, H. Sakurai, *J. Am. Chem. Soc.*, 1977, **99**, 1673.
- For reviews, see: (a) I. Fleming, J. Dungès, R. Smithers, *Org. React.*, 1989, **37**, 57; (b) I. Fleming, In *Comprehensive Organic Synthesis*, B. M. Trost, I. Fleming, Eds.; Pergamon: Oxford, 1991; Vol. 2, pp. 563.
- (a) S. Yamasaki, K. Fujii, R. Wada, M. Kanai, M. Shibasaki, *J. Am. Chem. Soc.*, 2002, **124**, 6536; (b) M. Wadamoto, H. Yamamoto, *J. Am. Chem. Soc.*, 2005, **127**, 14556.
- In contrast to the reaction of ketones, many successful methods for the catalytic enantioselective Hosomi–Sakurai reaction of aldehydes have been reported. For selected examples, see: (a) K. Furuta, M. Mouri, H. Yamamoto, *Synlett*, 1991, 561; (b) K. Ishihara, M. Mouri, Q. Gao, T. Maruyama, K. Furuta, H. Yamamoto, *J. Am. Chem. Soc.*, 1993, **115**, 11490; (c) D. R. Gauthier, Jr., E. M. Carreira, *Angew. Chem. Int. Ed.*, 1996, **35**, 2363; (d) S. A. A. El Bialy, H. Braun, L. F. Tietze, *Eur. J. Org. Chem.*, 2005, 2965. (e) D. A. Evans, Y. Aye, J. Wu, *Org. Lett.*, 2006, **8**, 2071; (f) Y. N. Belokon, D. Chusov, D. A. Borkin, L. V. Yashkina, P. Bolotov, T. Skrupskaya, M. North, *Tetrahedron: Asymmetry*, 2008, **19**, 459; (g) M. Mahlau, P. García-García, B. List, *Chem. Eur. J.*, 2012, **18**, 16283.
- (a) E. M. Carreira, R. A. Singer, *Tetrahedron Lett.*, 1994, **35**, 4323; (b) T. K. Hollis, B. Bosnich, *J. Am. Chem. Soc.*, 1995, **117**, 4570.
- (a) K. Zheng, B. Qin, X. Liu, X. Feng, *J. Org. Chem.*, 2007, **72**, 8478; (b) M. Takahashi, Y. Murata, M. Ishida, F. Yagishita, M. Sakamoto, T. Sengoku, H. Yoda, *Org. Biomol. Chem.*, 2014, **12**, 7686; (c) D. W. Robbins, K. Lee, D. L. Silverio, A. Volkov, S. Torker, A. H. Hoveyda, *Angew. Chem. Int. Ed.*, 2016, **55**, 9610.
- (a) A. Sakakura, R. Kondo, Y. Matsumura, M. Akakura, K. Ishihara, *J. Am. Chem. Soc.*, 2009, **131**, 17762; (b) A. Sakakura, K. Ishihara, *Chem. Soc. Rev.*, 2011, **40**, 163; (c) Y. Matsumura, T. Suzuki, A. Sakakura, K. Ishihara, *Angew. Chem. Int. Ed.*, 2014, **53**, 6131.
- For selected reviews of chiral bis(oxazoline) ligands, see: (a) A. K. Ghosh, P. Mathivanan, J. Cappiello, *Tetrahedron: Asymmetry*, 1988, **9**, 1; (b) J. S. Johnson, D. A. Evans, *Acc. Chem. Res.*, 2000, **33**, 325; (c) G. Desimoni, G. Faita, K. A. Jørgensen, *Chem. Rev.*, 2011, **111**, PR284.
- H. Mayr, M. Patz, *Angew. Chem. Int. Ed.*, 1994, **33**, 938.
- (a) I. Fleming, D. A. Perry, *Tetrahedron*, 1981, **37**, 4027; (b) B. Psaume, M. Montury, J. Goré, *Synth. Commun.*, 1982, **12**, 40; (c) R. Hunter, C. D. Simon, *Tetrahedron Lett.*, 1988, **29**, 2257;
- (d) I. Fleming, J. Dunoguès, R. Smithers, *Org. React.*, 1989, **37**, 57; (e) D. Serramedan, B. Delmond, G. Deleris, J. Dunogues, M. Pereyre, C. Filliatre, *J. Organomet. Chem.*, 1990, **398**, 79; (f) C.-M. Yu, S.-K. Yoon, S.-J. Lee, J.-Y. Lee, S. S. Kim, *Chem. Commun.*, 1998, 2749.