

Anti-tumor effect of neratinib against lung cancer cells harboring *HER2* oncogene alterations

YUSUKE OGOSHI¹, KAZUHIKO SHIEN¹, TAKAHIRO YOSHIOKA¹, HIDEJIRO TORIGOE¹, HIROKI SATO¹, MASAKIYO SAKAGUCHI², SHUTA TOMIDA³, KEI NAMBA¹, EISUKE KURIHARA¹, YUTA TAKAHASHI¹, KEN SUZAWA¹, HIROMASA YAMAMOTO¹, JUNICHI SOH¹ and SHINICHI TOYOOKA¹

Departments of ¹Thoracic, Breast and Endocrinological Surgery, ²Cell Biology and ³Biobank, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Okayama 700-8558, Japan

Received June 27, 2018; Accepted December 31, 2018

DOI: 10.3892/ol.2019.9908

Abstract. Human epidermal growth factor receptor 2 (HER2) is a member of the ErbB family of receptor tyrosine kinases. Numerous studies have reported the amplification and overexpression of HER2 in several types of cancer, including non-small cell lung cancer (NSCLC). However, the benefits of HER2-targeted therapy have not been fully established. In the present study, the anti-tumor effect of neratinib, an irreversible pan-HER tyrosine kinase inhibitor (TKI), against NSCLC cells harboring *HER2* alterations was investigated. The sensitivity of normal bronchial epithelial cells (BEAS-2B) ectopically overexpressing wild-type or mutant *HER2* to neratinib was assessed. Furthermore, the anti-tumor activity of neratinib in several NSCLC cell lines harboring *HER2* alterations was determined *in vitro* and *in vivo*, and the association between their genetic alterations and sensitivity to neratinib treatment was investigated. BEAS-2B cells ectopically overexpressing wild-type *HER2* or mutants (A775insYVMA, G776VC, G776LC, P780insGSP, V659E, G660D and S310F) exhibited constitutive autophosphorylation of HER2, as determined by western blotting. While these BEAS-2B cells were sensitive to neratinib, they were insensitive to erlotinib, a first-generation epidermal growth factor receptor-TKI. Neratinib also exerted anti-proliferative effects on *HER2*-altered (H2170, Calu-3 and H1781) NSCLC cell lines. Neratinib was also demonstrated to exert strong tumor growth inhibitory activity in mouse xenograft models using *HER2*-altered lung cancer cells. The

results of the present study strongly suggest that neratinib has potential as a promising therapeutic option for the treatment of *HER2*-altered NSCLC.

Introduction

Recent progress in the identification of tumor-specific molecular alterations has contributed to novel therapeutic approaches, and the development of molecular-targeting anti-tumor drugs has improved patient survival. For example, epidermal growth factor receptor (EGFR)-targeted therapy for non-small cell lung cancer (NSCLC) cases harboring *EGFR* oncogenic alterations is a promising strategy for improving the clinical outcome of patients with NSCLC (1,2).

Human epidermal growth factor receptor 2 (HER2) is part of the ErbB family of receptor tyrosine kinases. HER2 is activated by homodimerization or heterodimerization with other receptors in the ErbB family, particularly EGFR (3). HER2 has important roles in pathogenesis of certain types of human cancer, and numerous studies have reported the amplification and overexpression of HER2 in cancer, particularly breast cancer (4-6). The reported frequencies of HER2 overexpression and *HER2* amplification in NSCLC range from 11-32 and 2-23%, respectively (7-10). *HER2* mutations have been identified in 2-4% of all NSCLCs, and are usually mutually exclusive with other driver mutations (11,12). Several *HER2* variants have been reported previously, the majority of which are in-frame insertions in exon 20 of the kinase domain, including A775insYVMA, G776VC, P780insGSP and G776LC (12). Our previous study identified two novel mutations in the HER2 transmembrane domain, which is encoded by exon 17 (V659E and G660D), as rare HER2 variants in lung adenocarcinoma, and the preliminary data suggested that these mutations may be oncogenic (13). An extracellular domain point mutation, S310F, in exon 8 has also been reported to be oncogenic (14). However, the benefit of HER2-targeted therapy against NSCLC harboring *HER2* alterations is far less well defined than the known benefit against breast cancer and gastric cancer with *HER2* alterations (15).

Afatinib (BIBW 2992) is a pan-HER tyrosine kinase inhibitor (TKI) that has been approved for the treatment of patients with NSCLC harboring EGFR mutations. Recently,

Correspondence to: Dr Kazuhiko Shien, Department of Thoracic, Breast and Endocrinological Surgery, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, 2-5-1 Shikata-cho, Kita-ku, Okayama 700-8558, Japan
E-mail: k.shien@okayama-u.ac.jp

Abbreviations: HER2, human epidermal growth factor receptor 2; NSCLC, non-small cell lung cancer; TKI, tyrosine kinase inhibitor

Key words: neratinib, HER2, lung cancer, NSCLC, pan-HER tyrosine kinase inhibitor

afatinib has attracted attention as a HER2-targeting treatment agent. Afatinib was reported to exhibit good clinical activity in patients with lung adenocarcinoma carrying *HER2* mutations (16,17). In preclinical studies, afatinib inhibited the growth of *HER2*-altered NSCLC cells (18).

Neratinib (HKI-272) is another pan-HER TKI that has been reported to improve the overall survival of post-operative patients with HER2-positive breast cancer previously treated with trastuzumab-based adjuvant therapy (19). A phase II clinical trial (PUMA-NER-4201) evaluating the usefulness of neratinib combined with the mechanistic target of rapamycin kinase inhibitor temsirolimus for patients with NSCLC harboring *HER2*-mutations (insertions in exon 20) in currently ongoing; patient accrual has been completed, and the final results of the trial are being awaited (20,21). Thus far, the available data is limited, and the benefits of neratinib treatment remain unclear, particularly for cases with relatively rare mutations. Therefore, the aim of the present study was to investigate the potential use of neratinib against *HER2*-altered NSCLC, including cases with relatively uncommon mutations.

Materials and methods

Cell lines and reagents. Four lung cancer cell lines (A549, Calu-3, NCI-H2170 and NCI-H1781) and one normal human bronchial epithelial cell line (BEAS-2B) were used in the current study. Calu-3, H2170 and H1781 cells were received as gifts from Dr Adi F. Gazdar (University of Texas Southwestern Medical Center at Dallas, Dallas, TX, USA) (22,23). A549 was purchased from American Type Culture Collection (Manassas, VA, USA). BEAS-2B was purchased from European Collection of Authenticated Cell Cultures (Public Health England, Porton Down, UK). All the cancer cell lines were cultured in RPMI-1640 medium supplemented with 10% fetal bovine serum (FBS), and the BEAS-2B cells were maintained in Dulbecco's modified Eagle's medium supplemented with 10% FBS. They were cultured in a humidified atmosphere containing 5% CO₂ at 37°C. Neratinib and erlotinib were purchased from Selleck Chemicals (Houston, TX, USA).

Plasmid constructs and transfection. Human cDNAs encoding full-length *HER2* (wild-type and its variants, A775insYVMA, G776VC, G776LC, P780insGSP, V659E, G660D and S310F) were inserted into the pIDT-SMART (C-TSC) vector, pCMViRTSC (24). Transient transfection of the BEAS-2B cells with the mammalian expression vectors was performed using Lipofectamine[®] 3000 (Thermo Fisher Scientific, Inc., Waltham, MA, USA), according to the manufacturer's protocol.

Western blot analysis and immunohistochemistry. Total cell lysates were extracted using a mixture of radioimmunoprecipitation assay lysis buffer, phosphatase inhibitor cocktails 2 and 3 (Sigma-Aldrich; Merck KGaA, Darmstadt, Germany) and Complete Mini protease inhibitor cocktail (Roche Diagnostics, Basel, Switzerland). Western blot analysis was performed using the conventional method with the following primary antibodies: Anti-EGFR (4267S; 1:1,000), phospho-(p-) EGFR (Tyr1068) (3777S; 1:1,000), *HER2* (4290S; 1:1,000), p-*HER2* (Tyr1221/1222) (2243S; 1:1,000), Akt

(9272S; 1:1,000), p-Akt (Ser473) (4060S; 1:1,000), p44/p42 mitogen-activated protein kinase (MAPK) (9102S; 1:1,000), p-p44/p42 MAPK (4370S; 1:1,000), cleaved poly (ADP-ribose) polymerase (PARP; Asp214) (5625S; 1:1,000) were purchased from Cell Signaling Technology, Inc. (Danvers, MA, USA). β -actin (used as loading control) (MAB1501R; 1:1,000) was purchased from EMD Millipore (Billerica, MA, USA). The secondary antibodies were horseradish peroxidase-conjugated anti-mouse IgG (sc-2031; 1:2,500) or anti-rabbit IgG (sc-2030; 1:2,500; Santa Cruz Biotechnology, Inc., Dallas, TX, USA).

To detect specific signals, the membranes were examined using the ECL Prime Western Blotting Detection System (GE Healthcare Life Sciences, Little Chalfont, UK) and a LAS-3000 imager (Fujifilm, Tokyo, Japan). Immunohistochemical staining with anti-p-*HER2* (1:300 Y1221/1222; Cell Signaling Technology, Inc.) was conducted. The detailed protocol for the immunohistochemical staining has been described previously (25).

Cell growth inhibition assay. Cells were cultured with or without the appropriate drugs for 72 h and the sensitivities of the cells to the drugs were determined via a modified MTS assay using Cell Titer 96 Aqueous One Solution Reagent (Promega Corporation, Madison, WI, USA), as described previously (26). The anti-proliferative activity of each drug is presented as the IC₅₀, which is the concentration of the drug required to inhibit cell proliferation by 50%.

Cell cycle analysis. The effects of neratinib on the cell cycle distribution were assessed using a propidium iodide staining-based assay (CycleTEST PLUS DNA reagent kit; BD Biosciences, Franklin Lakes, NJ, USA) and a BD Accuri C6 flow cytometer (BD Biosciences). Doublets, cell debris and fixation artifacts were gated out, and cell cycle analysis was performed.

Xenograft model. NOD/SCID female mice [n=18, body weight (mean \pm standard)=19.9 \pm 0.6 g, 6-weeks-old] were purchased from Charles River Laboratories, Inc. (Wilmington, MA, USA). All the mice were provided with sterilized food and water, and housed in a barrier facility under a 12:12-h light/dark cycle. Each cell line (5 \times 10⁶ cells) was suspended in 200 μ l RPMI-1640 medium mixed with Matrigel Basement Membrane Matrix (Corning Incorporated, Corning, NY, USA) and subcutaneously injected into the backs of the mice. The tumor volume was calculated using the empirical formula, $V=1/2 \times [(shortest\ diameter)^2 \times (longest\ diameter)]$. When the tumor volume exceeded \sim 50 mm³, the mice were orally administered with vehicle alone or neratinib (40 mg/kg, 6 days a week). Neratinib was prepared in 0.5 w/v (%) methyl cellulose. The tumor volume was measured three times a week using calipers. After 4 weeks of treatment, or when humane endpoints were reached, mice were euthanized by cervical dislocation.

All animal experiments were performed in accordance with protocols approved by the Animal Care and Use Committee of Okayama University (Okayama, Japan; permission no. OKU-2018215) and were conducted in accordance with recent legislation of the National Institutes of Health Guide for the Care and Use of Laboratory Animals.




Figure 1. Overexpression of wild-type or mutant *HER2* activates *HER2* signaling, and neratinib inhibits this signaling pathway. BEAS-2B cells were transiently transfected with WT *HER2*, A775insYVMA, G776VC, G776LC, P780insGSP, G660D, V659E or S310F mutants, or vector control. At 48 h post-transfection, cells were treated with 1.0 μ M erlotinib or 0.1 μ M neratinib for 6 h. Cells were cultured with media supplemented with fetal bovine serum. Lysates were subjected to western blot analysis using the indicated antibodies. *HER2*, human epidermal growth factor receptor 2; WT, wild-type; p-, phosphorylated; EGFR, epidermal growth factor receptor.

Statistical analysis. Statistical analysis was performed using EZR software (Saitama Medical Center, Jichi Medical University, Saitama, Japan), which is a graphical user interface for R (The R Foundation for Statistical Computing, Vienna, Austria). EZR is a modified version of R commander software (version 1.6-3) with additional statistical functions frequently used in biostatistics (27). Data from two groups were compared using t-test. All tests were two-sided. $P < 0.05$ was considered to indicate a statistically significant difference.

Results

***HER2* mutations activate *HER2* signaling, which is inhibited by neratinib.** To examine the effect of *HER2* alterations on the signal transduction pathways, normal bronchial epithelial cells (BEAS-2B) were transiently transfected with vectors containing wild-type *HER2* or one of seven *HER2* mutations: Four kinase domain mutations (A775insYVMA, G776VC, G776LC, and P780insGSP), two transmembrane domain mutations (V659E and G660D) and one extracellular domain mutation (S310F).

The sensitivity of BEAS-2B cells ectopically expressing wild-type or mutant *HER2* to erlotinib (an EGFR-TKI) or neratinib (a pan-HER-TKI) was examined. At 48 h after transfection, the cells were cultured in the presence or absence of erlotinib or neratinib for 6 h. Erlotinib had minimal effect on the phosphorylation of *HER2* and EGFR, whereas neratinib strongly inhibited the phosphorylation of *HER2* and EGFR compared with untreated cells (Fig. 1). These results suggest that the *HER2* alterations were activating mutations and that neratinib treatment had an inhibitory effect on *HER2* activation. Furthermore, the activation of EGFR via cross-phosphorylation of *HER2* was not suppressed by erlotinib treatment.

Neratinib inhibits the growth of *HER2*-amplified and *HER2*-mutant lung cancer cells. The anti-tumor activity of neratinib (a pan-HER-TKIs) and erlotinib (an EGFR-TKI) against *HER2*-driven NSCLC cell lines was subsequently

examined (Table I; Fig. 2B). The cell lines in this panel consisted of two *HER2*-amplified NSCLC cell lines (H2170 and Calu-3) and one *HER2*-mutant NSCLC cell line (H1781). The detailed *HER2* genetic profiles of these three cell lines are presented in Table I based on the results of a previous study (18). The proliferation of the two *HER2*-amplified lung cancer cell lines, H2170 and Calu-3, was inhibited by neratinib, with IC_{50} values of 4.7 and 16.5 nM, respectively. Neratinib also exerted a strong cytotoxic effect against the H1781 cells, with an IC_{50} of 13.6 nM. By contrast, the H2170 and H1781 cell lines were resistant to erlotinib treatment, with IC_{50} values of 1150 and 1080 nM, respectively. Calu-3 cells were partially sensitive to erlotinib, with an IC_{50} of 316 nM. These results were consistent with those of a previous report (Table I; Fig. 2A) (28).

Subsequently, the effect of neratinib on signal transduction pathways in *HER2*-amplified or *HER2*-mutant lung cancer cells was assessed. After 6 h of treatment with neratinib, the cells were lysed and then subjected to western blot analysis. As demonstrated in Fig. 2B, neratinib potently inhibited the phosphorylation of *HER2* and EGFR when administered at concentrations as low as 0.01 μ M, and downstream signals, including phosphorylation of Akt and MAPK, were also inhibited by neratinib in the *HER2*-amplified and the *HER2*-mutant lung cancer cells. Taken together, these results suggest that neratinib has strong anti-tumor activity against *HER2*-amplified and *HER2*-mutant lung cancer cells *in vitro*.

Neratinib induces cell cycle arrest and apoptosis in *HER2*-dependent cells. The effect of neratinib on the cell cycle and apoptosis in *HER2*-driven cells (H2170, Calu-3 and H1781) and *KRAS* mutant cells (A549) was also examined to determine the mechanism of growth inhibition. Cells were treated with 0.1 μ M neratinib for 48 h and then analyzed using flow cytometry. To assess the cell cycle distribution, the sub-G1 fraction was excluded and the percentage of cells in each cycle phase was measured (Fig. 3A). Neratinib treatment caused an increase in the number of H2170, Calu-3 and H1781 cells in G1 phase compared with the distribution

Table I. Characteristics and IC₅₀ values for pan-HER tyrosine kinase inhibitor and EGFR tyrosine kinase inhibitor in non-small cell lung cancer cell lines.

Cell line	Histologic subtype	Gene copy number		Mutation status		Genetic alteration	IC ₅₀ (nM)	
		EGFR	HER2	EGFR	HER2		Neratinib	Erlotinib
H2170	SQ	2	95	WT	WT	HER2 amplification	4.7	1,150
Calu-3	AD	4	111	WT	WT	HER2 amplification	16.5	316
H1781	AD	2	3	WT	G776VC	HER2 mutation	13.6	1,080

SQ, squamous cell carcinoma; AD, adenocarcinoma; IC₅₀, half-maximal inhibitory concentration; WT, wild-type; HER2, human epidermal growth factor receptor 2; EGFR, epidermal growth factor receptor.




Figure 2. Neratinib inhibits *HER2*-amplified and *HER2*-mutant non-small cell lung cancer. (A) *HER2*-amplified and *HER2*-mutant non-small cell lung cancer cells were treated with neratinib or erlotinib for 72 h and the half-maximal inhibitory values were determined using an MTS assay. Error bars indicate the standard deviation. (B) Cells were treated with the indicated concentrations of neratinib for 6 h and the cell lysates were subjected to western blot analysis with the indicated antibodies. HER2, human epidermal growth factor receptor 2; p-, phosphorylated; EGFR, epidermal growth factor receptor; AKT, protein kinase B; MAPK, mitogen activated protein kinase.

in untreated cells; a similar increase was not detected in A549 cells. Subsequently, western blotting was performed to evaluate cell apoptosis, using cleaved PARP antibody as an apoptosis marker. Neratinib induced apoptosis in the H2170, Calu-3 and H1781 cells; however, an increase in cleaved PARP was not detected in A549 cells following neratinib treatment (Fig. 3B). These results suggest that neratinib induces anti-proliferative effects via G1 arrest and apoptotic cell death in *HER2*-altered cells (H2170, Calu-3 and H1781 cells), whereas *HER2*-independent NSCLC cells were not sensitive to neratinib.

Anti-tumor effect of neratinib in a mouse xenograft model of HER2-altered lung cancer. Based on the *in vitro* data, the anti-tumor effect of neratinib was investigated using mouse xenograft models of *HER2*-driven lung cancer. Two *HER2*-amplified (H2170 and Calu-3) cell lines and one *HER2*-mutant (H1781) cell line were used for this experiment. The dose of neratinib was selected based on the results of previous reports (29,30). Once the xenograft tumor volume reached ~50 mm³, the mice were orally treated with the vehicle alone or neratinib (40 mg/kg, 6 days a week). As demonstrated in Fig. 4, neratinib treatment significantly inhibited the tumor




Figure 3. Neratinib induces cell cycle arrest and apoptosis in *HER2*-dependent lung cancer cells. (A) Cells were treated with 0.1 μM neratinib for 48 h and subjected to cell cycle analysis using flow cytometry. The graph represents the percentages of each phase in live cells (mean \pm standard deviation) of triplicate cultures. * $P < 0.05$ vs. neratinib non-treated cells. (B) Cells were treated with 0.1 μM neratinib, then lysates were collected and subjected to western blot analysis with the indicated antibodies. *HER2*, human epidermal growth factor receptor 2; PARP, poly (adenosine diphosphate-ribose) polymerase.

growth of the H2170, Calu-3 and H1781 xenografts compared with the vehicle control ($P < 0.001$). In the H2170 xenograft group treated with the vehicle control, the mice were sacrificed on day 18, as the tumor volumes had reached the humane endpoint ($\sim 2,000 \text{ mm}^3$). The levels of p-*HER2* in the xenografts were examined using immunohistochemistry. The level of p-*HER2* was suppressed in tumors from mice treated with neratinib compared those that received the vehicle control (Fig. 4G-I).

Discussion

The findings of the current study demonstrated the *in vivo* and *in vitro* anti-tumor efficacy of neratinib against lung cancer cells harboring *HER2* alterations. Neratinib is an irreversible human EGFR-TKI that also binds to the tyrosine kinase domains of *HER2* and *HER4*. In patients with breast cancer, neratinib treatment has improved overall survival among post-operative patients with *HER2*-positive breast cancer previously treated with trastuzumab, and neratinib has been approved for extended adjuvant treatment in patients with early-stage *HER2*-positive breast cancer (19). By contrast, only a few reports have discussed the efficacy of neratinib against NSCLC. Furthermore, recent advances in clinical tumor sequencing have identified a large number of variants in oncogenic driver genes with unknown significance, including *EGFR*, *anaplastic lymphoma kinase* and *HER2* variants. Heterogeneity among the functions and/or drug sensitivities of tumors with these mutations has been reported.

Notably, suitable TKI selection for patients with individual oncogene variants, including relatively rare mutations, has been proposed (31). Thus, further preclinical and clinical investigations of patients with tumors harboring uncommon *HER2* gene variants are warranted.

In the current study, neratinib inhibited the growth of cells harboring *HER2* mutations in the transmembrane domain, extracellular domain and kinase domain. Although neratinib has been previously reported to inhibit the growth of cells harboring *HER2* mutations in the kinase domain, to the best of our knowledge, there are no reports of neratinib inhibiting the growth of cells with *HER2* mutations in the transmembrane and extracellular domains (32). Genomic and functional analyses have suggested that mutations in the transmembrane domain of *HER2* encoded by exon 17 (V659E and G660D) are oncogenic in lung adenocarcinoma. Additionally, although the variant rate is low, mutations in the extracellular domain (such as S310F) are also considered to be oncogenic alterations in lung adenocarcinoma (14). For cells harboring mutations in the extracellular domain, anti-*HER2* monoclonal antibodies, such as trastuzumab, can target this region of the receptor and prevent homo-dimerization and receptor activation (33). However, because the kinase domain is constitutively activated in tumors harboring kinase domain mutations, the anti-proliferative effects of monoclonal antibodies may be limited even if dimerization is inhibited (34). The effect of trastuzumab and other antibodies may also be




Figure 4. Neratinib exhibits a strong tumor growth inhibitory effect in a mouse xenograft model of *HER2*-altered lung cancer cells. Mice with (A) H2170, (B) Calu-3 and (C) H1781 tumors were treated with vehicle or neratinib. Tumor volumes were determined on the indicated days following the onset of treatment. Data are presented as the mean \pm standard (n=6). * $P < 0.05$ vs. control. Appearance of (D) H2170, (E) Calu-3 and (F) H1781 tumors following treatment at the time the mice were sacrificed. Representative images of immunohistochemical staining in (G) H2170, (H) Calu-3 and (I) H1781 tumors resected from xenograft model mice. In the H2170 xenograft group treated with the vehicle control, the mice were sacrificed on day 18 for humane reasons as the tumor volumes had reached $\sim 2,000$ mm³. Scale bars, 100 μ m. HER2, human epidermal growth factor receptor 2; p-, phosphorylated; H&E, hematoxylin and eosin.

limited in tumors harboring mutations in the transmembrane domain, as *HER2* dimerization is thought to be stable even if trastuzumab or other antibodies bind to the extracellular domain (35). On the other hand, neratinib targets the kinase domain of *HER2* and inhibits the phosphorylation and activity of *HER* receptors, and therefore may have a therapeutic advantage over trastuzumab and other monoclonal antibodies, as neratinib can exert anti-tumor effects regardless of the domain in which the mutations exist, as demonstrated in the present report. Although *HER2* mutations in the extracellular domain are rare (36), neratinib may be a useful therapeutic option in patients with such mutations.

The findings of the present study demonstrated the efficacy of neratinib against NSCLC cell lines harboring several *HER2* variants. The anti-tumor activity of neratinib was also evaluated in a mouse xenograft model using lung cancer cells with *HER2* amplification or *HER2* mutation. To the best of our knowledge, this is the first report of *in vivo* experiments using neratinib for mouse xenograft models of lung cancer cells harboring *HER2* amplification or *HER2* mutations.

In conclusion, the anti-tumor effect of neratinib against lung cancers harboring *HER2* alterations was demonstrated *in vitro* and *in vivo*. The findings suggest that neratinib has potential as a promising therapeutic option for the treatment of *HER2*-altered NSCLC.

Acknowledgements

The abstract was presented at the Annual Meeting of the American Association for Cancer Research Apr 14-18 2018 in Chicago, IL, USA and published as abstract no. 4777 in *Cancer Res* 78 (Suppl 13), 2018. The authors would like to thank Ms. Fumiko Isobe (Department of Thoracic, Breast and Endocrinological Surgery, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Okayama, Japan) for her technical assistance.

Funding

This study was supported by a Management Expenses Grant.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

YO, KSh, KSu, JS, HY and SToy conceived and designed the study. YO, KSh, and MS performed majority of the *in vitro* experiments. MS performed cell transfection. YO, YT, STom

and KSh performed the statistical analysis. JS, HY, TY, HT, HS, EK and KN participated in the animal experiments and helped perform the analysis with constructive discussion. MS, STom, KSu, KSh and SToy supervised the study.

Ethics approval and consent to participate

All animal experiments were performed in accordance with protocols approved by the Animal Care and Use Committee of Okayama University (Okayama, Japan; permission no. OKU-2018215) and were conducted in accordance with recent legislation of the National Institutes of Health Guide for the Care and Use of Laboratory Animals.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

- Mok TS, Wu YL, Thongprasert S, Yang CH, Chu DT, Saijo N, Sunpaweravong P, Han B, Margono B, Ichinose Y, *et al*: Gefitinib or carboplatin-paclitaxel in pulmonary adenocarcinoma. *N Engl J Med* 361: 947-957, 2009.
- Reck M, Heigener DF, Mok T, Soria JC and Rabe KF: Management of non-small-cell lung cancer: Recent developments. *Lancet* 382: 709-719, 2013.
- Oxnard GR, Binder A and Janne PA: New targetable oncogenes in non-small-cell lung cancer. *J Clin Oncol* 31: 1097-1104, 2013.
- Youlden DR, Cramb SM and Baade PD: The international epidemiology of lung cancer: Geographical distribution and secular trends. *J Thorac Oncol* 3: 819-831, 2008.
- Moasser MM: The oncogene HER2: Its signaling and transforming functions and its role in human cancer pathogenesis. *Oncogene* 26: 6469-6487, 2007.
- Yu D and Hung MC: Overexpression of ErbB2 in cancer and ErbB2-targeting strategies. *Oncogene* 19: 6115-6121, 2000.
- Tan D, Deeb G, Wang J, Slocum HK, Winston J, Wiseman S, Beck A, Sait S, Anderson T and Nwogu C: HER-2/neu protein expression and gene alteration in stage I-III non-small-cell lung cancer: A study of 140 cases using a combination of high throughput tissue microarray, immunohistochemistry, and fluorescent in situ hybridization. *Diagn Mol Pathol* 12: 201-211, 2003.
- Pellegrini C, Falleni M, Marchetti A, Cassani B, Miozzo M, Buttitta F, Roncalli M, Coggi G and Bosari S: HER-2/Neu alterations in non-small cell lung cancer: A comprehensive evaluation by real time reverse transcription-PCR, fluorescence in situ hybridization, and immunohistochemistry. *Clin Cancer Res* 9: 3645-3652, 2003.
- Cappuzzo F, Varella-Garcia M, Shigematsu H, Domenichini I, Bartolini S, Ceresoli GL, Rossi E, Ludovini V, Gregorc V, Toschi L, *et al*: Increased HER2 gene copy number is associated with response to gefitinib therapy in epidermal growth factor receptor-positive non-small-cell lung cancer patients. *J Clin Oncol* 23: 5007-5018, 2005.
- Swanton C, Futreal A and Eisen T: Her2-targeted therapies in non-small cell lung cancer. *Clin Cancer Res* 12: 4377s-4383s, 2006.
- Stephens P, Hunter C, Bignell G, Edkins S, Davies H, Teague J, Stevens C, O'Meara S, Smith R, Parker A, *et al*: Lung cancer: Intragenic ERBB2 kinase mutations in tumours. *Nature* 431: 525-526, 2004.
- Shigematsu H, Takahashi T, Nomura M, Majmudar K, Suzuki M, Lee H, Wistuba II, Fong KM, Toyooka S, Shimizu N, *et al*: Somatic mutations of the HER2 kinase domain in lung adenocarcinomas. *Cancer Res* 65: 1642-1646, 2005.
- Yamamoto H, Higasa K, Sakaguchi M, Shien K, Soh J, Ichimura K, Furukawa M, Hashida S, Tsukuda K, Takigawa N, *et al*: Novel germline mutation in the transmembrane domain of HER2 in familial lung adenocarcinomas. *J Natl Cancer Inst* 106: djt338, 2014.
- Greulich H, Kaplan B, Mertins P, Chen TH, Tanaka KE, Yun CH, Zhang X, Lee SH, Cho J, Ambrogio L, *et al*: Functional analysis of receptor tyrosine kinase mutations in lung cancer identifies oncogenic extracellular domain mutations of ERBB2. *Proc Natl Acad Sci USA* 109: 14476-14481, 2012.
- Mar N, Vredenburg JJ and Wasser JS: Targeting HER2 in the treatment of non-small cell lung cancer. *Lung Cancer* 87: 220-225, 2015.
- De Grève J, Teugels E, Geers C, Decoster L, Galdermans D, De Mey J, Everaert H, Umelo I, In't Veld P and Schallier D: Clinical activity of afatinib (BIBW 2992) in patients with lung adenocarcinoma with mutations in the kinase domain of HER2/neu. *Lung Cancer* 76: 123-127, 2012.
- Mazieres J, Peters S, Lepage B, Cortot AB, Barlesi F, Beau-Faller M, Besse B, Blons H, Mansuet-Lupo A, Urban T, *et al*: Lung cancer that harbors a HER2 mutation: Epidemiologic characteristics and therapeutic perspectives. *J Clin Oncol* 31: 1997-2003, 2013.
- Suzawa K, Toyooka S, Sakaguchi M, Morita M, Yamamoto H, Tomida S, Ohtsuka T, Watanabe M, Hashida S, Maki Y, *et al*: Antitumor effect of afatinib, as a human epidermal growth factor receptor 2-targeted therapy, in lung cancers harboring HER2 oncogene alterations. *Cancer Sci* 107: 45-52, 2016.
- Chan A, Delaloge S, Holmes FA, Moy B, Iwata H, Harvey VJ, Robert NJ, Silovski T, Gokmen E, von Minckwitz G, *et al*: Neratinib after trastuzumab-based adjuvant therapy in patients with HER2-positive breast cancer (ExteNET): A multicentre, randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet Oncol* 17: 367-377, 2016.
- Gandhi L, Besse B, Mazieres J, Waqar S, Cortot A, Barlesi F, Quoix E, Otterson G, Ettinger D, Horn L, *et al*: MA04.02 Neratinib ± temsirolimus in HER2-mutant lung cancers: An international, randomized phase II study. *J Thorac Oncol* 12 (Suppl): S358-S359, 2017.
- Mazieres J, Barlesi F, Filleron T, Besse B, Monnet I, Beau-Faller M, Peters S, Dansin E, Früh M, Pless M, *et al*: Lung cancer patients with HER2 mutations treated with chemotherapy and HER2-targeted drugs: results from the European EUHER2 cohort. *Ann Oncol* 27: 281-286, 2016.
- Gandhi J, Zhang J, Xie Y, Soh J, Shigematsu H, Zhang W, Yamamoto H, Peyton M, Girard L, Lockwood WW, *et al*: Alterations in genes of the EGFR signaling pathway and their relationship to EGFR tyrosine kinase inhibitor sensitivity in lung cancer cell lines. *PLoS One* 4: e4576, 2009.
- Girard L, Zöchbauer-Müller S, Virmani AK, Gazdar AF and Minna JD: Genome-wide allelotyping of lung cancer identifies new regions of allelic loss, differences between small cell lung cancer and non-small cell lung cancer, and loci clustering. *Cancer Res* 60: 4894-4906, 2000.
- Sakaguchi M, Watanabe M, Kinoshita R, Kaku H, Ueki H, Futami J, Murata H, Inoue Y, Li SA, Huang P, *et al*: Dramatic increase in expression of a transgene by insertion of promoters downstream of the cargo gene. *Mol Biotechnol* 56: 621-630, 2014.
- Shien K, Toyooka S, Ichimura K, Soh J, Furukawa M, Maki Y, Muraoka T, Tanaka N, Ueno T, Asano H, *et al*: Prognostic impact of cancer stem cell-related markers in non-small cell lung cancer patients treated with induction chemoradiotherapy. *Lung Cancer* 77: 162-167, 2012.
- Shien K, Toyooka S, Yamamoto H, Soh J, Jida M, Thu KL, Hashida S, Maki Y, Ichihara E, Asano H, *et al*: Acquired resistance to EGFR inhibitors is associated with a manifestation of stem cell-like properties in cancer cells. *Cancer Res* 73: 3051-3061, 2013.
- Kanda Y: Investigation of the freely available easy-to-use software 'EZ' for medical statistics. *Bone Marrow Transplant* 48: 452-458, 2013.
- Engelman JA, Janne PA, Mermel C, Pearlberg J, Mukohara T, Fleet C, Cichowski K, Johnson BE and Cantley LC: ErbB-3 mediates phosphoinositide 3-kinase activity in gefitinib-sensitive non-small cell lung cancer cell lines. *Proc Natl Acad Sci USA* 102: 3788-3793, 2005.
- Schwab CL, English DP, Black J, Bellone S, Lopez S, Cocco E, Bonazzoli E, Bussi B, Predolini F, Ferrari F, *et al*: Neratinib shows efficacy in the treatment of HER2 amplified carcinosarcoma in vitro and in vivo. *Gynecol Oncol* 139: 112-117, 2015.

30. Menderes G, Bonazzoli E, Bellone S, Black JD, Lopez S, Pettinella F, Masserdotti A, Zammataro L, Litkouhi B, Ratner E, *et al*: Efficacy of neratinib in the treatment of HER2/neu-amplified epithelial ovarian carcinoma in vitro and in vivo. *Med Oncol* 34: 91, 2017.
31. Kohsaka S, Nagano M, Ueno T, Suehara Y, Hayashi T, Shimada N, Takahashi K, Suzuki K, Takamochi K, Takahashi F and Mano H: A method of high-throughput functional evaluation of EGFR gene variants of unknown significance in cancer. *Sci Transl Med* 9: pii: eaan6566, 2017.
32. Minami Y, Shimamura T, Shah K, LaFramboise T, Glatt KA, Liniker E, Borgman CL, Haringsma HJ, Feng W, Weir BA, *et al*: The major lung cancer-derived mutants of ERBB2 are oncogenic and are associated with sensitivity to the irreversible EGFR/ERBB2 inhibitor HKI-272. *Oncogene* 26: 5023-5027, 2007.
33. Tai W, Mahato R and Cheng K: The role of HER2 in cancer therapy and targeted drug delivery. *J Control Release* 146: 264-275, 2010.
34. Wang SE, Narasanna A, Perez-Torres M, Xiang B, Wu FY, Yang S, Carpenter G, Gazdar AF, Muthuswamy SK and Arteaga CL: HER2 kinase domain mutation results in constitutive phosphorylation and activation of HER2 and EGFR and resistance to EGFR tyrosine kinase inhibitors. *Cancer Cell* 10: 25-38, 2006.
35. Ou SI, Schrock AB, Bocharov EV, Klempner SJ, Haddad CK, Steinecker G, Johnson M, Gitlitz BJ, Chung J, Campregher PV, *et al*: HER2 transmembrane domain (TMD) mutations (V659/G660) that stabilize homo- and heterodimerization are rare oncogenic drivers in lung adenocarcinoma that respond to Afatinib. *J Thorac Oncol* 12: 446-457, 2017.
36. Eng J, Hsu M, Chaft JE, Kris MG, Arcila ME and Li BT: Outcomes of chemotherapies and HER2 directed therapies in advanced HER2-mutant lung cancers. *Lung Cancer* 99: 53-56, 2016.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.