

RESEARCH PAPER

Inhibition of light-induced stomatal opening by allyl isothiocyanate does not require guard cell cytosolic Ca^{2+} signaling

Wenxiu Ye^{1,2,3}, Eigo Ando⁴, Mohammad Saidur Rhaman², Md. Tahjib-Ul-Arif², Eiji Okuma², Yoshimasa Nakamura², Toshinori Kinoshita^{3,4} and Yoshiyuki Murata^{2,*}

¹ School of Agriculture and Biology, Shanghai Jiao Tong University, Shanghai, 200240 China

² Graduate School of Environmental and Life Science, Okayama University, Tsushima-Naka, Okayama 700-8530, Japan

³ Institute of Transformative Bio-Molecule, Nagoya University, Chikusa, Nagoya, 464-8602, Japan

⁴ Graduate School of Science, Nagoya University, Chikusa, Nagoya, 464-8602, Japan

* Correspondence: muta@cc.okayama-u.ac.jp

Received 31 August 2019; Editorial decision 30 January 2020; Accepted 26 February 2020

Editor: Ian Dodd, Lancaster University, UK

Abstract

The glucosinolate–myrosinase system is a well-known defense system that has been shown to induce stomatal closure in Brassicales. Isothiocyanates are highly reactive hydrolysates of glucosinolates, and an isothiocyanate, allyl isothiocyanate (AITC), induces stomatal closure accompanied by elevation of free cytosolic Ca^{2+} concentration ($[\text{Ca}^{2+}]_{\text{cyt}}$) in *Arabidopsis*. It remains unknown whether AITC inhibits light-induced stomatal opening. This study investigated the role of Ca^{2+} in AITC-induced stomatal closure and inhibition of light-induced stomatal opening. AITC induced stomatal closure and inhibited light-induced stomatal opening in a dose-dependent manner. A Ca^{2+} channel inhibitor, La^{3+} , a Ca^{2+} chelator, EGTA, and an inhibitor of Ca^{2+} release from internal stores, nicotinamide, inhibited AITC-induced $[\text{Ca}^{2+}]_{\text{cyt}}$ elevation and stomatal closure, but did not affect inhibition of light-induced stomatal opening. AITC activated non-selective Ca^{2+} -permeable cation channels and inhibited inward-rectifying K^+ (K^+_{in}) channels in a Ca^{2+} -independent manner. AITC also inhibited stomatal opening induced by fusicoccin, a plasma membrane H^+ -ATPase activator, but had no significant effect on fusicoccin-induced phosphorylation of the penultimate threonine of H^+ -ATPase. Taken together, these results suggest that AITC induces Ca^{2+} influx and Ca^{2+} release to elevate $[\text{Ca}^{2+}]_{\text{cyt}}$, which is essential for AITC-induced stomatal closure but not for inhibition of K^+_{in} channels and light-induced stomatal opening.

Keywords: Allyl isothiocyanate, *Arabidopsis*, calcium channel, potassium channel, proton pump, stomatal closure, stomatal opening.

Introduction

Stomata, surrounded by pairs of guard cells, function as the main window for gas exchange and are in the frontline for defense against microbe invasion in the phyllosphere. To deal with the changing growth and environmental cues, guard cells have evolved to be specialist responders to various stimuli, such as light, drought stress, CO_2 , phytohormones,

and microbe-derived signals, resulting in stomatal movement (Murata *et al.*, 2015; Ye & Murata, 2016; Inoue & Kinoshita, 2017; Takahashi *et al.*, 2018; Zhang *et al.*, 2018).

The glucosinolate–myrosinase system is a well-known defense system against herbivores and pathogens in Brassicales (Halkier & Gershenzon, 2006; Bednarek *et al.*, 2009; Clay *et al.*,

2009). Myrosinases are highly abundant proteins in guard cells and required for abscisic acid (ABA)-induced stomatal movement (Zhao *et al.*, 2008; Islam *et al.*, 2009). Recent results suggest that aliphatic glucosinolates are involved in stomatal movement regulated by ABA and auxin (Zhu *et al.*, 2014; Salehin *et al.*, 2019). Isothiocyanates (ITCs) are highly reactive hydrolyzates of glucosinolates and an ITC, allyl isothiocyanate (AITC), induces stomatal closure in *Arabidopsis* (Khokon *et al.*, 2011; Hossain *et al.*, 2014). AITC reversibly induces stomatal closure in *Vicia faba*, which does not belong to Brassicales (Sobahan *et al.*, 2015). Since many ITCs including AITC are volatile, the results imply that ITCs function as signals for plant–plant interaction (Sobahan *et al.*, 2015). To elucidate the mechanism of AITC-induced stomatal closure, it has been shown that AITC induces production of reactive oxygen species (ROS) and elevation of free cytosolic Ca^{2+} concentration ($[\text{Ca}^{2+}]_{\text{cyt}}$) in guard cells (Khokon *et al.*, 2011). Pharmacological studies indicate that ROS are essential for AITC-induced stomatal closure (Hossain *et al.*, 2014; Sobahan *et al.*, 2015).

ABA-induced stomatal movement has long been studied as a signaling model in guard cells. ABA induces stomatal closure and inhibits light-induced stomatal opening. It was widely thought that the molecular mechanisms underlying these two responses overlap, but recent studies provide evidence that the two signaling pathways can be genetically dissected (Yin *et al.*, 2013). It is important to find out whether other guard cell signaling pathways can be dissected. Though AITC induces stomatal closure, it remains unknown whether it inhibits light-induced stomatal opening.

Cytosolic Ca^{2+} is a well-known second messenger in both stomatal closure and stomatal opening (Shimazaki *et al.*, 2007; Hubbard *et al.*, 2012; Murata *et al.*, 2015). Exogenous Ca^{2+} application is known to induce stomatal closure and inhibit light-induced stomatal opening (Peiter *et al.*, 2005; Islam *et al.*, 2010). Elevations of $[\text{Ca}^{2+}]_{\text{cyt}}$ are triggered by influx of Ca^{2+} from the apoplast and release of Ca^{2+} from intracellular stores in guard cell signaling (Leckie *et al.*, 1998; Hamilton *et al.*, 2000; Pei *et al.*, 2000; Lemtiri-Chlieh *et al.*, 2003; Garcia-Mata *et al.*, 2003). The influx of Ca^{2+} is carried by non-selective Ca^{2+} -permeable cation channels (I_{Ca} channels) that are activated by plasma membrane hyperpolarization (Hamilton *et al.*, 2000; Pei *et al.*, 2000). Pharmacological studies using nicotinamide (NA), an inhibitor of cyclic adenosine diphosphate ribose (cADPR) synthesis (Dodd *et al.*, 2007), suggest that the release of Ca^{2+} is mediated by a cADPR-dependent pathway (Allen *et al.*, 1995; Klüsener *et al.*, 2002; Hossain *et al.*, 2014). It is known that cADPR stimulates ryanodine receptors in the endoplasmic reticulum in animal cells, but homologous genes have not been identified in the *Arabidopsis* genome (Hetherington and Brownlee, 2004), and therefore, it is likely that cADPR functions via a different mechanism in plant cells. Elevation of $[\text{Ca}^{2+}]_{\text{cyt}}$ activates S-type anion channels in guard cells and inhibits plasma membrane H^+ -ATPase and inward-rectifying potassium (K^+) channels (Schroeder & Hagiwara, 1989; Kinoshita *et al.*, 1995; Siegel *et al.*, 2009; Wang *et al.*, 2013a). On the other hand, these transporters are subject to regulation by other components in addition to Ca^{2+} , which probably determine signaling specificity (Allen *et al.*, 2002; Siegel *et al.*, 2009; Xue *et al.*, 2011; Ye

et al., 2015). Furthermore, a Ca^{2+} -independent pathway exists to regulate stomatal movement (Roelfsema & Hedrich, 2010; Laanemets *et al.*, 2013). Though AITC induces $[\text{Ca}^{2+}]_{\text{cyt}}$ elevation, it remains unknown how AITC does this and whether the elevation is essential for AITC-induced stomatal movement. The effects of AITC on I_{Ca} channels and K^+ channels in guard cells remain to be clarified.

In the present study, we aimed to investigate the role of $[\text{Ca}^{2+}]_{\text{cyt}}$ elevation in AITC-induced stomatal closure and inhibition of light-induced stomatal opening in *Arabidopsis*. The presented results reveal that AITC induced influx of Ca^{2+} through activation of I_{Ca} channels and Ca^{2+} release possibly through a cADPR-dependent pathway, resulting in the elevation of $[\text{Ca}^{2+}]_{\text{cyt}}$ in guard cells. While $[\text{Ca}^{2+}]_{\text{cyt}}$ elevation was essential for AITC-induced stomatal closure, it was not essential for inhibition of light-induced stomatal opening. The inhibition was attributed to Ca^{2+} -independent inhibition of K^+ channels by AITC in guard cells.

Materials and methods

Plant materials and growth conditions

Arabidopsis wild-type plants (Columbia) were grown in pots containing a mixture of 70% (v/v) vermiculite (Asahi-kogyo, Okayama, Japan) and 30% (v/v) Kureha soil (Kureha Chemical, Tokyo, Japan) in a growth chamber (photon flux density of $80 \mu\text{mol m}^{-2} \text{s}^{-1}$ under a 16 h light–8 h dark regime). The temperature and relative humidity in the growth chamber were $22 \pm 2 \text{ }^\circ\text{C}$ and $60 \pm 10\%$, respectively. Twice or three times a week, 0.1% Hyponex solution (Hyponex, Osaka, Japan) as a fertilizer was provided to the plants. All the plants used for experiments were from 4 to 6 weeks old.

Stomatal aperture measurement

Fully expanded young leaves from 4- to 5-week-old plants were excised for stomatal aperture measurements as described previously (Ye *et al.*, 2013). For assays of light-induced stomatal opening, leaves were floated on assay solution containing 5 mM KCl, 50 μM CaCl_2 , and 10 mM MES–Tris (pH 6.15) with their adaxial surface upward in the dark for 2 h to close the stomata. After adding AITC, the leaves were kept in the light ($80 \mu\text{mol m}^{-2} \text{s}^{-1}$) for 2 h before measurement. For assays of stomatal closure, leaves were floated on the assay solution in the light for 2 h to open the stomata. Then AITC was added, and the leaves were kept in the light for 2 h before measurement. Inhibitors were added 30 min before AITC treatment. For measurement of stomatal apertures, the leaves were shredded for 30 s, and epidermal tissues were collected using nylon mesh. Thirty stomatal apertures were measured for each sample.

Imaging of $[\text{Ca}^{2+}]_{\text{cyt}}$ in guard cells

Four- to six-week-old wild-type plants expressing YC3.6 were used for the measurement of $[\text{Ca}^{2+}]_{\text{cyt}}$ in guard cells as described previously (Ye *et al.*, 2013, 2015). The abaxial side of an excised leaf was gently mounted on a glass slide with a medical adhesive (stock no. 7730; Hollister) followed by removal of the adaxial epidermis and the mesophyll tissue with a razor blade in order to keep the lower epidermis intact on the slide. The remaining abaxial epidermis was incubated in solution containing 5 mM KCl, 50 μM CaCl_2 , and 10 mM MES–Tris (pH 6.15) in the light for 2 h at $22 \text{ }^\circ\text{C}$ to promote stomatal opening. Turgid guard cells were used to measure $[\text{Ca}^{2+}]_{\text{cyt}}$. The observation chamber was perfused with the solution using a peristaltic pump. Guard cells were incubated in the absence of AITC for 5 min and then in the presence of AITC. Inhibitors were added at the time point indicated. For dual-emission ratio imaging of YC3.6, we used an FF-02-438/24-25 (Semrock) excitation filter, a 445DRLP

(Omega) dichroic mirror, an XF3075 480AF30 (Omega) emission filter for cyan fluorescent protein (CFP), and an XF3011 535DF25 (Omega) emission filter for yellow fluorescent protein (YFP). The CFP and YFP fluorescence intensity of guard cells were imaged and analysed using the W-View system and AQUA COSMOS software (Hamamatsu Photonics). CFP and YFP fluorescence were simultaneously monitored.

Patch-clamp measurement

Current measurements of I_{Ca} and K^+ channels in Arabidopsis guard cells were performed as described previously (Mori et al., 2006; Ye et al., 2013). Arabidopsis guard cell protoplasts were prepared from rosette leaves. For I_{Ca} current measurement, the pipette solution contained 10 mM BaCl₂, 0.1 mM dithiothreitol, 4 mM EGTA, and 10 mM HEPES–Tris, pH 7.1. The bath solution contained 100 mM BaCl₂, 0.1 mM dithiothreitol, and 10 mM MES–Tris, pH 5.6. For K^+ channel current measurement, the pipette solution contained 30 mM KCl, 70 mM K–Glu, 2 mM MgCl₂, 2.4 mM CaCl₂ (free Ca²⁺ concentration, 150 nM), 6.7 mM EGTA, and 10 mM HEPES–Tris, pH 7.1. The bath solution contained 30 mM KCl, 2 mM MgCl₂, 40 mM CaCl₂, and 10 mM MES–Tris, pH 5.5. In all cases, osmolality was adjusted to 500 mmol kg⁻¹ (pipette solutions) and 485 mmol kg⁻¹ (bath solutions) with D-sorbitol.

KAT1 current recording in *Xenopus laevis* oocytes

The expression of KAT1 in *Xenopus laevis* oocytes and current recording were performed according to our previous method (Islam et al., 2015). Before recording, the microinjected oocytes were incubated in ND96 solution containing 94 mM NaCl, 2 mM KCl, 1 mM CaCl₂, 1 mM MgCl₂, and 5 mM HEPES (pH 7.5) with different concentrations of AITC for 2 h.

Immunohistochemical detection of the plasma membrane H⁺-ATPase in guard cells using a whole leaf

Immunohistochemistry using a whole leaf was performed according to Sauer et al. (2006) and Hayashi et al. (2011) with modifications. Mature leaves were harvested from dark-adapted plants and floated on the basal buffer (5 mM MES–BTP (pH 6.5), 50 mM KCl, and 0.1 mM CaCl₂) containing 50 μM AITC for 20 min in the dark. After AITC treatment, 10 μM fusicoccin was added to the buffer and kept for a further 10 min. For the control, 0.1% (v/v) dimethyl sulfoxide was added to the buffer. After treatment, leaves were put into a syringe with fixative (4% (w/v) formaldehyde freshly prepared from paraformaldehyde and 0.3% (v/v) glutaraldehyde in 50 mM PIPES–NaOH (pH 7.0), 5 mM MgSO₄, and 5 mM EGTA), and negative pressure applied several times to infiltrate the fixative, followed by immersion in the solution for 1 h in the dark at room temperature. After washing with phosphate-buffered saline (PBS; 137 mM NaCl, 8.1 mM Na₂HPO₄, 2.68 mM KCl, and 1.47 mM KH₂PO₄), chlorophyll was removed by pure methanol (20 min incubation at 37 °C three or four times). Then, central areas of the leaves were cut out, and incubated with xylene at 37 °C for 2 min, pure ethanol at room temperature for 5 min, and 50% (v/v; in PBS) ethanol at room temperature for 5 min, and washed with Milli-Q water twice. The material was transferred to MAS-coated microscope slides (Matsunami) with a droplet of water, where the abaxial side of the leaf was attached to the slide, and freeze–thaw treatment applied followed by complete drying overnight at room temperature. Dried samples were rehydrated by PBS for 5 min at room temperature, and digested with 4% (w/v) Cellulase Onozuka R-10 (Yakult) with 0.5% (w/v) Macerozyme R-10 (Yakult) in PBS for 1 h at 37 °C. After digestion, leaf tissue except for the abaxial epidermis attached on the slide was removed stereomicroscopically in PBS, and the left epidermal tissue was washed four times for 5 min each with PBS, then permeabilized with 3% (v/v) IGEPAL CA-630 (MP Biomedicals) with 10% (v/v) dimethyl sulfoxide in PBS for 1 h at room temperature. Samples were washed five times for 5 min each with PBS and incubated with blocking solution (3% (w/v) bovine serum albumin Fraction V (BSA; Thermo Fisher Scientific) in PBS) for 1 h at room temperature. The primary antibody (anti-pThr; Hayashi et al., 2010) was treated with a dilution of 1:1000 in the blocking solution at 4 °C overnight.

Samples were washed five times for 5 min each with PBS; secondary antibody (Alexa Fluor 488-conjugated goat anti-rabbit IgG; Thermo Fisher Scientific) with a dilution of 1:500 in the blocking solution was applied at 37 °C for 3 h, followed by washing five times for 5 min each with PBS. The specimens were covered by a coverglass with 50% (v/v) glycerol. Fluorescence images were obtained (Hayashi et al., 2011) and analysed by ImageJ software (National Institutes of Health).

Statistical analysis

The significance of differences between datasets was assessed by Student's *t*-test and analysis of variance (ANOVA) with Tukey's test. The response of $[Ca^{2+}]_{cyt}$ was assessed by χ^2 test. Differences were considered significant for $P < 0.05$.

Results

Effect of inhibitors of Ca²⁺ influx and Ca²⁺ release pathways on allyl isothiocyanate-induced stomatal movement

To investigate AITC-induced stomatal closure, intact leaves were initially incubated in the light to open the stomata followed by AITC treatment in the light. For investigation of AITC inhibition of light-induced stomatal opening, leaves were initially incubated in the dark to close the stomata followed by AITC treatment in the light. AITC induced stomatal closure in a dose-dependent manner (Fig. 1A), which is consistent with previous results (Khokon et al., 2011). In addition, AITC inhibited light-induced stomatal opening in a dose-dependent manner (Fig. 1B).

Previous results have shown that AITC induces stomatal closure accompanied by $[Ca^{2+}]_{cyt}$ elevation. However, it remains unknown whether the $[Ca^{2+}]_{cyt}$ elevation is essential for AITC-induced stomatal closure. Application of a Ca²⁺ channel blocker, La³⁺, and a Ca²⁺ chelator, EGTA, inhibited AITC-induced stomatal closure in a dose-dependent manner (Fig. 2A). Nicotinamide, an inhibitor of Ca²⁺ release, also inhibited AITC-induced stomatal closure in a dose-dependent manner (Fig. 2B). These results suggest that $[Ca^{2+}]_{cyt}$ elevation induced by Ca²⁺ influx and Ca²⁺ release is essential for AITC-induced stomatal closure.

On the other hand, the three inhibitors even at the highest concentration for stomatal closure assay (Fig. 2) did not significantly affect AITC inhibition of light-induced stomatal opening (Fig. 3). These results suggest that $[Ca^{2+}]_{cyt}$ elevation is not essential for AITC inhibition of light-induced stomatal opening.

Since intact leaves were used for stomatal movement assay, AITC and inhibitors have to be transported via the petiole and/or penetrate cuticle directly to reach the guard cells. Therefore, it is possible that concentrations of AITC and inhibitors in the apoplast of intact leaves are lower than those in the assay solution.

Effect of inhibitors of Ca²⁺ influx and Ca²⁺ release pathways on allyl isothiocyanate-induced $[Ca^{2+}]_{cyt}$ elevation

Previous results have shown that AITC induced $[Ca^{2+}]_{cyt}$ elevations in guard cells (Khokon et al., 2011). Here, we investigated




Fig. 1. Effect of AITC on stomatal movement. (A) Stomatal closure induced by AITC. (B) Inhibition by AITC of light-induced stomatal opening. Averages from three independent experiments (90 stomata in total per bar) are shown. Error bars represent standard error of the mean ($n=3$). *Statistical significance compared with 0 µM AITC ($P<0.05$).




Fig. 2. Effects of La³⁺, EGTA, and NA on AITC-induced stomatal closure. (A) Effects of La³⁺ and EGTA on AITC-induced stomatal closure. (B) Effect of NA on AITC-induced stomatal closure. Averages from three independent experiments (90 stomata in total per bar) are shown. Error bars represent standard error of the mean ($n=3$). Different letters indicate statistical significance between groups ($P<0.05$).

how AITC induces $[Ca^{2+}]_{cyt}$ elevations in guard cells from isolated epidermal tissues. Mock treatment showed $[Ca^{2+}]_{cyt}$ elevations in 16.7% of guard cells (Fig. 4A, F), whereas application of 50 µM AITC induced elevations in 91.4% of wild-type guard cells (Fig. 4B, F), which is consistent with previous results (Khokon *et al.*, 2011). Application of 1 mM La³⁺, 2 mM EGTA and 50 mM NA abolished the AITC-induced $[Ca^{2+}]_{cyt}$ elevations (Fig. 4C–F). These results indicate that the inhibitors do suppress AITC-induced $[Ca^{2+}]_{cyt}$ elevations and that both Ca²⁺ influx and Ca²⁺ release contribute to the $[Ca^{2+}]_{cyt}$ elevations.

Other Ca²⁺ channel inhibitors, Gd³⁺, verapamil (VERA) and ruthenium red (RR) (Allen *et al.*, 1995; McAinsh *et al.*, 1995; Cessna *et al.*, 1998), were used to further investigate AITC-induced $[Ca^{2+}]_{cyt}$ elevations. One millimolar Gd³⁺

abolished and 1 mM VERA slightly impaired $[Ca^{2+}]_{cyt}$ elevations induced by 50 µM AITC (Fig. S1A, B, D). On the other hand, RR at 100 µM did not affect the $[Ca^{2+}]_{cyt}$ elevations significantly (Fig. S1C, D). These results suggest that Gd³⁺- and VERA-sensitive Ca²⁺ channels are involved in AITC-induced $[Ca^{2+}]_{cyt}$ elevations.

Effect of allyl isothiocyanate on I_{Ca} currents and K^+_{in} currents in guard cell protoplasts

Application of 50 µM AITC significantly activated I_{Ca} currents when the membrane was hyperpolarized, and the currents were inhibited by the Ca²⁺ channel inhibitor La³⁺ (Fig. 5). These results indicate that AITC induces Ca²⁺ influx through activation of I_{Ca} channels.




Fig. 3. Effects of La³⁺, EGTA, and NA on inhibition by AITC of light-induced stomatal opening. Averages from three independent experiments (90 stomata in total per bar) are shown. Error bars represent standard error of the mean ($n=3$). Different letters indicate statistical significance between groups ($P<0.05$).

Potassium influx through K⁺_{in} channels is critical for stomatal opening (Kwak et al., 2001; Takahashi et al., 2013). Resting [Ca²⁺]_{cyt} has been shown to be in the range of 120–150 nM in guard cells (Grabov & Blatt, 1999; Siegel et al., 2009). Since AITC inhibition of light-induced stomatal opening is not dependent on [Ca²⁺]_{cyt} elevation, the effect of AITC on K⁺_{in} currents at resting [Ca²⁺]_{cyt} buffered to 150 nM was investigated. Application of 50 µM AITC significantly suppressed K⁺_{in} currents in guard cell protoplasts (Fig. 6A, B), indicating that AITC suppresses K⁺_{in} channels in a Ca²⁺-independent manner.

The effect of AITC on a major K⁺_{in} channel in guard cells, KAT1, was investigated using the two-electrode voltage-clamp technique. AITC at 50, 100, and 500 µM had no significant effect on the currents seen in *Xenopus* oocytes expressing KAT1 (Fig. 6C).

Effect of allyl isothiocyanate on fusicoccin-induced stomatal opening and phosphorylation of penultimate threonine of plasma membrane H⁺-ATPases

To further investigate how AITC inhibits stomatal opening, the effect of AITC on stomatal opening induced by a plasma membrane H⁺-ATPase activator, fusicoccin (FC), was investigated. Treatment of 50 µM AITC significantly inhibited FC-induced stomatal opening in the dark (Fig. 7). Since FC induces stomatal opening through activation of H⁺-ATPases by increasing the phosphorylation level of the penultimate Thr (penThr) of H⁺-ATPases (Kinoshita & Shimazaki, 2001), the effect of AITC on FC-induced penThr phosphorylation was investigated. Application of 50 µM AITC did not affect FC-induced penThr phosphorylation significantly (Fig. 8). Taken together, these results indicate that AITC inhibits FC-induced stomatal opening without affecting penThr phosphorylation.

Discussion

Allyl isothiocyanate induces stomatal movement like many other abiotic and biotic stimuli

The glucosinolate–myrosinase system is widely known to be activated by tissue damage caused by herbivory, and hydrolysis products of glucosinolates have repellent effects on many herbivores (Halkier & Gershenzon, 2006). Studies have also shown that hydrolysis of glucosinolates is induced in live cells by fungal and bacterial infection and plant hormones such as ABA (Zhao et al., 2008; Bednarek et al., 2009; Clay et al., 2009; Fan et al., 2011; Zhu et al., 2014; Andersson et al., 2015). Guard cell responses to ABA were impaired in aliphatic glucosinolate-deficient (Zhu et al., 2014) and myrosinase-deficient (Zhao et al., 2008; Islam et al., 2009) mutants, suggesting that hydrolysis products of glucosinolates function as signaling component in ABA signaling. There is also evidence for involvement of glucosinolate hydrolysis in signaling induced by bacterial flagellin peptide, flg22 (Clay et al., 2009). These results motivate further study of the signaling by the hydrolysis product of glucosinolates. Among the hydrolysis products, isothiocyanates are highly active and their physiological functions in both abiotic and biotic stress are under intensive investigation (Hara et al., 2013; Andersson et al., 2015). Allyl isothiocyanate is biosynthesized in Arabidopsis (Lambrix et al., 2001) and is one of the most studied isothiocyanates so far (Khokon et al., 2011; Hossain et al., 2013; Åsberg et al., 2015; Øverby et al., 2015; Sobahan et al., 2015). While AITC at concentrations above millimolar is detrimental (Hara et al., 2010), AITC at lower concentrations functions as a signal to trigger physiological events (Khokon et al., 2011; Sobahan et al., 2015). It was also suggested that AITC, as a volatile compound, functions as a signal for plant–plant interaction (Sobahan et al., 2015). Previous results have shown that AITC reversibly induces stomatal closure (Khokon et al., 2011; Hossain et al., 2013; Sobahan et al., 2015). In the present study, it was further revealed that AITC also inhibited light-induced stomatal opening (Fig. 1). Many abiotic and biotic stimuli, such as ABA and flg22, also induce both stomatal closure and inhibition of light-induced stomatal opening (Melotto et al., 2006; Zhang et al., 2008; Yin et al., 2013). Therefore, it is likely that there is crosstalk between AITC signaling and signaling induced by other stimuli including ABA and flg22.

Allyl isothiocyanate induces Ca²⁺ influx and Ca²⁺ release leading to [Ca²⁺]_{cyt} elevations

Activation of I_{Ca} channels in guard cells is triggered by many stimuli, such as ABA, methyl jasmonate (MeJA), yeast elicitor, and amino acids (Murata et al., 2001; Munemasa et al., 2007; Ye et al., 2013; Yoshida et al., 2016; Kong et al., 2016). In the present study, AITC activated I_{Ca} channels in guard cells (Fig. 5). On the other hand, the molecular nature of I_{Ca} channels is mostly unknown. It is known that AITC activates transient receptor potential ankyrin 1, a Ca²⁺ channel, by reacting with several Cys residues in its N-terminus to induce Ca²⁺ influx in neuron cells (Bautista et al., 2006; Hinman et al., 2006). However,




Fig. 4. Effects of La^{3+} , EGTA, and NA on AITC-induced elevation of $[\text{Ca}^{2+}]_{\text{cyt}}$ in guard cells. (A–E) Representative traces of fluorescence emission ratios (535/480 nm) showing transient $[\text{Ca}^{2+}]_{\text{cyt}}$ elevations in guard cells; 1 mM La^{3+} , 2 mM EGTA, and 50 mM NA were added 4 min before 50 μM AITC treatment. (F) Percentage of guard cells showing 0, 1 or 2, or ≥ 3 transient $[\text{Ca}^{2+}]_{\text{cyt}}$ elevations. $[\text{Ca}^{2+}]_{\text{cyt}}$ elevations were counted when changes in fluorescence emission ratios were ≥ 0.1 U from the baseline.




Fig. 5. Activation of I_{Ca} channel currents by AITC in guard cell protoplasts (GCPs). (A) Representative I_{Ca} current traces in GCPs. (B) Average of current–voltage curves for AITC activation of I_{Ca} currents in GCPs ($n=5$) as recorded in (A) (open circles, control; filled circles, 50 μM AITC; open square, La^{3+} +AITC). A ramp voltage protocol from +20 to -180 mV (holding potential, 0 mV; ramp speed, 200 mV s^{-1}) was used. After attaining the whole-cell configuration, GCPs were recorded to obtain control data. To obtain data for AITC treatment and La^{3+} +AITC treatment, recordings were performed sequentially after extracellularly adding AITC and 1 mM La^{3+} . The GCPs were measured 16 times to get averages for each recording. The interpulse period was 1 min. *Statistical significance ($P < 0.05$). Results are from five independent experiments; error bars indicate SE.




Fig. 6. Effect of AITC on K^+ currents in GCPs and KAT1 activity expressed in *Xenopus* oocytes. (A) K^+ currents in GCPs treated without (top trace) or with (bottom trace) 50 μ M AITC. (B) Steady-state current-voltage relationship for AITC inhibition of K^+ currents in WT GCPs as recorded in (A) (open circles, control; filled circles, AITC). The voltage protocol was stepped up from 0 mV to -180 mV in 20 mV decrements (holding potential, -40 mV). GCPs were treated with AITC for 2 h before recordings. Each data point was obtained from at least seven GCPs in more than five independent experiments. Error bars represent standard errors. *Statistical significance compared with Control ($P < 0.05$). (C) Steady-state current-voltage relationship for KAT1-mediated currents in *Xenopus* oocytes. Oocytes were treated with AITC for 2 h before recordings. The voltage protocol was stepped up from 0 mV to -180 mV in 20 mV decrements (holding potential, -40 mV) with a pulse duration of 3 s. Each data point was obtained from seven oocytes in more than three independent experiments. Error bars represent standard errors.




Fig. 7. Effect of AITC on FC-induced stomatal opening in the dark. Averages from three independent experiments (90 stomata in total per bar) are shown. AITC was added 30 min before FC treatment. Error bars represent standard error of the mean ($n = 3$). *Statistical significance ($P < 0.05$).

plants do not harbor genes related to transient receptor potential channels in animal cells (Hedrich, 2012). Studies have identified that hyperosmolality-induced $[Ca^{2+}]_{cyt}$ increase 1 (OSCA1) family members, glutamate receptor homologs, and cyclic nucleotide-gated channels are I_{Ca} channels activated by different stimuli (Wang et al., 2013b; Yuan et al., 2014; Kong et al., 2016; Tian et al., 2019). It would be interesting to investigate whether these channels are activated by AITC.

Previous studies have shown not only Ca^{2+} influx but also Ca^{2+} release from intracellular stores is essential for $[Ca^{2+}]_{cyt}$ elevations induced by ABA, methyl jasmonate and flg22 (Leckie et al., 1998; Hamilton et al., 2000; Pei et al., 2000; Garcia-Mata et al., 2003; Lemtiri-Chlieh et al., 2003; Munemasa et al., 2007; Hossain et al., 2014; Thor and Peiter, 2014). ABA and MeJA recruit a Ca^{2+} release pathway involving cADPR, which is sensitive to NA (Allen et al., 1995; Klüsener et al., 2002; Hossain et al., 2014). In the present study, NA abolished AITC-induced $[Ca^{2+}]_{cyt}$ elevations, suggesting that the NA-sensitive Ca^{2+} release mechanism is conserved among different signaling pathways.




Fig. 8. Immunohistochemical detection of FC-induced penultimate Thr phosphorylation of H^+ -ATPases in guard cell plasma membrane. The vertical scale represents fluorescence levels of guard cells detected using penultimate Thr antiserum as primary antibody and Alexa Fluor 488-conjugated goat anti-rabbit IgG as secondary antibody. The fluorescence level was measured by ImageJ, expressed as relative values normalized to that of mock treatment. Typical fluorescence and the corresponding bright field images are shown at the top. Arrowhead indicates the guard cells. Scale bar: 50 μ m. Data represent means with SDs ($n=3$). NS, no significant difference observed ($P>0.1$; Student's t test). Results are from more than three independent experiments.

Elevation of $[Ca^{2+}]_{cyt}$ is essential for allyl isothiocyanate-induced stomatal closure but not inhibition of light-induced stomatal opening

The driving force for stomatal closure is membrane depolarization induced by activation of S-type anion channels, while the driving force for stomatal opening is membrane hyperpolarization induced by activation of H^+ -ATPase. Ca^{2+} positively regulates S-type anion channels but negatively regulates H^+ -ATPase through Ca^{2+} sensor-dependent pathways (Schroeder & Hagiwara, 1989; Kinoshita *et al.*, 1995), and thus Ca^{2+} is essential for stomatal closure induced by many stimuli such as ABA, MeJA, and yeast elicitor (Mori *et al.*, 2006; Munemasa *et al.*, 2011; Ye *et al.*, 2013). Recent studies have shown quadruple loss-of-function mutations of ABA receptors impair ABA-induced stomatal closure and $[Ca^{2+}]_{cyt}$ elevation but not inhibition of light-induced stomatal opening (Wang

et al., 2013a; Yin *et al.*, 2013). These results highlight that the signaling pathways underlying stomatal closure and inhibition of stomatal opening can be genetically dissected. In the present study, inhibition of $[Ca^{2+}]_{cyt}$ by three inhibitors impaired AITC-induced stomatal closure but not inhibition of light-induced stomatal opening (Figs 2, 3), indicating $[Ca^{2+}]_{cyt}$ elevation is essential for AITC-induced stomatal closure but not inhibition of light-induced opening. Taken together, these results suggest that signaling leading to stomatal closure does not overlap with signaling leading to inhibition of light-induced stomatal opening in different stress responses.

Inhibition of K^+ channels suppresses stomatal opening driven by activation of H^+ -ATPase (Kwak *et al.*, 2001; Takahashi *et al.*, 2013). In the present study, AITC inhibited K^+ channels in the absence of $[Ca^{2+}]_{cyt}$ elevation (Fig. 6). Further results showed that AITC inhibited FC-induced stomatal opening but not phosphorylation of penThr (Fig. 7, 8). These results

again suggest that inhibition of K^+_{in} channels contributes to AITC inhibition of light-induced stomatal opening.

As an electrophile, AITC is prone to forming covalent adducts with amino acid residues, such as Cys, Lys, and His, to modify protein function. It is known that the covalent modification of Cys is reversible under physiological conditions (Nakamura and Miyoshi, 2010; Higdon *et al.*, 2012). It has been shown that two electrophiles, acrolein and methylglyoxal, inhibit K^+_{in} channels in guard cells and target KAT1, a main K^+_{in} channel in guard cells, expressed in a heterologous system using *Xenopus* oocytes (Hoque *et al.*, 2012; Islam *et al.*, 2015, 2016). However, our experiments show that AITC at concentration of 50, 100, and 500 μ M did not significantly affect KAT1 activity expressed in oocytes (Fig. 6C), suggesting that AITC does not directly modify KAT1. It has been reported that one AITC target is microtubules in *Arabidopsis* (Overby *et al.*, 2015) and microtubules are critical for light-induced stomatal opening (Marcus *et al.*, 2001; Eisinger *et al.*, 2012). In the future, it would be interesting to investigate whether AITC modulates K^+_{in} channels through a microtubule-dependent pathway.

Conclusion

The presented results suggest that AITC triggers Ca^{2+} influx and Ca^{2+} release to induce $[Ca^{2+}]_{cyt}$ elevation, which is essential for AITC-induced stomatal closure but not for inhibition of K^+_{in} channels or light-induced stomatal opening.

Supplementary data

Supplementary data are available at *JXB* online.

Fig. S1. Effects of Gd^{3+} , VERA, and RR on AITC-induced elevation of $[Ca^{2+}]_{cyt}$ in guard cells.

Acknowledgements

This work was supported by National Natural Science Foundation of China (31901984 to WY), the Start-up Fund from Shanghai Jiao Tong University (WF220515002 to WY), Grants-in-Aid for Japan Society for the Promotion of Science Fellows from the Japan Society for the Promotion of Science (267977 to WY), Grants-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology, Japan (15H05956 to TK) and the Advanced Low Carbon Technology Research and Development Program from the Japan Science and Technology Agency (to TK).

Author contributions

WY, YN, TK, and YM conceived the research plans; YM and TK supervised the experiments; WY, EA, MSR, MT, and EO performed the experiments; WY wrote the article with the contribution of all the authors.

References

Allen GJ, Muir SR, Sanders D. 1995. Release of Ca^{2+} from individual plant vacuoles by both $InsP_3$ and cyclic ADP-ribose. *Science* **268**, 735–737.

Allen GJ, Murata Y, Chu SP, Nafisi M, Schroeder JI. 2002. Hypersensitivity of abscisic acid-induced cytosolic calcium increases in the *Arabidopsis* farnesyltransferase mutant era1-2. *The Plant Cell* **14**, 1649–1662.

Andersson MX, Nilsson AK, Johansson ON, *et al.* 2015. Involvement of the electrophilic isothiocyanate sulforaphane in *Arabidopsis* local defense responses. *Plant Physiology* **167**, 251–261.

Åsberg SE, Bones AM, Øverby A. 2015. Allyl isothiocyanate affects the cell cycle of *Arabidopsis thaliana*. *Frontiers in Plant Science* **6**, 364.

Bautista DM, Jordt SE, Nikai T, Tsuruda PR, Read AJ, Poblete J, Yamoah EN, Basbaum AI, Julius D. 2006. TRPA1 mediates the inflammatory actions of environmental irritants and proalgesic agents. *Cell* **124**, 1269–1282.

Bednarek P, Pislewska-Bednarek M, Svatos A, *et al.* 2009. A glucosinolate metabolism pathway in living plant cells mediates broad-spectrum antifungal defense. *Science* **323**, 101–106.

Cessna SG, Chandra S, Low PS. 1998. Hypo-osmotic shock of tobacco cells stimulates Ca^{2+} fluxes deriving first from external and then internal Ca^{2+} stores. *The Journal of Biological Chemistry* **273**, 27286–27291.

Clay NK, Adio AM, Denoux C, Jander G, Ausubel FM. 2009. Glucosinolate metabolites required for an *Arabidopsis* innate immune response. *Science* **323**, 95–101.

Dodd AN, Gardner MJ, Hotta CT, *et al.* 2007. The *Arabidopsis* circadian clock incorporates a cADPR-based feedback loop. *Science* **318**, 1789–1792.

Eisinger W, Ehrhardt D, Briggs W. 2012. Microtubules are essential for guard-cell function in *Vicia* and *Arabidopsis*. *Molecular Plant* **5**, 601–610.

Fan J, Crooks C, Creissen G, Hill L, Fairhurst S, Doerner P, Lamb C. 2011. *Pseudomonas sax* genes overcome aliphatic isothiocyanate-mediated non-host resistance in *Arabidopsis*. *Science* **331**, 1185–1188.

Garcia-Mata C, Gay R, Sokolovski S, Hills A, Lamattina L, Blatt MR. 2003. Nitric oxide regulates K^+ and Cl^- channels in guard cells through a subset of abscisic acid-evoked signaling pathways. *Proceedings of the National Academy of Sciences, USA* **100**, 11116–11121.

Grabov A, Blatt MR. 1999. A steep dependence of inward-rectifying potassium channels on cytosolic free calcium concentration increase evoked by hyperpolarization in guard cells. *Plant Physiology* **119**, 277–288.

Halkier BA, Gershenzon J. 2006. Biology and biochemistry of glucosinolates. *Annual Review of Plant Biology* **57**, 303–333.

Hamilton DW, Hills A, Kohler B, Blatt MR. 2000. Ca^{2+} channels at the plasma membrane of stomatal guard cells are activated by hyperpolarization and abscisic acid. *Proceedings of the National Academy of Sciences, USA* **97**, 4967–4972.

Hara M, Harazaki A, Tabata K. 2013. Administration of isothiocyanates enhances heat tolerance in *Arabidopsis thaliana*. *Plant Growth Regulation* **69**, 71–77.

Hara M, Yatsuzuka Y, Tabata K, Kuboi T. 2010. Exogenously applied isothiocyanates enhance glutathione *S*-transferase expression in *Arabidopsis* but act as herbicides at higher concentrations. *Journal of Plant Physiology* **167**, 643–649.

Hayashi M, Inoue S, Takahashi K, Kinoshita T. 2011. Immunohistochemical detection of blue light-induced phosphorylation of the plasma membrane H^+ -ATPase in stomatal guard cells. *Plant & Cell Physiology* **52**, 1238–1248.

Hayashi Y, Nakamura S, Takemiya A, Takahashi Y, Shimazaki K, Kinoshita T. 2010. Biochemical characterization of in vitro phosphorylation and dephosphorylation of the plasma membrane H^+ -ATPase. *Plant and Cell Physiology* **51**, 1186–1196.

Hedrich R. 2012. Ion channels in plants. *Physiological Reviews* **92**, 1777–1811.

Hetherington AM, Brownlee C. 2004. The generation of Ca^{2+} signals in plants. *Annual Review of Plant Biology* **55**, 401–427.

Higdon A, Diers AR, Oh JY, Landar A, Darley-Usmar VM. 2012. Cell signalling by reactive lipid species: new concepts and molecular mechanisms. *The Biochemical Journal* **442**, 453–464.

Hinman A, Chuang HH, Bautista DM, Julius D. 2006. TRP channel activation by reversible covalent modification. *Proceedings of the National Academy of Sciences, USA* **103**, 19564–19568.

Hoque TS, Okuma E, Uraji M, Furuichi T, Sasaki T, Hoque MA, Nakamura Y, Murata Y. 2012. Inhibitory effects of methylglyoxal on

- light-induced stomatal opening and inward K⁺ channel activity in *Arabidopsis*. *Bioscience, Biotechnology, and Biochemistry* **76**, 617–619.
- Hossain MA, Ye W, Munemasa S, Nakamura Y, Mori IC, Murata Y.** 2014. Cyclic adenosine 5'-diphosphoribose (cADPR) cyclic guanosine 3',5'-monophosphate positively function in Ca²⁺ elevation in methyl jasmonate-induced stomatal closure, cADPR is required for methyl jasmonate-induced ROS accumulation NO production in guard cells. *Plant Biology* **16**, 1140–1144.
- Hossain MS, Ye W, Hossain MA, Okuma E, Uraji M, Nakamura Y, Mori IC, Murata Y.** 2013. Glucosinolate degradation products, isothiocyanates, nitriles, and thiocyanates, induce stomatal closure accompanied by peroxidase-mediated reactive oxygen species production in *Arabidopsis thaliana*. *Bioscience, Biotechnology, and Biochemistry* **77**, 977–983.
- Hubbard KE, Siegel RS, Valerio G, Brandt B, Schroeder JI.** 2012. Abscisic acid and CO₂ signalling via calcium sensitivity priming in guard cells, new CDPK mutant phenotypes and a method for improved resolution of stomatal stimulus-response analyses. *Annals of Botany* **109**, 5–17.
- Inoue SI, Kinoshita T.** 2017. Blue light regulation of stomatal opening and the plasma membrane H⁺-ATPase. *Plant Physiology* **174**, 531–538.
- Islam MM, Munemasa S, Hossain MA, Nakamura Y, Mori IC, Murata Y.** 2010. Roles of AtTPC1, vacuolar two pore channel 1, in *Arabidopsis* stomatal closure. *Plant & Cell Physiology* **51**, 302–311.
- Islam MM, Tani C, Watanabe-Sugimoto M, Uraji M, Jahan MS, Masuda C, Nakamura Y, Mori IC, Murata Y.** 2009. Myrosinases, TGG1 and TGG2, redundantly function in ABA and MeJA signaling in *Arabidopsis* guard cells. *Plant & Cell Physiology* **50**, 1171–1175.
- Islam MM, Ye W, Matsushima D, Khokon MA, Munemasa S, Nakamura Y, Murata Y.** 2015. Inhibition by acrolein of light-induced stomatal opening through inhibition of inward-rectifying potassium channels in *Arabidopsis thaliana*. *Bioscience, Biotechnology, and Biochemistry* **79**, 59–62.
- Islam MM, Ye W, Matsushima D, Munemasa S, Okuma E, Nakamura Y, Biswas S, Mano J, Murata Y.** 2016. Reactive carbonyl species mediate ABA signaling in guard cells. *Plant & Cell Physiology* **57**, 2552–2563.
- Khokon MA, Jahan MS, Rahman T, Hossain MA, Muroyama D, Minami I, Munemasa S, Mori IC, Nakamura Y, Murata Y.** 2011. Allyl isothiocyanate (AITC) induces stomatal closure in *Arabidopsis*. *Plant, Cell & Environment* **34**, 1900–1906.
- Kinoshita T, Nishimura M, Shimazaki K.** 1995. Cytosolic concentration of Ca²⁺ regulates the plasma membrane H⁺-ATPase in guard cells of fava bean. *The Plant Cell* **7**, 1333–1342.
- Kinoshita T, Shimazaki K.** 2001. Analysis of the phosphorylation level in guard-cell plasma membrane H⁺-ATPase in response to fusicoccin. *Plant & Cell Physiology* **42**, 424–432.
- Klüsener B, Young JJ, Murata Y, Allen GJ, Mori IC, Hugouvieux V, Schroeder JI.** 2002. Convergence of calcium signaling pathways of pathogenic elicitors and abscisic acid in *Arabidopsis* guard cells. *Plant Physiology* **130**, 2152–2163.
- Kong D, Hu HC, Okuma E, et al.** 2016. L-Met activates *Arabidopsis* GLR Ca²⁺ channels upstream of ROS production and regulates stomatal movement. *Cell Reports* **17**, 2553–2561.
- Kwak JM, Murata Y, Baizabal-Aguirre VM, Merrill J, Wang M, Kemper A, Hawke SD, Tallman G, Schroeder JI.** 2001. Dominant negative guard cell K⁺ channel mutants reduce inward-rectifying K⁺ currents and light-induced stomatal opening in *Arabidopsis*. *Plant Physiology* **127**, 473–485.
- Laanemets K, Brandt B, Li J, Merilo E, Wang YF, Keshwani MM, Taylor SS, Kollist H, Schroeder JI.** 2013. Calcium-dependent and -independent stomatal signaling network and compensatory feedback control of stomatal opening via Ca²⁺ sensitivity priming. *Plant Physiology* **163**, 504–513.
- Lambrix V, Reichelt M, Mitchell-Olds T, Kliebenstein DJ, Gershenzon J.** 2001. The *Arabidopsis* epithiospecifier protein promotes the hydrolysis of glucosinolates to nitriles and influences *Trichoplusia ni* herbivory. *The Plant Cell* **13**, 2793–2807.
- Leckie CP, McAinsh MR, Allen GJ, Sanders D, Hetherington AM.** 1998. Abscisic acid-induced stomatal closure mediated by cyclic ADP-ribose. *Proceedings of the National Academy of Sciences, USA* **95**, 15837–15842.
- Lemtiri-Chlieh F, MacRobbie EA, Webb AA, Manison NF, Brownlee C, Skepper JN, Chen J, Prestwich GD, Brearley CA.** 2003. Inositol hexakisphosphate mobilizes an endomembrane store of calcium in guard cells. *Proceedings of the National Academy of Sciences, USA* **100**, 10091–10095.
- Marcus AI, Moore RC, Cyr RJ.** 2001. The role of microtubules in guard cell function. *Plant Physiology* **125**, 387–395.
- McAinsh MR, Webb A, Taylor JE, Hetherington AM.** 1995. Stimulus-induced oscillations in guard cell cytosolic free calcium. *The Plant Cell* **7**, 1207–1219.
- Melotto M, Underwood W, Koczan J, Nomura K, He SY.** 2006. Plant stomata function in innate immunity against bacterial invasion. *Cell* **126**, 969–980.
- Mori IC, Murata Y, Yang Y, et al.** 2006. CDPKs CPK6 and CPK3 function in ABA regulation of guard cell S-type anion- and Ca²⁺-permeable channels and stomatal closure. *PLoS Biology* **4**, e327.
- Munemasa S, Hossain MA, Nakamura Y, Mori IC, Murata Y.** 2011. The *Arabidopsis* calcium-dependent protein kinase, CPK6, functions as a positive regulator of methyl jasmonate signaling in guard cells. *Plant Physiology* **155**, 553–561.
- Munemasa S, Oda K, Watanabe-Sugimoto M, Nakamura Y, Shimoishi Y, Murata Y.** 2007. The coronatine-insensitive 1 mutation reveals the hormonal signaling interaction between abscisic acid and methyl jasmonate in *Arabidopsis* guard cells. Specific impairment of ion channel activation and second messenger production. *Plant Physiology* **143**, 1398–1407.
- Murata Y, Mori IC, Munemasa S.** 2015. Diverse stomatal signaling and the signal integration mechanism. *Annual Review of Plant Biology* **66**, 369–392.
- Murata Y, Pei ZM, Mori IC, Schroeder J.** 2001. Abscisic acid activation of plasma membrane Ca²⁺ channels in guard cells requires cytosolic NAD(P)H and is differentially disrupted upstream and downstream of reactive oxygen species production in *abi1-1* and *abi2-1* protein phosphatase 2C mutants. *The Plant Cell* **13**, 2513–2523.
- Nakamura Y, Miyoshi N.** 2010. Electrophiles in foods: the current status of isothiocyanates and their chemical biology. *Bioscience, Biotechnology, and Biochemistry* **74**, 242–255.
- Øverby A, Silihagen Bævre M, Bones A.** 2015. Disintegration of microtubules in *Arabidopsis thaliana* and bladder cancer cells by isothiocyanates. *Frontiers in Plant Science* **6**, 6.
- Pei ZM, Murata Y, Benning G, Thomine S, Klüsener B, Allen GJ, Grill E, Schroeder JI.** 2000. Calcium channels activated by hydrogen peroxide mediate abscisic acid signalling in guard cells. *Nature* **406**, 731–734.
- Peiter E, Maathuis FJ, Mills LN, Knight H, Pelloux J, Hetherington AM, Sanders D.** 2005. The vacuolar Ca²⁺-activated channel TPC1 regulates germination and stomatal movement. *Nature* **434**, 404–408.
- Roelfsema MRG, Hedrich R.** 2010. Making sense out of Ca²⁺ signals: their role in regulating stomatal movements. *Plant, Cell and Environment* **33**, 305–321.
- Salehin M, Li B, Tang M, Katz E, Song L, Ecker JR, Kliebenstein DJ, Estelle M.** 2019. Auxin-sensitive Aux/IAA proteins mediate drought tolerance in *Arabidopsis* by regulating glucosinolate levels. *Nature Communications* **10**, 4021.
- Sauer M, Paciorek T, Benková E, Friml J.** 2006. Immunocytochemical techniques for whole-mount in situ protein localization in plants. *Nature Protocols* **1**, 98–103.
- Schroeder JI, Hagiwara S.** 1989. Cytosolic calcium regulates ion channels in the plasma membrane of *Vicia faba* guard cells. *Nature* **338**, 427–430.
- Shimazaki K, Doi M, Assmann SM, Kinoshita T.** 2007. Light regulation of stomatal movement. *Annual Review of Plant Biology* **58**, 219–247.
- Siegel RS, Xue S, Murata Y, Yang Y, Nishimura N, Wang A, Schroeder JI.** 2009. Calcium elevation-dependent and attenuated resting calcium-dependent abscisic acid induction of stomatal closure and abscisic acid-induced enhancement of calcium sensitivities of S-type anion and inward-rectifying K channels in *Arabidopsis* guard cells. *The Plant Journal* **59**, 207–220.
- Sobahan MA, Akter N, Okuma E, Uraji M, Ye W, Mori IC, Nakamura Y, Murata Y.** 2015. Allyl isothiocyanate induces stomatal closure in *Vicia faba*. *Bioscience, Biotechnology, and Biochemistry* **79**, 1737–1742.
- Takahashi F, Suzuki T, Osakabe Y, Betsuyaku S, Kondo Y, Dohmae N, Fukuda H, Yamaguchi-Shinozaki K, Shinozaki K.** 2018. A small peptide modulates stomatal control via abscisic acid in long-distance signalling. *Nature* **556**, 235–238.

- Takahashi Y, Ebisu Y, Kinoshita T, Doi M, Okuma E, Murata Y, Shimazaki K.** 2013. bHLH transcription factors that facilitate K⁺ uptake during stomatal opening are repressed by abscisic acid through phosphorylation. *Science Signaling* **6**, ra48.
- Thor K, Peiter E.** 2014. Cytosolic calcium signals elicited by the pathogen-associated molecular pattern flg22 in stomatal guard cells are of an oscillatory nature. *New Phytologist* **204**, 873–881.
- Tian W, Hou C, Ren Z, et al.** 2019. A calmodulin-gated calcium channel links pathogen patterns to plant immunity. *Nature* **572**, 131–135.
- Wang Y, Chen ZH, Zhang B, Hills A, Blatt MR.** 2013a. PYR/PYL/RCAR abscisic acid receptors regulate K⁺ and Cl⁻ channels through reactive oxygen species-mediated activation of Ca²⁺ channels at the plasma membrane of intact *Arabidopsis* guard cells. *Plant Physiology* **163**, 566–577.
- Wang YF, Munemasa S, Nishimura N, et al.** 2013b. Identification of cyclic GMP-activated nonselective Ca²⁺-permeable cation channels and associated CNGC5 and CNGC6 genes in *Arabidopsis* guard cells. *Plant Physiology* **163**, 578–590.
- Xue S, Hu H, Ries A, Merilo E, Kollist H, Schroeder JI.** 2011. Central functions of bicarbonate in S-type anion channel activation and OST1 protein kinase in CO₂ signal transduction in guard cell. *The EMBO Journal* **30**, 1645–1658.
- Ye W, Adachi Y, Munemasa S, Nakamura Y, Mori IC, Murata Y.** 2015. Open stomata 1 kinase is essential for yeast elicitor-induced stomatal closure in *Arabidopsis*. *Plant & Cell Physiology* **56**, 1239–1248.
- Ye W, Murata Y.** 2016. Microbe associated molecular pattern signaling in guard cells. *Frontiers in Plant Science* **7**, 583.
- Ye W, Muroyama D, Munemasa S, Nakamura Y, Mori IC, Murata Y.** 2013. Calcium-dependent protein kinase CPK6 positively functions in induction by yeast elicitor of stomatal closure and inhibition by yeast elicitor of light-induced stomatal opening in *Arabidopsis*. *Plant Physiology* **163**, 591–599.
- Yin Y, Adachi Y, Ye W, Hayashi M, Nakamura Y, Kinoshita T, Mori IC, Murata Y.** 2013. Difference in abscisic acid perception mechanisms between closure induction and opening inhibition of stomata. *Plant Physiology* **163**, 600–610.
- Yoshida R, Mori IC, Kamizono N, Shichiri Y, Shimatani T, Miyata F, Honda K, Iwai S.** 2016. Glutamate functions in stomatal closure in *Arabidopsis* and fava bean. *Journal of Plant Research* **129**, 39–49.
- Yuan F, Yang H, Xue Y, et al.** 2014. OSCA1 mediates osmotic-stress-evoked Ca²⁺ increases vital for osmosensing in *Arabidopsis*. *Nature* **514**, 367–371.
- Zhang J, De-Oliveira-Ceciliato P, Takahashi Y, et al.** 2018. Insights into the molecular mechanisms of CO₂-mediated regulation of stomatal movements. *Current Biology* **28**, R1356–R1363.
- Zhang W, He SY, Assmann SM.** 2008. The plant innate immunity response in stomatal guard cells invokes G-protein-dependent ion channel regulation. *The Plant Journal* **56**, 984–996.
- Zhao Z, Zhang W, Stanley BA, Assmann SM.** 2008. Functional proteomics of *Arabidopsis thaliana* guard cells uncovers new stomatal signaling pathways. *The Plant Cell* **20**, 3210–3226.
- Zhu M, Zhu N, Song WY, Harmon AC, Assmann SM, Chen S.** 2014. Thiol-based redox proteins in abscisic acid and methyl jasmonate signaling in *Brassica napus* guard cells. *The Plant Journal* **78**, 491–515.