

指導教授氏名 Instructing Professor Commissioned Professor	指導役割 Role
印	研究全般の指導
印	
印	

## 学位論文要旨

## Dissertation Abstract

岡山大学大学院医歯薬学総合研究科

専攻分野 Department	機能再生・再建科学 歯科矯正学分野	身分 大学院生	氏名 Name	ワン 王 WANG	ズーイー 紫儀 ZIYI
論文題名 Title of Doctoral Dissertation	Bioinformatic analysis of mechano-sensitive genes and pathways in osteocyte under the mechanical loadings バイオインフォマティクス解析による骨細胞のメカノセンシティブ遺伝子と経路の解析				
論文内容の要旨 (2000字程度) Dissertation Abstract (approx. 800 words)					
<p><b>Background:</b> Review in <b>Chapter I</b> discussed the cell-cell communications among bone cells and those between the bone and other organs systems with an emphasis on the role of the energy metabolism. The intercellular network of cell-cell communications among osteocytes is mediated by gap junctions. Gap junctional intercellular communication (GJIC) is thought to play an essential role in the integration and synchronization of bone remodeling. To further understand the mechanism of bone development, it is of great importance to investigate the underlying mechanism of osteocyte differentiation-induced changes of GJIC capacity.</p> <p><b>Methods:</b> <b>Chapter II</b> emphasized on the mechano-sensitive genes to figure out a possible mechanism for the osteocyte differentiation-induced changes of GJIC capacity. Therefore, a bioinformatics analysis was applied to screen the key genes and pathways that are activated when different types of mechanical loading are applied to osteocytes. We retrieved the public mRNA expression datasets (series number of GSE62128 and GSE42874) from Gene Expression Omnibus database (GEO). High gravity-treated osteocytic MLO-Y4 cell-line samples from GSE62128 (Set1) and fluid flow-treated MLO-Y4 samples from GSE42874 (Set2) were employed. Functional enrichment was performed after the differentially expressed genes (DEGs) were identified. The common DEGs between Set1 and</p>					

論文内容の要旨（2000字程度） [Dissertation Abstract \(approx. 800 words\)](#)

Set2 were considered as key DEGs, and then the minimal nodes from all such key DEGs were used to construct a protein-protein interaction (PPI) network that linked most of the key DEGs. Several open source software programs were employed to process and analyze the original data. The bioinformatic results and the biological meaning were validated by *in vitro* experiments.

**Results:** Bioinformatic analysis in **Chapter II** demonstrated that high gravity and fluid flow induced opposite expression trends in the key DEGs. The hypoxia-related biological process and signaling pathway were the common functional enrichment terms among the DEGs from Set1, Set2, and the PPI network. The expression of almost all the key DEGs (*Pdk1*, *Ccng2*, *Eno2*, *Egln1*, *Higd1a*, *Slc5a3*, and *Mxi1*) was mechano-sensitive. *Eno2*, a gene related to adenosine triphosphate generation, was identified as the hub gene in the PPI network, and *Eno2* knockdown resulted in expression changes of some other key DEGs (*Pdk1*, *Mxi1*, and *Higd1a*).

**Conclusions:** A brief review in **Chapter III** elaborated on the changes of GJIC induced by hypoxia and several hypotheses were proposed for the future studies. Our findings based on the prediction of a bioinformatics analysis indicated that the hypoxia response might have an important role in the differential responses of osteocytes to the different types of mechanical force and the osteocyte differentiation-induced changes of GJIC capacity may be subject to the regulation of hypoxia signal pathway-mediated energy metabolism.