Supplementary Table 1. Characteristics of all POAF patients

Case No.	Age/Sex	LAd, mm	RCRI	Past history of PAF	Disease	Surgical procedure	Surgical time, min	Blood loss. mL	Days from surgery to onset of PAF	Days from surgery to discharge
Bisoprolol										
1	65/M	31	2	−	Esophageal cancer	Laparoscopic esophagectomy	640	110	1	22
2	90/M	41	2	−	Lung cancer	Lobectomy	257	130	1	15
3	72/M	38	2	−	Lung cancer	Lobectomy	173	60	1	11
4	73/M	40	3	−	Abdominal aortic aneurysm	Aortic replacement with Y graft	330	2450	1	21
Control										
1	76/M	30	2	+	Gastric cancer	Laparoscopic Gastrectomy	362	130	1	12
2	66/M	27	2	−	Lung cancer	Lobectomy	640	2425	3	28
3	73/F	35	2	+	Pancreatic cystic tumor	Distal Pancreatectomy	353	50	3	32

4	72/M	41	2	−	Lung cancer	Lobectomy	313	150	30	11
5	79/M	33	2	−	Lung cancer	Lobectomy	642	1375	3	12
6	69/M	28	2	−	Lung cancer	Lobectomy	385	375	8	11
7	82/F	33	3	+	Pancreatic head cancer	Pancreaticoduodenectomy	463	460	1	25
8	82/M	40	2	−	Lung cancer	Lobectomy	274	40	3	17
9	86/M	41	2	+	Lung cancer	Lobectomy	202	0	1	11
10	85/M	40	2	+	Rectal cancer	Rectal resection	341	50	1	19
11	74/M	35	2	+	Lung cancer	Lobectomy	258	170	1	12
12	80/M	36	2	−	Lung cancer	Lobectomy	185	50	0	35 POD in-hospital death due to pneumonia

POAF, perioperative atrial fibrillation; LAd, left atrial dimension; RCRI, revised cardiac risk index; PAF, paroxysmal atrial fibrillation; POD, postoperative day