

Supplementary Information for:

Effect of patient participation continuous nutritional counseling in gastric cancer patients who underwent gastrectomy

Nobuo Takata^{1,2}, Satoru Kikuchi^{1*}, Shinji Kuroda¹, Shunsuke Tanabe¹, Naoaki Maeda¹, Kazuhiro Noma¹,
Ayako Takahashi³, Yuzo Umeda¹, Kenichi Shikata^{3,4}, Kazuhide Ozaki²
and Toshiyoshi Fujiwara¹



¹Department of Gastroenterological Surgery, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Okayama 700-8558, Japan; ²Department of Gastroenterological Surgery, Kochi Health Sciences Center, Kochi 781-8555, Japan, ³Department of Clinical nutrition, Okayama University Hospital, Okayama 700-8558, Japan, ⁴Center for Innovative Clinical Medicine, Okayama University Hospital, Okayama 700-8558, Japan

Supplementary Fig. S1

Perioperative and post-discharge nutritional intervention protocols in the control and CNC groups

Supplementary Fig. S2


Rate of development of sarcopenia at POM12 between patients without preoperative sarcopenia in the control and CNC groups.



Supplementary Fig. S1

Perioperative and post-discharge nutritional interventions protocols in the control and CNC groups.

CNC, continuous nutritional counseling



Supplementary Fig. S2

Rate of development of sarcopenia at POM12 between patients without preoperative sarcopenia in the control and CNC groups.

CNC, continuous nutritional counseling