



Case Report

Korean J Anesthesiol 2024;77(5):565–569

<https://doi.org/10.4097/kja.24232>

pISSN 2005–6419 • eISSN 2005–7563

Received: April 9, 2024

Revised: May 15, 2024 (1st); June 3, 2024 (2nd)

Accepted: June 4, 2024

Corresponding author:

Daisuke Sugiyama, M.D., Ph.D.

Department of Anesthesiology, Kameda Medical Center, 929 Higashi-cho, Kamogawa City, Chiba, Japan

Tel: +81-470-92-2211

Fax: +81-470-99-7798

Email: sugiyama.daisuke@kameda.jp

ORCID: <https://orcid.org/0000-0002-4848-2059>

Previous presentation in conferences:

This work was previously presented at the 40th ESRA Annual Congress, 6th World Congress on Regional Anesthesia & Pain Medicine, held in September 6, 2023 in Paris, France.



© The Korean Society of Anesthesiologists, 2024

© This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Utilization of the pericapsular nerve group block in preoperative rehabilitation of patients with femoral neck fractures -a case series-

Zhuan Jin^{1,2}, Daisuke Sugiyama¹, Fumiya Higo³, Takahiro Hirata³, Osamu Kobayashi¹, Hiroshi Morimatsu², Kenichi Ueda^{1,4}

¹Department of Anesthesiology, Kameda Medical Center, Chiba, ²Department of Anesthesiology and Resuscitology, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Okayama, ³Department of Rehabilitation, Kameda Medical Center, Chiba, Japan, ⁴Department of Anesthesia, University of Iowa Hospitals and Clinics, Iowa City, IA, USA

Background: Elderly patients with femoral neck fractures, particularly those with severe comorbidities or living in regions with limited medical resources, may experience delays in surgical treatment. Although the benefits of preoperative rehabilitation (prehabilitation) in hip arthroplasty have been reported, pain management remains a challenge. The pericapsular nerve group (PENG) block, known for its exceptional analgesic effect and motor function preservation, may be a promising intervention during prehabilitation in these patients.

Case: We enrolled ten patients with Garden classification 3–4 femoral neck fractures scheduled for hip arthroplasty. After receiving a PENG block with 20 ml of 0.375% ropivacaine, all patients underwent initial prehabilitation sessions comprising 9 mobility levels, ranging from bed-sitting to walking. One patient was excluded due to experiencing high blood pressure during prehabilitation. Six of the nine remaining patients (66.7%) were successfully transferred from bed to wheelchair.

Conclusions: The PENG block enhanced prehabilitation for patients with femoral neck fractures undergoing hip arthroplasty.

Keywords: Conduction anesthesia; Femoral neck fractures; Hip fractures; Nerve block; Prehabilitation; Preoperative exercise; Preoperative rehabilitation; Regional anesthesia.

Timely surgical intervention for femoral neck fractures can be challenging, particularly in elderly patients with severe comorbidities and in regions with limited medical resources [1]. Preoperative rehabilitation (prehabilitation) has been shown to be effective at enhancing mobilization, preserving muscle strength, and improving postoperative outcomes for patients with femoral neck fractures scheduled for hip arthroplasty [2,3]; however, significant pain during prehabilitation remains a barrier. The pericapsular nerve group (PENG) block, which targets the articular sensory nerve branches of the anterior capsule of the femoral neck, provides excellent postoperative analgesia while preserving motor function [4–6]. This study aims to investigate the feasibility of utilizing the PENG block in prehabilitation for patients with femoral neck fractures scheduled for hip arthroplasty and to offer insights for a future large-scale randomized controlled trial.

Case Report

This study was approved by the Kameda Medical Center Clinical Research Review Board (approval number: 21-093) and registered in the Japan Registry of Clinical Trials (registration number: JRCT1031220004) prior to patient enrollment.

A total of 16 adult patients diagnosed with Garden classification 3 or 4 femoral neck fractures and scheduled for hip arthroplasty at Kameda Medical Center in Japan between April 2022 and July 2022, were assessed. Patients were excluded if they met any of the following criteria: 1) dementia, defined as scoring less than 23 points on the Mini-Mental Status Examination; 2) known allergies to local anesthetics; 3) a history of hip arthroplasty at the same extremity; 4) an infection at the site where the PENG block would be performed; 5) those bedridden prior to the injury; and 6) those determined to be inappropriate for participation at the discretion of the orthopedic surgeon. After excluding three patients with dementia, one patient who refused participation, one patient who was bedridden prior to the injury, and one patient who was assessed as inappropriate for participation due to a pathologic fracture, a total of ten eligible patients were enrolled in the study. Written informed consent was obtained from all the enrolled patients.

The study did not interfere with surgical treatment. The surgical procedure was scheduled to be performed as early as possible following current guidelines [7]. All patients were allowed preoperative mobilization with no weight bearing on the injured lower extremity, a practice that has been validated to be safe and effective in this patient cohort: patients with Garden 3–4 femoral neck fractures who are scheduled for a hip arthroplasty [3]. Additionally, each patient received standard preoperative pain management including prescriptions for acetaminophen, non-steroidal anti-inflammatory drugs, and tramadol.

The PENG block was performed at least 1 h before the initial prehabilitation session. Following local anesthesia, we performed a PENG block at the patient's bedside using the technique described by Giron-Arango et al. [4], with 20 ml of 0.375% ropivacaine solution. The procedure was performed using a handheld portable ultrasound device (Vscan Air, GE Healthcare) and a 22-gauge 80-mm needle (Stimuplex Ultra, B. Braun).

After PENG block administration, the initial prehabilitation session was conducted by a single qualified physical therapist. The sessions incorporated a prehabilitation program aimed at enhancing mobilization and preserving muscle strength, designed according to the ICU Mobility Scale (IMS) [8] and the Cumulated Ambulation Score [9,10]. The program comprises nine mobility levels, with each subsequent level being progressively more demanding (Table 1). Prehabilitation was performed following this sequence of mobility levels until the patient was unable to perform the exercises, requested to stop, or the discontinuation criteria were met [11]. The same prehabilitation protocol was provided to all patients once daily until surgery; however, only the initial session was conducted under the PENG block. Subsequent sessions were conducted using standard pain management.

Table 2 presents the baseline characteristics of the ten patients enrolled in this study. The outcomes of the prehabilitation program during the initial prehabilitation are shown in Fig. 1. Due to experiencing a high blood pressure exceeding 180/120 mmHg, one patient could not complete the initial prehabilitation [11]. The median Numerical Rating Scale pain score during prehabilitation was 6.0 (5.0, 7.0) points.

Four of the nine patients underwent surgery the day after their initial prehabilitation session, while the remaining five had a median wait time of 3.0 (2.0, 4.0) days before surgery. Regarding postoperative outcomes, patients achieved a median Functional Independence Measure (FIM motor) score of 80.0 (66.0, 83.0).

Table 1. The Prehabilitation Program

Level	Item	Explanation
1	Bed-sitting	Sitting up on the bed with the assistance of a single physical therapist.
2	Edge-sitting	Sitting on the edge of the bed with the assistance of a single physical therapist.
3	Stand-up	Standing up from the bed with the assistance of a single physical therapist.
4	Wheelchair-transfer	Transferring to wheelchair with the assistance of a single physical therapist.
5	Marching	Stepping at the bedside with the assistance of a single physical therapist.
6	Walk-with-two	Walking at least 5 meters with the assistance of a single physical therapist and either another physical therapist or a ward nurse.
7	Walk-with-one	Walking at least 5 meters with the assistance of a single physical therapist.
8	Walk-with-device	Walking at least 5 meters with a walking-device.
9	Walk-alone	Walking independently at least 5 meters without a walking-device.

Prehabilitation: preoperative rehabilitation.

Table 2. Patient Baseline Characteristics

Characteristic	Value
Age (yr)	75.5 ± 7.1
Female sex	10 (100.0)
Asian race	10 (100.0)
BMI	19.5 ± 3.6
ASA-PS	
I	0 (0.0)
II	6 (60.0)
III	4 (40.0)
Garden classification	
III: complete fracture, incompletely displaced	0 (0)
IV: complete fracture, completely displaced	10 (100)
Walking ability before fracture	
Independent	9 (90.0)
Walking-device use	1 (10.0)
Preadmission residence	
Home	10 (100.0)
Nursing home or other locations	0 (0.0)
FIM motor score*	26.5 (22.3, 40.3)
Barthel Index [†]	30.0 (18.8, 38.8)

Values are presented as mean ± SD, number (%) or median (Q1, Q3).

*The FIM motor score comprises 13 activities of daily living (ADL) items. Each item is scored on a scale of 1–7. The total scores range from 13 to 91, with higher scores indicating greater independence. [†]The Barthel Index measures 10 ADL items. The total scores range from 0 to 100, with higher scores indicating greater independence. BMI: body mass index, ASA-PS: American Society of Anesthesiologists physical status classification, FIM: Functional Independence Measure.

points at discharge [12]. Three out of the nine patients (33.3%) were discharged to rehabilitation hospitals and nursing homes, while six (66.7%) were discharged directly to home. No motor blockade or other adverse events associated with the PENG block were observed.

Discussion

In this study, we investigated the utilization of the PENG block in prehabilitation for patients with femoral neck fractures scheduled for hip arthroplasty. In the preoperative phase, patients often experience intolerable pain, which impedes basic daily activities such as eating and toileting and adversely affects muscle strength maintenance, thereby influencing postoperative outcomes [13]. Incorporating the PENG block in preoperative care resulted in enhanced performance in prehabilitation compared to conventional approaches, with more than half of the patients successfully transferred from bed to wheelchair.

While current guidelines recommend prompt surgical intervention for femoral neck fractures [7], a significant number of elderly patients, particularly those with severe comorbidities, encounter difficulties in adhering to these guidelines. Factors such as the requirement to discontinue anticoagulants or antiplatelets or the need for additional examinations can contribute to delays in surgery for these individuals. As demonstrated in this study, the integration of prehabilitation with the PENG block presents a promising treatment approach, particularly beneficial for individuals experiencing delays in surgical intervention.




Fig. 1. Prehabilitation program outcomes in the initial prehabilitation session. One patient who was unable to participate in prehabilitation due to high blood pressure was excluded from the outcome analysis.

Furthermore, although many patients with femoral neck fractures cannot fully regain the ability to perform activities of daily living during their hospital stay, often requiring discharge to rehabilitation hospitals or nursing homes after hospitalization for further care, the patients in this study exhibited enhanced postoperative outcomes, characterized by a high FIM motor score, and a large proportion were discharged directly to home. Therefore, receiving a PENG block only once during the initial prehabilitation session can have a substantial effect on prehabilitation and preoperative mobilization, with potential promising effects on postoperative outcomes. However, this needs to be further investigated in future studies.

Our study had several limitations. First, the case-series design may introduce potential biases, thereby affecting the generalizability of the findings. Moreover, the use of a single-shot PENG block only in the initial prehabilitation rather than catheter placement for continuous infusion limited our capacity to assess the long-term impact on postoperative outcomes.

Based on the results of this study, we are currently conducting a randomized controlled trial to provide further evidence of the effects of the PENG block in this context.

Funding

None.

Conflicts of Interest

No potential conflict of interest relevant to this article was reported.

Data Availability

The datasets generated during and/or analyzed during the current study are available from the corresponding author upon reasonable request.

Author Contributions

Zhuan Jin (Conceptualization; Data curation; Formal analysis; Investigation; Methodology; Project administration; Visualization; Writing – original draft; Writing – review & editing)
 Daisuke Sugiyama (Conceptualization; Data curation; Investigation; Methodology; Project administration; Resources; Supervision; Validation; Writing – review & editing)
 Fumiya Higo (Conceptualization; Data curation; Investigation;

Methodology; Project administration; Writing – review & editing)
 Takahiro Hirata (Conceptualization; Data curation; Investigation; Methodology; Project administration; Writing – review & editing)
 Osamu Kobayashi (Conceptualization; Supervision; Validation; Writing – review & editing)
 Hiroshi Morimatsu (Conceptualization; Supervision; Validation; Writing – review & editing)
 Kenichi Ueda (Conceptualization; Methodology; Project administration; Resources; Supervision; Validation; Writing – review & editing)

ORCID

Zhuan Jin, <https://orcid.org/0009-0008-0338-3664>
 Daisuke Sugiyama, <https://orcid.org/0000-0002-4848-2059>
 Fumiya Higo, <https://orcid.org/0009-0003-4697-6949>
 Takahiro Hirata, <https://orcid.org/0009-0001-7919-3698>
 Osamu Kobayashi, <https://orcid.org/0009-0001-2968-6814>
 Hiroshi Morimatsu, <https://orcid.org/0000-0002-6744-0508>
 Kenichi Ueda, <https://orcid.org/0000-0002-4559-4581>

References

1. Vidán MT, Sánchez E, Gracia Y, Marañón E, Vaquero J, Serra JA. Causes and effects of surgical delay in patients with hip fracture: a cohort study. *Ann Intern Med* 2011; 155: 226-33.
2. Sawaguchi A, Momosaki R, Hasebe K, Chono M, Kasuga S, Abo M. Effectiveness of preoperative physical therapy for older patients with hip fracture. *Geriatr Gerontol Int* 2018; 18: 1003-8.
3. Liu J, Chen L, Long C, Zhang X, Gao F, Duan X, et al. Effect of a preoperative mobilization program on perioperative complications and function recovery in older adults with femoral neck fracture. *Geriatr Nurs* 2022; 44: 69-75.
4. Girón-Arango L, Peng PW, Chin KJ, Brull R, Perlas A. Pericapsular nerve group (PENG) block for hip fracture. *Reg Anesth Pain Med* 2018; 43: 859-63.
5. Wang Y, Wen H, Wang M, Lu M. The efficiency of ultrasound-guided pericapsular nerve group block for pain management after hip surgery: a meta-analysis. *Pain Ther* 2023; 12: 81-92.
6. Huda AU, Ghafoor H. The use of pericapsular nerve group (PENG) block in hip surgeries is associated with a reduction in opioid consumption, less motor block, and better patient satisfaction: a meta-analysis. *Cureus* 2022; 14: e28872.
7. O'Connor MI, Switzer JA. AAOS clinical practice guideline summary: management of hip fractures in older adults. *J Am Acad Orthop Surg* 2022; 30: e1291-6.

8. Hodgson C, Needham D, Haines K, Bailey M, Ward A, Harrold M, et al. Feasibility and inter-rater reliability of the ICU Mobility Scale. *Heart Lung* 2014; 43: 19-24. Erratum in: *Heart Lung* 2014; 43: 388.
9. Parker MJ, Palmer CR. A new mobility score for predicting mortality after hip fracture. *J Bone Joint Surg Br* 1993; 75: 797-8.
10. McDonough CM, Harris-Hayes M, Kristensen MT, Overgaard JA, Herring TB, Kenny AM, et al. Physical therapy management of older adults with hip fracture. *J Orthop Sports Phys Ther* 2021; 51: CPG1-81.
11. Fletcher GF, Ades PA, Kligfield P, Arena R, Balady GJ, Bittner VA, et al. Exercise standards for testing and training: a scientific statement from the American Heart Association. *Circulation* 2013; 128: 873-934.
12. Keith RA, Granger CV, Hamilton BB, Sherwin FS. The functional independence measure: a new tool for rehabilitation. *Adv Clin Rehabil* 1987; 1: 6-18.
13. Carpintero P, Caeiro JR, Carpintero R, Morales A, Silva S, Mesa M. Complications of hip fractures: a review. *World J Orthop* 2014; 5: 402-11.